

# Potentially Important Food Plants of Vietnam



**FOOD PLANT  
SOLUTIONS  
ROTARIAN ACTION GROUP**

*Solutions to Malnutrition  
and Food Security*



A Project of the Rotary Club of Devonport North,  
District 9830 & Food Plants International

[www.foodplantsolutions.org](http://www.foodplantsolutions.org)



# Potentially Important Food Plants of Vietnam

## **Dedication**

This book is dedicated to the 3 billion hard working farmers and families around the world who cultivate these, and other, food plants for their own subsistence, and who help conserve them in their rich diversity for other people to enjoy.

## **Preface**

This guide is based on information from the Food Plants International (FPI) database developed by Tasmanian agricultural scientist Bruce French. The source material and guidance for the preparation of the book has been made possible through the support of Food Plants International, the Rotary Clubs of District 9830, particularly the Rotary Club of Devonport North who founded Food Plant Solutions, (previously the Learn♦Grow project), and many volunteers who have assisted in various ways.

The selection of plants included in this guide has been developed by Kirk Schmidt (Rotary Club of Freedom, California, U.S.A., District 5170), working in a voluntary capacity using the selection criteria developed by Food Plant Solutions. These selection criteria focus on the local plants from each of the main food groups with the highest levels of nutrients important to human nutrition and alleviation of malnutrition. It is intended as a **Draft Guide only** to indicate some important food plants that serve as examples for this purpose. Other important nutritious plants may be equally useful, and it is recommended that the FPI database be used to source information on the full range of plants known to occur in Vietnam. This guide has been developed with the best intention to create interest and improve understanding of the important local food plants of Vietnam, and on the understanding that it will be further edited and augmented by local specialists with appropriate knowledge and understanding of local food plants.

Food Plant Solutions was initiated by the Rotary Club of Devonport North to assist in creating awareness of the edible plant database developed by Food Plants International, and its potential in addressing malnutrition and food security in any country of the world. In June 2007, Food Plant Solutions was established as a project of Rotary District 9830, the Rotary Club of Devonport North and Food Plants International. The primary objective of the project is to increase awareness and understanding of the vast food resource that exists in the form of local plants, well adapted to the prevailing conditions where they naturally occur, and how this resource may be used to address hunger, malnutrition and food security. For more information, visit the website [www.foodplantsolutions.org](http://www.foodplantsolutions.org). More detailed or specific information on plants, including references to material by other authors, is available on DVD on request.

**Disclaimer:** This Field Guide has been produced using information from the “Edible Plants of the World” database compiled by Bruce French of Food Plants International. Although great care has been taken by Food Plants International and Food Plant Solutions, neither organisation, or the people involved in the compilation of the database or this Field Guide:

- makes any expressed or implied representation as to the accuracy of the information contained in the database or the Field Guide, and cannot be held legally responsible or accept liability for any errors or omissions
- can be held responsible for claims arising from the mistaken identity of plants or their inappropriate use
- assume responsibility for sickness, death or other harmful effects resulting from eating or using any plant described in the database or this Field Guide

Always be sure you have the correct plant, and undertake proper preparation methods, by consulting with specialist scientists or local users of the plant. The Food Plants International database, from which the information in this Field Guide is drawn, is a work in progress and is regularly being amended and updated.

# Contents

- INTRODUCTION ..... 1**
- STARCHY STAPLES ..... 11**
- LEGUMES..... 20**
- LEAFY GREENS ..... 26**
- FRUIT ..... 32**
- VEGETABLES..... 43**
- NUTS, SEEDS, HERBS AND OTHER FOODS ..... 51**
- NUTRITIONAL VALUES OF FOOD PLANTS BY PLANT FAMILY..... 56**



## ***Introduction***

This book is designed as a simple introduction to the more common food plants of Vietnam. It is hoped people will take greater pride and interest in these plants and become confident and informed about how to grow and use them. Many of the local food plants that occur in every country are very good quality foods. Unfortunately, people often reject traditional food plants and grow more of the introduced vegetables, such as ballhead cabbage. These do not have the same food value as many traditional, tropical, dark green, leafy vegetables.

### **Growing food**

Growing food to feed a family is, without doubt, one of the most important things anyone can do. The more interest you take in your garden and the more you learn about plants and how to grow them well, the more interesting and fun food gardening becomes.

### **A country with very special plants**

The local food plants of most countries have not been promoted and highlighted in the way they deserve. Visiting a local food market will quickly show what a rich variety of food plants can be grown in this country. Good information about these plants is often still in the minds and experience of local farmers, and has not been written down in books. This can make it hard for the next generation of young people to find out how to grow them.

In many countries, some of the traditional food plants are only harvested from the wild and others are only known in small areas. Others have hundreds of varieties and are the main food for people in different regions. Information on all these plants, their food value and the pest and diseases that damage them is available in the Food Plants International database.

### **Getting to know plants**

People who spend time in gardens and with their food plants get to know them very well. It is a good idea to learn from someone who grows plants well. Each plant grows best in certain conditions and there are often special techniques in getting it to grow well. For example, sweet potato will not form tubers if the soil is too wet, but it may still grow lots of green leaves. Taro will grow in light shade, but sweet potato will not. Ginger can grow in fairly heavy shade. Pruning the tips of betel leaf or pepper vines will cause more side branches to grow and therefore, produce more fruit. Stored yam tubers need special treatment if you want them to put out shoots early. There are lots of unique things about every plant and learning about these helps a good gardener produce more food.


## Naming of plants

Many food plants have local names, as well as a common English name. Every type of plant also has its own scientific name. Although the scientific name might not be widely recognised, this is the link by which people in different countries and with different languages can recognise the same plant. We know that many plants are grown in many different countries, but relying on local or common names, we might not recognise the same plant grown in different places. By using scientific names to accurately identify plants, we can get useful information from people in other countries. Wherever possible, plants in this book are named by their common English name and their scientific name.

## Local food plants are often very good

People sometimes think that local food plants are not very special and that any food plant that is new or comes from another country must be a lot better. This is often not true. Many of the newer or introduced food plants, such as the round or ballhead cabbages, have very little food value. Many traditional tropical green, leafy vegetables and ferns have 10 times or more food value as ballhead cabbage or lettuce. It is important to find out more information about the food value of different foods if we want to eat well. Citrus fruit, such as lemons and oranges, are often grown for vitamin C that helps keep people healthy. These fruits do not grow well in the tropics - the common guava fruit has three times as much vitamin C and is loved by children. This is just one example that there are often much better choices of local foods with higher levels of important nutrients.

Our bodies need a variety of food plants to enable us to grow, stay healthy and have enough energy to work. Different foods are needed to provide energy, protein, vitamins and minerals. The following diagram highlights the iron content value of some traditional edible, tropical plant leaves, compared with cabbage. Iron is a nutrient that is very important for our bodies and especially our blood. People who are short of iron become anaemic and lack energy.



The relative iron content of some edible leaves

## **A healthy balanced diet**

Good nutrition, or eating a healthy balanced diet, is really very simple. If people eat a wide range of food plants, their bodies will normally get a balanced amount of all the different nutrients they require. If a nutrient is lacking in one food plant, then they are likely to get it from another plant if they are eating a range of food plants. For this reason, everybody should eat a range of different food plants every day. The food group that is especially important for young people is the dark green leaves. Everyone should eat a good serving of dark green leaves every day. They have many vitamins and minerals, as well as protein. There are many spices or flavouring plants that can improve the taste of foods, but taste should be considered separately from food value.

## **Learning to cook well**

Even though some nutrients in food can lose some of their value during cooking, it is normally much safer to cook all food plants, at least for a short time. Bacteria, which cause diarrhoea, can occur in gardens and on food plants. These are killed during cooking. Many plants in the tropics develop cyanide, a chemical that makes them bitter and poisonous. This happens often with cassava (tapioca, manioc) and beans, but can also occur in many other plants. Boiling the food for two minutes normally destroys cyanide and makes the food safe to eat. Some of the nutrients our bodies need (such as vitamin A for good eyesight) only become available when food is cooked in oil.

## **Learning to grow “wild” food plants**

Many plants grow wild in the bush and are not cultivated by people. We can normally find someone who has taken an interest in them and has learned to grow them. This may be people from a different language group. It may be that in their area they have found better types than the ones that simply grow wild.

## **Saving better types of plants**

If we simply allow plants to grow from seed, the improvements that have been made in finding sweeter or better types may get lost. Some fruit trees are like this and the fruit produced may not be sweet at all. It is often necessary to take cuttings from a tree to be sure the new plant is exactly the same as the old one. If the plants won't easily grow from cuttings simply by sticking a piece of the branch in the ground, there are other ways of helping these plants to form roots and start to grow. One good way is to make a small cut in the bark of a young branch and then wrap soil around the cut and cover it with plastic. With plants like guava, new roots will start to grow from this cut and grow into the soil wrapped around the branch. It can then be cut off and planted. This is called air-layering. A similar method is used with the roots of breadfruit. A shallow root is uncovered and a small cut made from which a new sucker will start to grow. This can be cut off and replanted.

## **Growing from cuttings and suckers**

Many food plants are grown from cuttings and suckers. This is very important, as it allows all the different kinds of yams, taros, bananas, sweet potato and sugarcane to be continually grown and ensures the varieties are preserved. Each plant has its own special propagation method. It is important to use healthy planting material, as diseases can be spread in planting material.

## **Saving seed**

Some food plants are grown from seed. Sometimes this is very easy as the seeds are large, store well, grow easily and grow the same as the original plant. It is more difficult with other plants. Many large fleshy seeds, such as breadfruit, need to be planted while still fresh as they do not store easily. Other seeds do not “breed true” or do not grow into new plants that are the same as the original plants. For example, the fruit may not be as large or sweet or have the same colour or taste. With many of these plants, it may be necessary to find ways of growing them from cuttings or other methods such as grafting. Some plants “inbreed” and get smaller or poorer. This happens when a plant self-pollinates or receives pollen from a close relative. Corn grown in small plots normally does this and the plants grown from seed grown in this situation get smaller and smaller each year. The seed needs to be saved from several different plants with different history and then mixed together before sowing. All the seeds on one cob are related and will inbreed. Some seeds develop a hard seed coat and need to be scratched, soaked in water, or even put into hot water, before they will start to grow. Saving local seeds is often a good idea as they are already adapted to local conditions. For example, seed saved from pumpkins grown locally will produce plants with less pest and disease damage than those grown from imported seed. *If you can't get seeds or planting material from local gardens – it is probably not a suitable local plant!*

## **Growing a garden of mixed plants**

In nature, one variety of one plant never grows alone. There are always lots of different plants of different kinds and sizes, all growing together. Anyone who has ever walked into a tropical jungle will know this very well. The reason people all over the world want to save the rainforest is because it has so many different kinds of plants all growing together. Growing plants in a food garden in a way similar to how they grow in nature, as a mixed group of plants, is very good agriculture. Mixing plants in a garden usually gives more reliable food production, as any disease from one plant will wash off in the rain onto a different plant, where it cannot survive. Small plants fill the gaps and reduce the need for weeding.

## **Different types of plants for food security**

There is another reason for growing a range of food plants in a local garden or around a village. If something goes wrong, like extreme insect damage to plants, some disease occurring in the garden, or a poor growing season, some plants will be more damaged than others. With a variety of plants, there will still be some food to eat

until the other plants recover and grow again. Also, a wide variety of plants will mean that different ones will be maturing at different times, which helps ensure a continuous supply of food. There are shrubs that can be planted as edible hedges around houses, and fruit and nut trees that need to be planted as a gift for your children, several years before they will be able to enjoy them. Some nuts can be stored and eaten when other foods are not available. Most yams will store well for a few months.

### **Looking after the soil**

Gardeners in traditional tropical agriculture usually move their gardens often by shifting to a new piece of land. There are usually three reasons for this:

- In the tropical lowlands, weeds can become a very big problem. There are usually a lot fewer weeds in the first year or two after clearing and burning the land, but weeds increase in the following years.
- Some of the nutrients in the soil are used each year and the soil becomes poorer and plants do not grow as well. There are ways of reducing this loss of nutrients.
- Very small worms called nematodes build up in the soil after a few years and get into the roots, especially of annual vegetable plants, and stop their roots working properly. For example, root knot nematode will cause the roots of plants like tomatoes and beans to become twisted resulting in poor growth of the plant.

### **Building up the soil**

When a new garden has been cleared, it has lots of leaf mulch and other old plant material. This provides plant nutrients for new plants to grow. There is a simple rule for growing plants and improving the soil - "If it has lived once, it can live again." Any old plant material can provide nutrients for new plants to grow, but it must be allowed to rot into mulch or compost for this to happen. If this plant material is burnt, some nutrients, especially phosphorus and potassium ("potash"), get left behind in the ashes for new plants to use, although it also allows these important nutrients to be lost by being washed away by rain. But with burning other important nutrients, such as nitrogen and sulphur, get lost in the smoke and disappear from the garden and soil. These last two plant nutrients are especially important for growing green leaves and when their levels are low, plants grow small or pale green. When nitrogen is lacking, the old leaves of the plant go pale and fall off early, and when sulphur is lacking, the young leaves go pale. Wherever possible, old plant material should be covered with some soil to allow it to rot down and not simply dry out or get burnt.

### **Poor soils where crops won't grow**

When soils are very acid (or sour), plants cannot get the necessary nutrients. Natural chemicals in the soil that are toxic to plants when present at higher levels become soluble, get into plants, and stop them growing. Adding limestone to these soils can improve them. Using compost will not make them less acid, but will keep the plant nutrients in the soil in a more readily available form that plants can use.

### **Soil nutrients**

Plants need 16 different kinds of plant food or nutrients in different amounts to grow properly. A plant that has already been growing will have these nutrients in them and probably even have them in a balanced amount. That is why composting old plant material is so important. Plants usually show some signs or symptoms if any of these nutrients is running out.

One of the most common and important nutrients for plant growth is nitrogen, which actually comes from the air, but gets into plants through the soil. When plants are short of nitrogen, their older leaves often become yellow or pale. When grass family plants, like sugarcane and corn, are short of nitrogen, the centre of the oldest (lowest) leaves starts to develop a dry or dead V-shape. The plant cannot find enough nitrogen in the soil so it gets it from an old leaf to grow a new leaf. This causes the old leaf to die, forming a characteristic V-shape in the centre of the leaf. The plant does not get any bigger as an old leaf dies each time a new leaf is produced. Village farmers often walk through grassland before they clear it for gardens, looking to see if the grass leaves are dry and dead, because they know gardens on this soil won't grow well. It is necessary to use compost or legumes (such as beans) to put nitrogen back into the soil. Growing plants from the bean family (legumes) is the most efficient way to increase the level of nitrogen in the soil.

Corn is a good plant for indicating which nutrients are running short in the soil. If the older leaves go dry along the edges, the soil is running out of potash. If leaves that are normally green develop a bluish colour, the soil is short of phosphorus. Generally, leafy crops need lots of nitrogen, and root crops need lots of potash.

### **Making compost**

Compost is old plant material that has been allowed to rot down into a fine, sweet smelling mulch that is full of nutrients that can be put back on the soil to grow new plants. Making good compost is very simple. A simple heap of plant material can be made in the corner of a garden or near a house. The composting process is carried out by small bacteria that live in the soil and feed on decaying plants. They break down old plant material into compost. These bacteria are living, so they need air, water and food. A good compost heap must have air, so don't cover it with plastic or put it in a container. This makes a foul smelling compost, as different bacteria that don't need air turn it into an acid mixture that preserves it. Good compost must have

moisture, so keep the heap damp, but not too wet. The compost bacteria like a balanced diet, which means that both green material and dried material is needed to balance the carbon and nitrogen in the compost pile. If the compost material gets too dry and brown, it will not break down, and if it gets too green, it will go slimy. Using a little bit of compost from an old heap will make sure the right bacteria are there to start the whole process off. As soon as the plant material is broken down to a fine mulch it can be put onto the garden. It is best if it is dug in, but if it is regularly put onto the surface of the garden, worms will mix it into the soil.



## **Pests**



There are a large number of insects that enjoy sharing our food with us! We should not try to kill all these insects as they have an important role to play in keeping everything in nature in balance. What we need to do is to learn to manage these insects so we can all get some food to eat! Some insects are attracted to lights, and if the garden is near village lights some insects can cause a lot of damage. If large areas of one particular crop are planted, insects can breed more quickly and cause a lot of damage. As an example, insects called armyworms can breed up in large numbers on the shade trees of cacao and then move “like an army” into gardens. Some insects are large and breed slowly and can be picked off and removed. The large, green grubs with pointy tips that hide under taro leaves are best controlled by simply picking them off. Some insects, like taro beetles, can be a serious problem, but the young curl grubs of this insect are tasty if you catch and cook them. Some insects do not like sunlight. The very small moth than damages banana fruit is like this. Simply pulling off the leafy bracts over the banana fruit reduces the damage, as this lets sunlight in and the insect flies away. The best rule for reducing pest damage is to grow healthy plants, as they suffer less damage.


## **Diseases**

The living organisms that cause disease are much smaller than insects. These disease organisms can often only be seen with a microscope. There are three main kinds of disease organisms - fungi, bacteria and viruses. Fungi are like the mushrooms we eat, only very much smaller. They usually make distinct dry spots on leaves and other plant parts. Fungi have spores that often blow in the wind. Bacteria are often smaller and live in damp places. They usually make plants go soft and squashy, and they may cause a smell. Bacteria are mostly spread with rain and in water. Viruses are very, very small and usually make irregular stripes and patterns on leaves and other plant parts. Viruses usually spread in planting material or in the mouths of small sucking insects. One common fungus disease on sweet potato causes the leaves to become wrinkled and twisted. It usually gets worse in old gardens and where soils are running out of nutrients. It doesn't affect all kinds of sweet potato to the same extent. The answer is not to stop the disease, but to improve the soil. The general rule is that healthy plants that are growing well will suffer less damage from disease.

## Food value charts for a selection of plants from Vietnam







**Note regarding plant selection:** In compiling these field guides, we acknowledge that some staple foods and commercial crops which are grown widely in the target country may be omitted. Such foods are often in the starchy staple category (e.g. rice). This does not mean that they are not useful, but merely reflects a desire to concentrate on plants that are less well known and/or underutilised.

## *Starchy staples*

**English:** Cassava

**Local:** Sǎn

**Scientific name:** *Manihot esculenta*

**Plant family:** EUPHORBIACEAE

**Description:** A plant which can re-grow year after year from the thickened roots. It has several stems. The stems are woody and have some branches. Plants grow up to 3 metres tall. Stalks have distinct scars where leaves have fallen. The leaves tend to be near the ends of branches. The leaves are divided like the fingers on a hand. The leaves have long leaf stalks. The leaves have 3 - 7 long lobes which can be 20 cm long. These are widest about 1/3 of the distance from the tip and taper towards the base. The colour varies. It produces several long tubers. These can be 50 cm long by 10 cm across. The flowers are on short stalks around a central stalk. They are produced near the ends of branches. The female flowers are near the base of the flower stalk and the male flowers higher up.



**Distribution:** A tropical plant. Plants grow from sea level up to about 1,650 m. In Fiji they grow to 900 m. They can grow in poor soil and can survive drought. It is native to tropical America. It grows between 25°N and 25°S and needs a rainfall above 750 mm. It suits hardiness zones 10 - 12.

**Use:** The tubers are eaten after thorough cooking. They are boiled, roasted or made into flour. The starch is used in puddings, soups and dumplings. Young leaves are edible after cooking. They are also sometimes dried and stored. Seeds are also eaten. **Caution:** Bitter kinds of cassava contain poison but this is destroyed on heating. This kind of cassava should be cooked, sun dried, soaked and cooked again.

**Cultivation:** Cassava is planted from sections of the stalk. Sections about 15 - 20 cm long of the more mature woody stem are cut and stuck into the ground. They can be completely buried or put at almost any angle and it affects the growth little. Soon roots form and leaves start to sprout from the stalk. Cassava seeds need a soil temperature of 30°C for their germination. Flower and fruit production is more common under lower temperatures such as in highland or less equatorial conditions.

It is not necessary to dig a hole to plant cassava and on many soils where the soil is loose it can be planted without digging the soil first. Cassava does not suit waterlogged soils and preferably they should not be too shallow or stony.

Cassava can be planted at any time of the year but to get started it needs moisture so is often planted near the beginning of the wet season. The crop once established can survive for several months without rain. The ability to tolerate drought varies significantly with cultivar. During drought less and smaller leaves are produced and leaves die off more quickly but storage roots can be increased in the short term.

Because cassava can still grow satisfactorily in poorer soils it is often put last in a rotation after others crops have already been grown on the piece of land. Cassava is more responsive to nitrogen and potassium than phosphorus under many field situations. Nitrogen can increase cyanide levels. Under very acid conditions with high soluble aluminium levels, cassava has been able to achieve and maintain top growth but with significantly reduce root yields. When drainage is good and soil

moisture is adequate, cassava stalks can be planted at any orientation from horizontal to vertical, but in very sandy soils horizontal planting is best and in heavy clay soils vertical planting is best.

Because of the slow growth in early establishment stages, soil loss from erosion with heavy rains can be significant. To avoid this planting should be timed so that the maximum vegetative growth is occurring during the heaviest rains. A leaf area index between 2.5 - 3.5 is optimal for cassava yield. The critical period for weed control is the time from 2 - 8 weeks after planting. Cassava tuber bulking is delayed under shaded conditions. Yields are also reduced. In mixed cropping situations using crops which mature early, allowing the cassava time to recover, is one possible strategy. For optimum production shading should be avoided.

Cassava takes about 10 - 12 months to produce mature tubers in the lowlands tropics although some varieties produce a smaller yield earlier. Yields in the range of 20 - 45 t/ha have been recorded for 12 - 14 month crops. The plants can be left growing and the tubers stored in the soil for considerable time. Crops of 24 months duration occur. Once the tubers have been dug they do not keep for more than a few days. Pre-harvest pruning of plants increases the storage time of tubers after harvest.

Spacing and plant density varies with soil climatic conditions and variety. Plant densities from 10,000 to 30,000 plants per hectare are used. Plants from the higher density crops have been shown to have quick post harvest deterioration. Mulching has given significant yield increases in some conditions. It also reduces the incidence and damage of some root boring insects.

**Production:** Plants can be harvested after 10 months in the lowlands. There are some faster growing varieties. Yields in the range of 20 - 45 t/ha have been recorded for 12 - 14 month crops.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber	62.8	625	1.4	30	15	0.23	0.48
leaf	82.0	382	7.1	57	275	7.6	-

## Starchy staples

**English:** Elephant foot yam    **Scientific name:** *Amorphophallus paeonifolius* var. *campanulatus*

**Local:** Cây chột nưa, cây nưa chuông, khoai nưa hoa chuông    **Plant family:** ARACEAE

**Description:** A taro family plant but with a very divided leaf. It grows to 1 m tall. It is a herbaceous plant with rough and mottled leaf stalks. It has a straight stem and the leaf is divided into leaflets. The leaves can be 1 m wide. The leaves usually come singly from the ground. The flower stalk can be 3 - 20 cm long. The edge is curved back and wavy. The flower is dull purple and up to 30 cm across. The flower gives off a bad smell like rotting meat and this attracts flies. The flower only develops after the leaves have died off. The leaves and corms especially in the wild varieties contain many stinging crystals. It has a large round tuber up to 25 cm across.



The large round underground corm produces small corms around the side. These can be 10 cm long.

**Distribution:** A tropical plant. It occurs mainly in seasonally dry areas and grassland up to 800 m altitude in equatorial zones. It requires an average temperature of 25 - 35°C and rainfall of 1,000 - 1,500 mm during the growing season. Soils need to be well drained as it cannot stand waterlogging.

**Use:** The corm is cooked and eaten. The leaves are edible. The young petioles or leaf stalks are eaten cooked.

**Cultivation:** The cormels are planted. Seeds will grow but flowers need hand pollination. Small corms from around the side are the normal part planted. If a very small corm is planted, the plant may need to grow for several seasons to produce a large yield. Setts or small cormels of 200 g are suitable for use planted at 30 cm x 30 cm spacing and produce seed corms of about 500 g. Larger corms take 3 - 4 years to produce. This is achieved by digging up corms and replanting next season. Each crop takes about 8 months to mature. Corms are planted 15 cm deep. Spacing is increased between plants in successive years of growth. After harvest, the corm needs to be kept for a few months before it is ready to produce a new shoot and re-grow.

**Production:** The stalk dies back when the plant is mature. The corm will keep for several months. An individual corm can finally weigh 8 kg. Eventually the corm under the ground increases in size then the leaf dies back. The corm could be harvested and stored, or eaten at this stage. If it is just left, a very large flower is produced. This type of growth pattern where vegetative growth is followed by a storage organ with dormancy is the type of growth that suits areas with a distinct wet and dry season. It has the advantage that the corm will store well after harvest and can be eaten in the dry season when food is short.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber	84.5		0.9			1.4	0.2

## Starchy staples

**English:** Potato yam

**Local:** Khoai dái, củ dại, khoai tròi

**Scientific name:** *Dioscorea bulbifera*

**Plant family:** DIOSCOREACEAE

**Description:** A yam with a long smooth stemmed vine, round in cross section and without spines. The vine winds to the left, can climb into trees and grow to long lengths. The large leaves (14 - 30 cm across and slightly longer than wide) have pointed tips and round bases. About 7 veins arise from the tip of the leaf stalk. It produces often flattened bulbils (potatoes) in the leaf angles along the vine. They can be grey brown or purple. The smaller tuber underground is normally covered with roots. The flowers are large. The male flowers are in spikes up to 20 cm long. The female spikes are usually in pairs. The winged fruit are about 2.5 cm long by 1.5 cm across. The seeds have wings. The bulbils normally have few fibres through the tissue compared to some yam tubers. Many varieties have yellow flesh.



**Distribution:** An annual tropical plant. It will grow from the coast up to about 1700 m altitude in equatorial zones. It is common near the edge of grassland and forest at mid altitudes. Both wild and cultivated forms occur. It is common near secondary forest at low and medium altitudes.

**Use:** The cooked tubers are eaten. More commonly the cooked aerial bulbils are eaten. Some kinds are bitter and inedible or at least require special processing and cooking. Some varieties are poisonous.

**Cultivation:** Either the vine bulbils or the underground tubers are planted. It is convenient to train the long vines up trees. The bulbils need a set storage time before sprouting. The leaves die off for 1 - 4 months each year before re-sprouting from the tuber. Bulbils only grow shoots from one end unless the bulbil is cut into pieces. If the larger bulbils are cut, the cut surfaces should be dried and healed in a shady place for 2 - 3 days before planting. Bulbils are planted 8 - 12 cm below ground at a spacing of about 100 cm by 100 cm. Normally nitrogen and potassium fertilisers give greater responses than phosphorus. Friable well drained soils are most suitable. Often little cultivation or mounding is done. A high level of organic matter improves yield. Strong staking is required with branched 2 m stakes, or with trees or living stakes. Vine and tuber growth can be extensive and heavy.

**Production:** Bulbils or aerial yams are produced as leaves begin to unfold, continue until plant maturity, and often fall. Harvesting can start 3 months after planting but immature tubers have less starch. Underground tubers are normally not harvested until leaf die back. Wounds and damage to the tubers normally heal naturally given dry aerated conditions. Some varieties have seasonally dormant aerial tubers which only grow after an extended storage period. Others germinate quickly.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber	70.8	357	2.7	-	78	3.1	0.4
bulbil	79.4	326	1.4	-	-	2.0	-

## Starchy staples

**English:** Sweet potato

**Local:** Khoai lang

**Scientific name:** *Ipomoea batatas*

**Plant family:** CONVULVACEAE

**Description:** This is a root crop which produces long creeping vines. The leaves are carried singly along the vine. Leaves can vary considerably from divided like fingers on a hand to being entire and rounded or heart shaped. Purple trumpet shaped flowers grow at the end of the vine. Under the ground fattened tubers are produced. There are a large number of varieties which vary in leaf shape and colour, tuber shape, colour, texture and in several other ways.



**Distribution:** A tropical and subtropical plant. They grow from sea level up to about 2,700 m altitude in the tropics. Plants can grow with a wide range of rainfall patterns and in different soils. Plants are killed by frost and can't stand water-logging. Plants grow well with temperatures between 21 - 26°C. It can grow with a pH between 5.2 - 6.8. Sweet potato are not tolerant to shading. Under shaded conditions, both foliage growth and storage root production are decreased. Some cultivated varieties can be selected for increased production under mild shade but not heavy shade. The survival of cuttings at planting is also reduced under shaded conditions. Under shaded conditions plant become more climbing and with fewer leaves which are however larger. With increasing shade less tubers are produced and these grow more slowly. Sweet potato tends to be responsive to potassium fertiliser. cultivated varieties are often selected for yield under low fertility conditions. Under lowland conditions in the tropics sweet potato tubers undergo active tuber enlargement from 6 - 16 weeks. Weed control is essential especially during early stages of growth. The rate of ground coverage by foliage varies greatly with growing conditions and cultivar but once ground coverage has occurred weed control is less of a problem. Sweet potato tuber initiation is subject to aeration in the soil. Either heavy clay soils, waterlogged conditions or other factors reducing aeration can result in poor tuber production. For this reason sweet potatoes are often grown on mounded beds. It suits hardiness zones 9 - 12.

**Use:** Tubers are boiled or baked. They can be steamed, fried, mashed or dried. They can be fermented into alcoholic drinks. They can also be used in pies, cakes, puddings and candies and jams. They can be used in noodles. The chopped and dried tubers can be boiled with rice or ground into flour and mixed with wheat flour to make cakes or bread. The young leaves are edible.

**Cultivation:** Vine cuttings are used for planting. In grassland soils it is grown in mounds, ridges or other raised beds. In bush fallow, it is mostly planted in undug loose soils. It needs a sunny position. Tubers won't form if the ground is waterlogged when tubers start to develop. Sweet potato is grown by cuttings of the vine. About 33,000 cuttings are required per hectare. These weigh about 500 kg. Vine lengths of about 30 cm are optimum. As long as the vine is adequately inserted in the soil, the length of vine inserted does not significantly affect yield. Fresh sweet potato seeds germinate relatively easily and lead to continuous production of new cultivars under tropical conditions. Excess nitrogen restricts storage root initiation and therefore excess leaves are produced without significant tuber yield. Dry matter percentage increases with increasing age of the crop. Higher dry matter tubers are normally preferred.

Sweet potato are not tolerant to shading. Under shaded conditions, both foliage growth and storage root production are decreased. Some cultivars can be selected for increased production under mild shade but not heavy shade. The survival of cuttings at planting is also reduced under shaded

conditions. Under shaded conditions plant become more climbing and with fewer leaves which are however larger. With increasing shade less tubers are produced and these grow more slowly. Sweet potato tends to be responsive to potassium fertiliser. Cultivars are often selected for yield under low fertility conditions.

Under lowland conditions in the tropics sweet potato tubers undergo active tuber enlargement from 6 - 16 weeks. Weed control is essential especially during early stages of growth. The rate of ground coverage by foliage varies greatly with growing conditions and cultivar but once ground coverage has occurred weed control is less of a problem. Sweet potato tuber initiation is subject to aeration in the soil. Either heavy clay soils, waterlogged conditions or other factors reducing aeration can result in poor tuber production. For this reason sweet potatoes are often grown on mounded beds. In well drained or high organic matter soils digging or mounding is not as essential. Leaf scab (*Elsinoe batatas*) can significantly reduce yield especially in sites where leaf production is low due to low soil fertility. To reduce sweet potato weevil damage plants need to be hilled or have the tubers well covered with soil. Cracking soils can allow the weevil access to tubers.

**Production:** The time to maturity ranges from 5 months to 12 months depending on the variety planted and the altitude at which it is being grown. Yields range from 6 - 23 t/ha.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber (baked)	72.9	431	1.7	961	24.6	0.5	0.3
tuber (raw)	70.0	387	1.2	709	25	0.7	0.4
tuber (boiled)	72.0	363	1.1	787	15	0.6	0.3
leaf	86.3	168	3.9	105	58	2.9	-

## Starchy staples

**English:** Taro  
**Local:** Khoai môn

**Scientific name:** *Colocasia esculenta*  
**Plant family:** ARACEAE

**Description:** This plant has large flat leaves on the end of upright leaf stalks. It grows up to 1 m high. The leaf stalk or petiole joins the leaf towards the centre of the leaf. The leaves are 20 - 50 cm long. Near the ground a thickened rounded corm is produced. Around this plant there is normally a ring of small plants called suckers. Many different varieties occur. If left to maturity, a lily type flower is produced in the centre of the plant. It has a spathe 15 - 30 cm long which is rolled inwards. The flowers are yellow and fused along the stalk. There are many named cultivated varieties. Taro comes in two basic forms. The Dasheen type *Colocasia esculenta* var. *esculenta* and *Colocasia esculenta* var. *antiquorum* or the Eddoe type. The basic difference is the adaptation of the Eddoe type to storage and survival in seasonally dry places, while the dasheen type needs to be maintained in a more or less continuously growing vegetative stage.



**Distribution:** It is a tropical plant. Taro grows from sea level up to about 2,300 m altitude in the tropics. It grows well in humid places. It can stand damp soil and grow under light shade. It suits hardiness zones 9 - 12.

**Use:** The corms, petioles and leaves are all edible after cooking. The leaves are also dried and stored. Fresh leaves can be stored for 4 - 5 days. **Caution:** Some varieties burn the throat due to oxalate crystals.

**Cultivation:** Taro can be planted from cormels or from the top of the central corm. Other sections of the corm could also be used but this is not commonly done. Flowering of taro and seed production can lead to new cultivars. Flowering can be promoted by the use of gibberellic acid. The general growth pattern is for an increase in top growth, in terms of leaf number, leaf area and petiole length, to continue for about 6 months under tropical lowland conditions then for each of these to decrease and tuber storage to continue to increase. Corm weight increases significantly from 5 - 11 months. Starch content also increases with time but protein content declines over the corm development period.

Taro can be grown under flooded conditions but root rots develop if the water becomes stagnant. For flooded cultivation, the land is cleared, ploughed, cultivated and puddled. The aim is to get a field that is flat with embankments allowing the impounding of water. Planting is done into 2 - 5 cm of standing water.

For dryland taro, the soil is prepared by digging, unless a fresh bush fallow is used where the natural friability of the soil allows plants to be put into the undug soil in a small hole that is prepared. Plants are put into a hole 5 - 7 cm deep or deeper. Mulching to conserve moisture and reduce weed growth is beneficial. Setts from corms normally give higher yield than that from cormels. The greater leaf area and root production may be responsible for this. Setts of about 150 g are optimum.

The time of planting is primarily determined by the availability of moisture. Planting is done shortly after the rainfall has become regular, if seasonally distinct wet and dry occur. Higher rainfall, higher temperatures, and higher hours of sunlight, enhance production and determine seasonality of production.

Evapotranspiration for flooded taro averages about 4 mm per day, ranging from 1.5 - 7.2 mm, with a total of about 1,200 mm for the crop. Intermittent moisture can result in irregular shaped corms. Flooding has been found to be more effective than sprinkler irrigation, or furrow irrigation. Increased suckering, giving greater leaf area, seems to be the reason for this.

Taro is sensitive to weed competition throughout most of its growth, but it is more critical during early growth up to 3 - 4 months. About 7 - 9 weedings are required, to keep the crop clean under tropical lowland conditions, where flooding is not used. Due to the decrease in height and leaf area towards the end of the growth cycle when starch accumulation in the corms is maximum, weed competition and weed control are again significant. Mechanical weeding needs to be shallow to avoid damaging the superficial taro roots. A range of herbicides have been recommended in various situations.

Taro produces the highest dry matter yield under full sunlight, but it can still grow under moderate shade. Under shaded conditions it grows more slowly and develops fewer cormels. They require good moisture conditions and have little tolerance for drought. Taro residue has an allelopathic factor which can reduce the germination and growth of other plants, for example, beans.

Taro tends to demand high fertility, and is responsive to additional NPK fertiliser. Higher doses of K increases starch content and higher doses of N increases protein content. Both N and K applications increase oxalic acid content of the tubers.

Spacing affects total yield, and marketable, harvestable yield, of corms. Close spacing increases the corm yield per area, and the shoot yield per area, but decreases the corm yield per plant, and the contribution of sucker corms, to the yield. Where spacings of 30 cm x 30 cm are used, giving about 110,000 plants per hectare, a very large amount of planting material is required, which reduces the net return per unit of planting material. A spacing of 60 cm x 60 cm is more common. Wider spacings of 90 cm x 90 cm reduces overall yield.

**Production:** Crops mature in 6 - 18 months. Yields of 5 - 15 tonnes per hectare are probably average.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
root	66.8	1231	1.96	3	5	0.68	3.2
leaf	85.0	210	5.0	57	90	0.62	0.7
leaf stalk	93.0	101	0.5	180	13	0.9	-
leaf (cooked)	92.2	92.2	2.7	424	35.5	1.2	0.2

## Starchy staples

**English:** Mung bean

**Local:** Cây đậu xanh

**Scientific name:** *Vigna radiata*

**Plant family:** FABACEAE

**Description:** An upright hairy bean plant which can grow to 1 m tall. It has many branches. The leaves have 3 leaflets, are dark green and grow on long leaf stalks. There are oval stipules at the base of the leaf. Flowers are pale yellow and small. They occur in bunches of 10 - 20 on the ends of long hairy flower stalks. Pods are black and straight. They do not have a beak. Pods contain 10 - 20 seeds which are usually green or golden yellow. They are smaller than black gram. The beans can be black. They have a flat white hilum. There are 2,000 varieties.



**Distribution:** A tropical and subtropical plant. The plant will grow from sea level up to about 2000 m in the tropics. It is drought resistant but can't stand water-logging. Plants are damaged by frost. They cannot stand salinity. Rainfall at flowering is detrimental. It requires a deep soil. Both short day and long day varieties occur. It can grow where annual temperatures are from 8 - 28°C. It can tolerate a pH from 4.3 - 8.1. It suits a drier climate and can grow in arid places. It suits hardiness zones 10 - 11.

**Use:** Seeds are eaten ripe, raw or roasted. They are added to soups and stews. They are also fermented. Young pods and leaves can be eaten. The seeds can be germinated for sprouts and used in salads and stir-fried dishes. The seeds are ground and used for starch to make noodles.

**Cultivation:** Plants are grown from seed. In some areas these are broadcast while for small plots often 2 - 3 seeds are sown in holes 50 - 60 cm apart. Seeding rates of 6 - 22 kg per ha are used in different locations. It normally requires phosphorus fertiliser for adequate growth. Seeds germinate in 3 - 5 days.

**Production:** Green pods are ready after about 2 months and ripe pods may take another 1 - 2 months. For ripe beans the whole plant is harvested and dried before threshing. Yields of 450 - 560 kg/ha of seeds are common.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	11.0	1432	22.9	55	4	7.1	-
seed (cooked)	-	439	7.0	2.4	1.0	1.4	-
seed (sprouted)	90.4	126	3.0	2	13.2	0.9	0.4

## Legumes


**English:** Jack bean

**Local:** Cây đậu kiể

**Scientific name:** *Canavalia ensiformis*

**Plant family:** FABACEAE

**Description:** A perennial climber, although short kinds do occur. Often it is a more bushy plant than the sword bean. Plants grow up to 1.5 m long. Stems can be hairy. Leaves have 3 leaflets. The leaflets are oval and 5.7 - 20 cm long by 3.2 - 11.5 cm wide. The leaf tends to be wedge shaped at the base. The leaf stalks are 2.5 - 11 cm long. Flowers are red/purple. They occur on flower clusters 5 - 12 cm long and with flower cluster stalks which are 10 - 34 cm long. The individual flower stalks are 2 - 5 mm long. Pods are long and sword shaped. Pods can be 15 - 35 cm long. Seeds are white with a light brown hilum half as long as the seed. Seeds are 2 cm long, by 1 cm across.



**Distribution:** It grows in tropical and subtropical places. It requires a fairly high temperature (15° - 30°C). It will possibly grow up to 900 m altitude. It is fairly drought resistant and also has some resistance to water-logging and salt in the soil. It can tolerate shade. It can tolerate pH from 4.5 - 8.0 but does best at about 6.1. The optimum mean annual temperature is 14.4° - 27.8°C. Seed germinate between 24 - 27.5°C. It is a short day plant growing well with a daylength of 10 - 12 hours of sunlight. It can grow in arid places.

**Use:** The leaves and top shoots are eaten. The very young pods are boiled and eaten. The flowers can be eaten. The young seeds are eaten boiled, roasted, or peeled and cooked. The seeds are also fermented. The ripe seeds are roasted and used as a coffee substitute. **Caution:** The ripe seeds can contain poison and need to be well cooked and the water changed before eating. They are also often left under running water or fermented.

**Cultivation:** It is grown from seeds. Seeds need to be 2 cm deep. A spacing of about 60 cm is suitable. Plants preferably need a support to climb over. It benefits from a fertile soil but adding nitrogen depresses yield.

**Production:** Green pods are produced in 3 - 4 months, but ripe seeds need 6 - 9 months. Yield of seeds can range from 700 - 5,400 kg/ha.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	10.0	1423	20.4	160	0	4.9	-
pod (fresh)	88.0	155	2.4	-	-	-	-

## Legumes

**English:** Sweet acacia, Prickly Moses

**Local:** Keo thom, keo ta, mâm xôi

**Scientific name:** *Acacia farnesiana*

**Plant family:** FABACEAE

**Description:** An evergreen shrub. It grows 5 - 7 m tall and 3 m across. The stem is slender and erect. The crown is open. It is a spreading, densely branched shrub. The bark is smooth and brown. The leaves are branched and green. There are 4 - 6 pairs of larger leaves and 10 - 20 pairs of small leaflets. They have tiny leaflets and thorns up to 2 cm long, occur in pairs. The leaf stalk has a gland at or above the middle. The flowers are large orange balls. They are strongly perfumed. (The oil is used as a perfume in France.) The pods are long and dark brown to black. They are 5 - 8 cm long by 0.5 - 1 cm wide. They are inflated and sausage like. Often they are curved. They are marked with narrow lines. The pods have hard grey seeds imbedded in a pithy substance. The pods do not split open at maturity. The seeds are chestnut brown and 7 - 8 mm long by 5.5 mm wide.



**Distribution:** It is a tropical plant. This tree occurs naturally in Australia, Asia and Africa. It will grow on most soils. It is drought and frost resistant. It most commonly grows naturally on clay soils. In Papua New Guinea the plants are coastal below 60 m altitude. It grows in areas with an annual rainfall between 400 - 4,000 mm. It can grow in acid or alkaline soils. It can grow in arid places. It suits hardiness zones 11 - 12.

**Use:** The pods have been recorded as eaten after cooking. The gum is eaten. The ground up seeds are eaten. The germinated seeds are claimed to be eaten. The gum is used to prepare sweets. The young leaves are used in India as a substitute for tamarind in chutneys.

**Cultivation:** It is grown from seed.

**Production:** It is fast growing. Flowering can occur almost continuously if watering is regular. In northern Australia, flowering is normally May to July, with pods available from September to November.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	8.1	1522	36.6	-	-	6.0	0.6

## Legumes

**English:** Pigeon pea  
**Local:** Cây đậu triều

**Scientific name:** *Cajanus cajan*  
**Plant family:** FABACEAE

**Description:** An upright perennial shrubby legume that can live for 3 - 4 years. They can grow up to 4 m tall and spread to 1.5 m wide. It has a bushy appearance and a strong deep taproot. The root nodules are round and sometimes lobed. The leaf consists of 3 narrow, green leaflets which are silvery-green underneath. The end leaflet is larger with a longer leaf stalk. The pea shaped flowers are red and yellow and occur on branched flower stalks which stick upwards in the axils of leaves. Pods are long, straight and narrow, often with 4 - 8 seeds. Seeds vary in shape, size and colour. The pods are slightly hairy. Pods are often 4 - 8 cm long and have a beak at the end. Pods are constricted between the seeds. Many varieties of pigeon pea occur. Some are dwarf and day length neutral.



**Distribution:** A tropical plant that requires a tropical or subtropical climate. Plants grow from sea level up to about 1,800 m in the tropics. They can tolerate drought and are suited to a drier climate. They can grow in places with less than 600 mm rainfall per year. They do less well in the wet tropics. They suffer in waterlogged soils and are damaged by frost. It can also tolerate heat. It will grow on poor soils cannot grow on salty soils. It can grow in arid places and suits hardiness zones 10 - 12.

**Use:** Young leaves, shoots and pods are eaten. The pods can be used in curries. The leaves and shoots as potherbs. Young seeds are cooked and eaten like peas. Ripe seeds are also cooked and eaten in soups and curries. Bean sprouts can be produced and eaten. Preparation of the seeds for dahl is somewhat complicated.

**Cultivation:** They are grown from seeds. It is best to sow seeds where the plants are to grow. Seeds normally germinate easily and well. Before sowing seed it helps to soak them in cold water for one day. Seeds store well if kept cool and dry. A spacing of 1.5 m x 1.5 m is suitable. Plants can be cut back and allowed to re-grow. Plants can also be grown from cuttings.

**Production:** Plants are fast growing. Pods are ready after 5 months. Mature seeds take about 8 months. Plants will often live for 3 - 4 years. Plants are cross pollinated by insects, or self pollinated.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	10.0	1449	19.5	55	-	15.0	-
pod (young)	64.4	477	8.7	-	-	2.0	-
seed (young, boiled)	71.8	464	6.0	2	28.1	1.6	0.8

## Legumes

**English:** Soybean

**Local:** Cây đậu nành, đậu tương

**Scientific name:** *Glycine max*

**Plant family:** FABACEAE

**Description:** A small erect bean growing up to 60 cm tall. It grows each year from seed. Straggling kinds can occur. Stems, leaves and pods are softly hairy. The leaves have 3 leaflets. The leaflets have stalks. Flowers are small and white or blue. They occur in groups in the axils of leaves. The pods are broad, flat and hairy. Pods have 2 - 4 seeds. The seeds can be yellow to black.



**Distribution:** It is a temperate plant that suits lowland areas. It can be grown from sea level to 2,000 m altitude. Many varieties will not flower in the tropics (short days). It needs fertile soil. The best soil acidity is pH 5.5 - 7.0. It is damaged by frost.

**Use:** The young pods and ripe seeds are eaten. They are used for flour. The dried seeds are boiled or baked and used in soups, stews and casseroles. The seeds are used for oil. Toasted seeds are eaten like a snack. Strongly roasted seeds are used for coffee. Soy flour is used for noodles, and confectionary. The beans are fermented and used in a range of foods. Sometimes the young leaves are eaten. The seeds are also used for sprouts and for making cooking oil and soya sauce etc. Because soybean contains a trypsin inhibitor they should be cooked and even the sprouts should be lightly cooked.

**Cultivation:** It is grown from seed. Seeds need to be inoculated with bacteria before planting. Plants need to be about 20 cm apart.

**Production:** Plants flower about 8 weeks after sowing and pods mature about 16 weeks after sowing. Often plants are pulled up and hung up before threshing out the seed.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	9.0	1701	33.7	55	-	6.1	-

## Legumes

**English:** Winged bean

**Local:** Cây đậu rồng

**Scientific name:** *Psophocarpus tetragonolobus*

**Plant family:** FABACEAE

**Description:** A climbing perennial bean up to 4 m tall. It can re-grow each year from the fattened roots. The stems twine around supports or trail over the ground. The leaves have 3 leaflets 8 - 15 cm long. The leaf stalks are long. The flowers are blue or white. They occur on the ends of branches from within the axils of leaves. Pods have wavy wings and are roughly square in cross section. They are 6 - 36 cm long with 5 - 30 seeds. Seeds can be white, yellow, brown or black. They are bedded in the solid tissues of the pod. The seeds are round, smooth and brown with a small hilum.



**Distribution:** A tropical plant that grows from sea level up to about 1850 m altitude in the tropics. It normally only produces tubers at 1200 - 1850 m altitude. Winged bean is a short day plant and needs a daylength less than 12 hours. Because of the effect of daylength, it will not produce flowers or pods at places far removed from the equator. The main areas of production occur between 20°N and 10°S latitudes. It is ideally suited to the tropics including the hot humid lowlands. For maximum seed production, winged beans need temperatures of 23 - 27°C and for tubers the temperatures should be 18 - 22°C. Winged beans can grow on a wide variety of soils and have been grown on soils with pH from 3.6 - 8.0. Soils which are very acid have soluble aluminium to which winged beans are sensitive. Soils should not be waterlogged.

**Use:** The young leaves, flowers, young pods, ripe seeds and root tubers are edible. The seeds can be used to extract an edible oil.

**Cultivation:** Seeds are sown at the beginning of the rainy season. Seeds germinate and grow slowly for the first 3 - 5 weeks. For tubers, vines are pruned off at about 1 m high (or left unstaked) and some flowers are removed. Cultivation procedures vary slightly depending on which part of the plant is to be eaten. Short podded winged bean are used for tubers and long podded ones have poor tubers. Tuber production is not as efficient in tropical lowland conditions.

**Production:** The first green pods are ready about 10 weeks after sowing. Tubers are ready after 4 - 8 months. Seed yields of 1.2 tons/ha and tuber yields of 4 tons/ha are possible. A single plant can produce up to 75 pods. Dry bean yields of 45 - 330 g per plant can be produced depending on variety. Tuber yields of 5500 - 12000 kg per hectare have been produced. Seeds can contain a trypsin inhibitor which reduces protein digestibility. This inhibitor is destroyed by soaking seeds then boiling them well. Tubers can also contain this chemical and need to be well cooked.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	8.5	1764	41.9	-	-	15.0	4.5
pod (fresh)	92.0	105	2.1	-	-	-	-
leaf	95.0	197	5.0	809	30	6.2	1.3
seed (young)	87.0	205	7.0	13.0	18.3	1.5	0.4
root	57.4	619	11.6	0	0	2.0	1.4

## Legumes

**English:** Yard long bean

**Scientific name:** *Vigna unguiculata subsp. sesquipedalis*

**Local:** Cây đậu đũa

**Plant family:** FABACEAE

**Description:** A climbing bean with long pods. The vines can be 3 m long. They normally twine around sticks. Dwarf kinds also occur. Leaves have 3 leaflets. The leaflets are oval and side leaflets are at an angle. Leaflets can be 2 - 16 cm long by 1 - 12 cm wide. There can be few or several flowers. The flowers can be white, yellow or blue. Flowers are 1 - 3 cm long by 1 - 3 cm wide. Pods are long (up to 90 cm) and flexible. The seeds can vary between white to dark brown. They are oblong or kidney shaped. Seeds are 4 - 12 mm long by 2 - 6 mm wide.



**Distribution:** A tropical plant that grows in coastal areas in the tropics from sea level up to about 300 m in equatorial regions. Seeds germinate in moist soil over 22°C. For growth, day temperatures between 25 - 35°C and night temperatures above 15°C are required. It suits wet areas. Plants are day length neutral. They perform best under full sunlight but can tolerate some shade. It has a high water requirement for the full grown crop (6 - 8 mm per day). It can tolerate a wide range of soils with pH of 5.5 - 7.5.

**Use:** The young pods, ripe seeds and leaves are eaten. They can be steamed, stir-fried and used in other dishes. The seeds can be sprouted and eaten.

**Cultivation:** Plants are grown from seed. Seeds germinate quickly (2 - 3 days) and plants grow rapidly. Flowering occurs after 5 weeks and harvesting of young pods can start 2 weeks later. Plants die after about 3 - 4 months. A spacing of 60 cm is suitable. Plants need sticks about 2.5 m long to climb up. Often 5 or 6 seeds are sown around the one stick. Plants are often topped when growing too vigorously. It grows as an annual bean so seeds need to be replanted each year. The pods need to be harvested every 2 - 3 days. Diseases get worse if the bean does not have sticks to climb. The damage by bean pod borer is less if snake beans are grown inter-cropped with maize.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (fresh, raw)	87.9	197	2.8	11.3	18.8	0.47	0.37
seed (boiled)	87.5	197	2.5	20	16.2	0.98	0.36
seed (dry, raw)	8.43	1453	24.33	0.16	1.6	8.61	3.5

Food values obtained from: U.S. Department of Agriculture, Agricultural Research Service. 2011. USDA National Nutrient Database for Standard Reference, Release 24. Nutrient Data Laboratory Home Page, <http://www.ars.usda.gov/ba/bhnrc/ndl>

## Leafy greens

**English:** Amaranths

**Local:** Rau dền

**Scientific name:** *Amaranthus tricolor*

**Plant family:** AMARANTHACEAE

**Description:** A small, annual, leafy green herb about 1 m high, spreading to 45 cm wide. An upright, much-branched annual with a thin membrane covering the stems. Sometimes the plant lies over. The stems are angular. The plant branches in the upper part of the plant. It does not have thorns and grows from seed each year. Leaves have long leaf stalks which can be 5 - 10 cm long. Leaves vary in shape, size and colour. The leaf blade can be 5 - 25 cm long by 2 - 6 cm wide. Leaves are dull-purplish and the top leaves can be yellow or red. Some types have coloured leaves or patterns on the leaves. It has a clumpy seed head at the top. The flower spike at the top can be 30 cm long. The seeds are 1 - 1.2 mm across and black.



**Distribution:** It grows in most tropical countries, including the Pacific and Solomon Islands. It will grow in warm, temperate places. Plants grow wild in waste places. Amaranths grow from sea level to 2,400 m altitude in the equatorial tropics. Amaranth seeds need a temperature above 15 - 17°C to germinate. In areas of the equatorial highlands above 1,800 m, average temperatures are probably below this during the cooler months. It may be more difficult to get Amaranths started during these months. It suits hardiness zones 8 - 11.

**Use:** The young leaves and stems are cooked and eaten as a vegetable. The seeds can be eaten. It is a very important tropical vegetable. It grows quickly, produces well and is nutritious.

**Cultivation:** The very small seeds of this plant are scattered over ashes or fine soil in fertile ground. The seeds are normally spread by rubbing the dry seed heads between the hands. Some types are self-sown. These plants grow in most tropical countries. The soil must be fertile. If they are put in an old garden, they will grow very poorly. They should either be put in a new garden site, when it is cleared from bush, or in old ground that is had compost added. Small gardens close to a house can often be built up to a good fertility by using food scraps and ashes that are left over near houses. Amaranths need high amounts of two nutrients, nitrogen and potash. The ashes from fires are high in potash, so farmers scatter seeds of Amaranth over areas they have burnt. Normally, the hotter it is, the better they grow. They also like plenty of sunlight and do not suit shaded places. They need to have water most of the time they are growing. This is mostly not an issue in areas with high rainfall.

**Production:** Plants can be harvested when small by thinning out closely-spaced plants. These can be either transplanted or eaten cooked. Plants can be harvested whole or have top leaves harvested several times. Harvesting begins after 4 - 7 weeks and can continue over the next 2 months.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	91.7	96	2.5	292	43.3	2.3	0.9

## Leafy greens

**English:** Horseradish tree

**Local:** Cây chùm ngây, cây cải ngựa

**Scientific name:** *Moringa oleifera*

**Plant family:** MORINGACEAE

**Description:** A small, soft-wooded tree that grows 9 - 12 m tall. The tree loses its leaves during the year. The bark is grey, thick, corky and peels off in patches. The leaves are pale green and the leaf is divided 3 times. The whole leaf is 30 - 60 cm long and the leaflets are usually oval and 1 - 2 cm long. The leaflets are jointed with a gland near the joint. The flowers are pale yellow. They occur in long sprays 30 cm long. Each flower has 5 petals and of these one is erect and 4 are bent backwards. The fruit is a long capsule 30 - 100 cm long by 2 cm wide. The seed capsules are up to 45 cm long. They are roughly triangular in shape. The seeds have 3 wings. Often the fruiting kinds are grown as annual plants.



**Distribution:** A tropical and subtropical plant. They suit the dry lowland areas and grow up to 1,350 m altitude in the tropics. They are not hardy to frost. They cannot tolerate water-logging. A pH of 6 - 7.5 is suitable. It can grow in arid places. It suits hardiness zones 9 - 12.

**Use:** The young tops and leaves are eaten cooked. They are eaten as potherbs or used in soups and curries. They can be dried and stored for later use. The very young long pods are eaten cooked, especially in curries and soup. They are also pickled. The young seeds are eaten roasted or fried. Sometimes the roots are used as a horseradish substitute. A gum from the bark is used as seasoning. The bark is used for tea. The roots, leaves, flowers and fruits are eaten cooked in water and mixed with salt and chili peppers. The oil expressed from the seeds is used in salads.

**Cultivation:** It is best to grow plants from 1 metre long cuttings but they can be grown from seed. They can be used as a hedge and pruned regularly to produce more leaves. Properly dried seed can be stored for a long time in sealed containers in a cool place. Normally perennial types are grown from cuttings and annual types are grown from seed.

**Production:** Trees are fast growing. They can be pruned or topped. With one variety the tree flowers and fruits continuously while with the other variety there are flowers and fruit once per year. The fruit ripens 3 months after flowering. Annual types produce fruit 6 months after planting. Leaves are best dried in the shade to retain more of their Vitamin A.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	76.4	302	5.0	197	165	3.6	-
flower	84.2	205	3.3	-	-	5.2	-
leaf (boiled)	87	189	4.7	40	31.0	2.0	0.2
pod (raw)	88.2	155	2.1	4	141	0.4	0.5
seed	6.5	-	46.6	-	-	-	-

## Leafy greens

**English:** Indian mustard, Leaf mustard

**Local:** Cây xanh, cải cay, cải canh (cải mù tạt)

**Scientific name:** *Brassica juncea*

**Plant family:** BRASSICACEAE

**Description:** It is an erect leafy annual cabbage plant. It grows to about 1 m high. The plant can vary a lot. The dark green leaves are elliptical and deeply divided. The end segment of the leaves is oval. The leaves taper towards the stem and have a strong mustard flavour. The flowers are pale yellow. They have 4 petals. It produces a flower and seed pods at the top. The seed pod is 3 - 5 cm long and narrow. The seeds are reddish-brown.



**Distribution:** A warm temperate plant. It is grown in some highland areas in the tropics. It is often grown as a cool season crop. Most varieties are not frost tolerant. It needs a fertile, well drained soil. A pH of 5.5 - 6.8 is suitable. It can tolerate poor soils. It can grow in arid places. It suits hardiness zones 9 - 11.

**Use:** The leaves are cooked and eaten. They have a bitter taste, so the cooking water needs changing. They can be stir-fried, or added to soups and stews. They can be eaten crystallised in vinegar or salt. They can be used in salads. The seeds can be fried then used as a spice. They also yield an edible oil. They can be sprouted. The leaves are also pickled.

**Cultivation:** The seed is broadcast. They can be put in a nursery and transplanted. A spacing of 25 cm x 25 cm is suitable. Seed germinate in 5 days at 20 - 25°C.

**Production:** Plants grow rapidly. Leaves can be harvested one month after planting. Leaves can be harvested several times. Whole plants can be harvested in about 45 days from transplanting.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	6.9	1964	24.9	-	3	10.0	-
leaf	92.0	108	2.4	31	73	2.7	-

## Leafy greens


**English:** Indian spinach

**Local:** Cây mồng tơi

**Scientific name:** *Basella alba*

**Plant family:** BASELLACEAE

**Description:** An annual or perennial climbing herb with thick fleshy leaves. The vine is smooth and juicy and can be 10 m long. It branches freely. The vine and leaves can be red or green. The leaves are fleshy and pointed at the tip. They can be 8 - 18 cm long and 8 - 10 cm across. They are carried alternately along the vine. Leaves can be heart shaped or oval. It has white, pink or red flowers in short spikes which are in the axils of the leaves. The fruit are round and soft. They can be red, white or black and are 6 - 8 mm across. The seeds are round and black. They are 3 mm across. (Often the ones with heart shaped leaves are called *Basella cordifolia*, the ones with a red stem *Basella rubra* and the short day flowering dark green kind *Basella alba*.)



**Distribution:** A tropical plant. It occurs mostly in the tropical lowlands and is best below 500 m but will grow up to about 1600 m. in the equatorial tropics. It will grow quite well in the temperature range 15 - 35°C. It does not like water-logging but can survive 4 - 12 weeks drought once well established. It requires adequate water during the growing season. The best pH is 5.5 - 7.0. It cannot tolerate salty conditions. Flowering does not occur when day lengths are over 13 hours.

**Use:** The young shoots and leaves are eaten cooked. They are somewhat slimy. In soups and stews the mucilage can be used as thickening. The purple colour of fruit is harmless and is used to colour vegetables and agar-agar. Some lemon juice added to the dye enhances the colour. The leaves can be eaten raw in salads or cooked like a vegetable. The leaves are used to make tea and can also be dried and stored. The seeds can be crushed to use as an edible dye for jellies.

**Cultivation:** It can be sown from seeds or cuttings. Seeds germinate in a few days. Sticks can be provided for support, or it can grow over fences and stumps. If seeds are used, 3 kg of seed will sow one hectare. They are best sown in a nursery and transplanted. A spacing of 1 m is suitable. Plants grown from seed are more productive than those grown from cuttings. When cuttings are used, 20 - 25 cm long cuttings are suitable. Where the plant grows over light soil it can root at the nodes and continue growing. Partial shade, rich fertile soil and adequate moisture favour abundant leaf production. It is responsive to nitrogen fertiliser. Light shade gives bigger leaves. It requires a trellis to climb over. Frequently picking of the bud encourages branching.

**Production:** It is 4 - 6 weeks until the first harvest. It grows reasonably well on poor soils and is fairly resistant to pest and disease. Leaves will only store for one day at 20 - 30°C. Yields of 40 kg of leaves from a 10 metre square bed is possible over 75 days.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	85.0	202	5.0	56	100	4.0	-

## Leafy greens

**English:** Kale

**Local:** Cải xoăn

**Scientific name:** *Brassica oleracea* var. *acephala*

**Plant family:** BRASSICACEAE

**Description:** A cabbage plant. The leaves are ruffled or crinkled but some are more smooth than normal kales. The leaf shape and colour varies considerably. The flowers are yellow or white.

**Distribution:** A temperate plant. Young leaves are used in salads while older leaves are used in cooking.

**Use:** The leaves are eaten boiled, steamed, used in soups and stews. The unopened flower buds are used like broccoli.

**Cultivation:** Plants are grown from seed or transplanted. Plants should be spaced 30 - 45 cm apart no more than 1cm deep.



**Production:** Leaves are harvested 40 - 60 days.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf (raw)	84	210	3.3	70	120	1.7	0.3
leaf (boiled)	91.7	109	2.1	313	18	0.5	0.4

Distribution, cultivation and production information and photograph from: Canberra organic growers society 1977-2010, <http://www.cogs.asn.au/growing-guides/kale/>

## Leafy greens

**English:** Kangkong

**Local:** Rau muống

**Scientific name:** *Ipomoea aquatica*

**Plant family:** CONVULVULACEAE

**Description:** Kangkong is a creeping sweet potato-like plant. It has hollow stems and can float on water. The leaves are green and are normally not divided like some sweet potato leaves, but the shape and size varies a little between different kinds. The trumpet shaped flower looks like a sweet potato flower and is normally white. The runners develop roots at the nodes and also branch. This branching increases when tips are picked off. Some variation in leaf shape can be observed. Leaf shape is less variable than in the related sweet potato, but narrow and broad leafed kinds occur. White and green stemmed kinds occur. Green stemmed kinds have more cold tolerance than white stemmed.



Green stemmed kinds have more cold tolerance than white stemmed.

**Distribution:** Kangkong is a tropical plant. It grows best in short day, stable high temperature, moist conditions. Temperatures need to be above 25°C for satisfactory growth. In equatorial regions plants probably grow up to 1,000 m altitude. Below 23°C the growth rate is too slow for economic production, so production is mainly in the lowland tropics. Optimum soil pH is between 5.3 - 6.0. It suits damp places and grows well in swamps. It can grow as a partly floating plant in swamps and lagoons behind the beach along the coast. In some countries they grow the dry land form in gardens.

**Use:** The young tips of shoots are cooked and eaten. They can be boiled, steamed, stir-fried, or added to soups, stews or curries. The young stems can be used in pickles. The young tips can be eaten raw in salads and the roots are occasionally cooked and eaten.

**Cultivation:** Dryland kangkong is normally grown from seed. Sometimes seed are pre-soaked for 12 - 24 hours prior to sowing. Plants can also be grown from cuttings and establishment is rapid. Top cuttings 25 - 40 cm long can be planted beside a pond.

**Production:** Young tips can be harvested 30 days after planting, and subsequent harvests every 7 - 10 days. Production of new shoots probably declines at flowering. Yields up to 60,000 kg/ha have been recorded.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	90.3	126	3.9	315	60	4.54	-
leaf (boiled)	92.9	84	2.1	-	16	1.3	0.2

## ***Fruit***


**English:** Banana

**Local:** Cây chuối

**Scientific name:** *Musa sp* (A &/or B genome) cv.

**Plant family:** MUSACEAE

**Description:** These are the main group of cultivated bananas. They can be classed into diploid, triploid and tetraploid kinds with various amounts of the A or B parents. They grow 2 - 9 m high. They are large non woody herbs with broad long leaves. Most kinds have several suckers. Bananas grow a soft firm false stem from an underground corm. The fruiting stalk eventually emerges from the top of this false stem and normally curves over pointing towards the ground. Fruit occur in clumps or hands along this stem. The male flowers are in a red bud at the end of the flower stalk.



The colour of the stem, bracts, bud and fruit varies considerably depending on the variety. The fruit can be 6 - 35 cm long depending on variety. They can also be 2.5 - 6 cm across.

**Distribution:** A tropical and subtropical plant. They grow from sea level up to about 2,000 m altitude in the tropics. They are rarely an important food above about 1,600 m. In Nepal they grow to about 1,800 m altitude. They do best in warm and humid tropical climates. Temperatures need to be above 15°C. The best temperature is 27°C. The maximum temperature is 38°C. Bananas grow best in full sun. For best growth, a rainfall of 200 - 220 mm per month is needed. A deep friable soil is best. They can tolerate a pH between 4.5 - 7.5. It suits hardiness zones 10 - 12. It is widely grown in many countries.

**Use:** Fruit are eaten raw or cooked depending on variety. Male buds and flowers are eaten on some varieties. They are cooked as a vegetable. The central pith of the false stem and the underground rhizome are also sometimes eaten.

**Cultivation:** They are planted from sword suckers. Diploids need re-planting annually but many triploids can be re-suckered from the base on the same site. Spacing depends on variety. A spacing of 1,000 – 3,000 plants per hectare is used depending on variety. Suckers are usually put 30 cm deep.

**Production:** Time to maturity varies from 6 - 18 months depending on variety and altitude. Triploids have larger bunches than diploids. Tetraploids are very large plants.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit (cooking)	65.3	510	2.0	113	18.4	0.6	0.1
fruit (sweet)	70.7	365	1.7	-	2	0.9	0.4
flower buds	91.3	109	1.6	-	-	1.0	-

## Fruit

**English:** Dragon fruit

**Local:** Thanh long

**Scientific name:** *Hylocereus undatus*

**Plant family:** CACTACEAE

**Description:** A cactus which climbs. It has triangular stalks. Stalks can be 7 m long. It grows in soil and on rotting plant material. It has aerial roots enabling it to climb. The stems have joints and branched. The spine spots are spaced along the edge of the stems. They have a few short, dark spines. The flowers are white but yellowish-green on the outside and open at night. They are 30 cm long. The fruit are round and red, with long scales. Some varieties are pink or yellow. The skin is thin. The flesh is white with small black seeds.



**Distribution:** A tropical plant that grows best in dry tropical and subtropical climates. It suits the lowlands and needs good light. It suits areas where rainfall is 500 – 1,250 mm per year. It is best in seasonally wet and dry climates. In wet tropical zones, plants may grow well but have trouble setting fruit. It needs a temperature above 13°C to grow. It can tolerate temperatures to 40°C and light frosts. It suits hardiness zones 10 - 12.

**Use:** The fruit is eaten fresh. It is also used for drinks. A syrup of the fruit is used to colour pastries and candy. The unopened flower buds are cooked and eaten as a vegetable.

**Cultivation:** Plants can be grown by seed or stem cuttings. The cuttings are usually dried out for a week or two. The plants are self-sterile and have to be pollinated by bats and moths. Hand pollination is recommended for good fruit production.

**Production:** They can have 4-6 fruiting cycles per year.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	89.4	0	0.5	-	25	0.4	-

## Fruit

**English:** Durian

**Local:** Cây sầu riêng

**Scientific name:** *Durio zibethinus*

**Plant family:** DURIONACEAE

**Description:** A large evergreen tree with buttresses that grows 30 - 50 m tall. The bark is dark red-brown and rough and peels off irregularly. The leaves are alternate and oval with a cover of silver scales on the lower surface. The flowers are on the older branches and form bunches of flowers where the lower flower stalks are longer. The fruit are green to yellow, fat and have sharp spines over the surface. The seeds are completely covered with a yellowish soft, very sweet seed covering. The fruit drop unopened. As the fruit ripens it splits open naturally into 3 - 5 sections which have large seeds embedded in a yellow coloured pulp.



**Distribution:** A tropical plant that grows in the humid tropics, preferably below 300 m altitude, but they may grow up to about 800 m altitude in the equatorial tropics. They need plenty of soil moisture and a rich soil. It can tolerate temperatures down to 23°C and up to 46°C. A rainfall of over 2,000 mm evenly distributed throughout the year is best. A drier period during flowering is beneficial. Poorly drained or sandy soil are not suitable. It can grow well in acid soils with pH 5 - 6.5.

**Use:** The flesh around the seeds is eaten. Fruit need to be eaten within 2 days of falling from the tree. Fruit should only be opened at time of eating as the flesh goes sour. The unripe fruit can be cooked as a vegetable. Fruit are also processed for ice-creams and desserts. The seeds are edible, usually cooked. The young leaves and shoots can be eaten. The raw fruit is used as a vegetable or in soups.

**Cultivation:** Trees grow readily from seed. Seeds need to be fresh. Seeds germinate in about 3 days and can be transplanted in about 4 weeks. Seeds do not breed true. Trees can be grown by budding or grafting. Trees fruit about 7 years after planting. A spacing of at least 14 m between plants is needed. Grafted trees seldom reach 20 m height. Heavy mulching near the trunk can help patch canker (*Phytophthora palmivora*) to develop. Trees should be pruned and shaped for uniform branching and by topping to reduce excessive growth and give optimum production.

**Production:** Grafted trees can produce in 4 - 5 years but seedling trees take 10 years. Flowers are cross pollinated by bats. Normally flowers cannot self pollinate. Flowers open in the afternoon and fall by next morning. Up to 50 fruit can be produced per tree per year. Up to 400 fruit can grow on one tree and this varies with season. Fruit can weigh 3 - 5 kg. Fruiting is seasonal. Fruit take from 90 - 130 days from flowering to maturity depending on variety.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	51.5	794	2.6	-	-	1.0	-
fruit (raw)	61.1	602	2.5	5	24	0.7	0.3

## Fruit

**English:** Mango

**Local:** Cây xoài

**Scientific name:** *Mangifera indica*

**Plant family:** ANACARDIACEAE

**Description:** An erect, branched evergreen tree. It can grow to 10 - 40 m high and is long lived. (Trees grown by vegetative means are smaller and more compact.) Trees spread to 15 m across. It has strong deep roots. The trunk is thick. The bark is greyish-brown. The leaves are simple and shaped like a spear. Some kinds of mangoes have leaves with a wavy edge. They can be 10 - 30 cm long and 2 - 10 cm wide. They are arranged in spirals. The leaf stalk is 1 - 10 cm long and flattened. Leaves are often brightly coloured and brownish-red when young.



These tender leaves which are produced in flushes become stiff and dark-green when mature. The flower stalks are at the ends of branches. They are 10 - 50 cm long and branching. Up to 6,000 flowers can occur on a stalk. Most of these are male and up to 35% have both male and female flower parts. Fruit are green, yellow or red and 2.5 - 30 cm long. The fruit hang down on long stalks. The outside layer of the seed is hard and fibrous and there is one seed inside. Several embryos can develop from one seed by asexual reproduction. The fruit shape and colour vary as well as the amount of fibre and the flavour. India has many varieties and they cannot tolerate humidity.

**Distribution:** A tropical and subtropical plant. It grows in the lowlands. It grows from sea level up to 1300 m altitude in the tropics. It does best in areas below 700 m and with a dry season. Rain and high humidity at flowering reduces fruit set. It thrives best where temperatures are about 25°C but will grow with temperatures from 10 - 42°C. Temperatures of 0°C will damage young trees and flowers. Low temperatures (10 - 20°C) at flowering time will reduce fruiting. As temperatures get lower due to latitude or altitude, fruit maturity is later and trees become more likely to only have good crops every second year. Mangoes can grow on a range of soils. In wetter areas soils with less clay are better. They can withstand occasional flooding. A soil pH of 5.5 - 6.5 is best. Soils with pH above 7.5 cause plants to develop iron deficiency. It grows in the Sahel. It can grow in arid places. It suits hardiness zones 11 - 12.

**Use:** Ripe fruit are eaten raw. Unripe fruit is pickled. Seeds can be eaten cooked. They are boiled or roasted. They are made into meal by powdering. Young leaves can be eaten raw or cooked. Amchur is made from the dried unripe fruit. This is used in curries, and pickles and chutneys. The seed kernels are used for famine food in India. They are boiled, roasted or soaked to remove the bitterness. **Caution:** The sap from the tree or fruit can cause skin problems with some people.

**Cultivation:** Trees are grown by planting fresh seed and they can be transplanted. Mangoes vary in their ability to breed true from seed. When more than one seedling emerges from the seed some of these are asexual and breed true. Clean seed germinate best if they are treated at 50°C for 20 minutes, then planted on their edge with the round bulge upwards and near the soil surface. The husk around the seed should be removed. Seeds germinate in 3 - 6 weeks. The strongest growing seedlings from this seed are used and the others thrown away. The seedlings from the folds of the seed are vegetative while the seedling from the centre of the seedling near the stalk end may be sexual and show variation from type. Other seeds only produce one seedling and these normally vary and can be different from the parent tree. Plants can be propagated by budding, or by grafting using in-arching. This is not easy and care is required. In wetter places, flowers need to be

protected with fungicides to enable fruit to form. If organic manure is used this should not be directly in the planting hole nor immediately against the new plant. Young transplanted seedlings need regular watering. A spacing of 6 - 12 m between plants is used. Wind protection is advisable to prevent fruit rubbing and getting damaged. Trees should only ever be lightly pruned as fruit develop on new growth and heavy pruning can reduce flowering. Flowering can be brought about by foliar sprays of potassium nitrate.

**Production:** Seeds germinate after about 20 days. Seedling trees produce after 4 - 6 years and increase in production up to 20 years. Trees often bear better each second year. Rain at flowering reduces fruit setting. Fruiting is at the end of the year. Fruit take 4 - 5 months to mature. Fruit vary in weight from 200 - 1,000 g. Trees can produce one million flowers but only 500 fruit. Trees last for many years.

**Food Value:** Per 100 g edible portion

<b>Edible part</b>	<b>Moisture %</b>	<b>Energy kJ</b>	<b>Protein g</b>	<b>proVit A µg</b>	<b>proVit C mg</b>	<b>Iron mg</b>	<b>Zinc mg</b>
fruit	83.0	253	0.5	54	30	0.5	0.04
leaf	82.1	226	3.9	-	60	2.8	-

## Fruit

**English:** Mangosteen

**Local:** Cây măng cụt

**Scientific name:** *Garcinia mangostana*

**Plant family:** CLUSIACEAE

**Description:** Mangosteens are a medium sized tree, growing up to 20 m high. Trees keep their leaves all year. Leaves are 15 - 25 cm long, bright, glossy and leathery. Male and female flowers occur on separate trees. Commercial trees only have female flowers. Male trees occur but are rare. Flowers are produced on shoots near the end of branches. Female flowers have 4 cream petals and 4 sepals and a round ovary with a thick stigma composed of 4 - 8 lobes. No pollination is required because fruit develop without fertilisation. The fruit is dark blue with 4 - 8 white juicy segments inside the thick skin. Each segment is enclosed in fine pinkish veins. It is 8 cm across and has flower sepals attached to the stalk end. It leaks yellow sap from wounds. The fruit is produced on side branches near to the tips of the main branches. Fruit often have no seeds or up to 2. These are not true seeds but they will grow.



**Distribution:** Mangosteens are tropical plants and require a hot humid climate (25 - 35°C). Temperatures below 5°C will kill the plant and temperatures below 20°C slow growth. It grows from sea level to 1,000 m altitude in the equatorial tropics although it grows very slowly at higher altitudes. It can stand light shade but does not do well with sea breezes, and needs protection from wind. It cannot tolerate drought or alkaline soils. It needs fertile soil. Good fertility enables earlier bearing. It suits high rainfall areas over 2,500 mm per year. It suits hardiness zones 11 - 12.

**Use:** Fruit is best eaten fresh, (It is the layer around the seed). Can be cooked and used in desserts. Seeds are eaten after boiling or roasting and can be ground to produce a vegetable butter.

**Cultivation:** Trees breed true from seed because they are produced asexually. This means trees are very uniform. Some seeds can produce more than one seedling. They germinate between 10 and 54 days after planting. Seeds need to be fresh (less than 5 days after extraction), undamaged, and cleaned of pulp. They quickly lose their viability. Transplanting is done with care after 2 years. Long tap roots make this difficult. Young trees require shade during hot weather. The slow early growth is because the original roots of the seedling are replaced by new roots from the base of the stem. A spacing of 7 - 10 m suits. Grafting onto vigorous root stocks is difficult. It has been performed on *Garcinia tinctoria*. Budding, cuttings and layering have been unsuccessful.

**Production:** Trees are grow slowly and begin to bear after 8 - 20 years. Fruit are produced on shoots more than two years old so pruning is unnecessary. Fruiting is seasonal once or twice a year. Often more fruit are produced every second year. The main fruiting season is November to March. Fruit need to be harvested when mature and ripe. Fruit can only be transported with difficulty. They can be stored under refrigeration (10°C for up to eight weeks). Between 500 and 1,500 fruit are produced per tree. Fruit are best opened by cutting the skin around the middle to prevent tannins from the skin spoiling the flavour. Fruit can be stored for 3 weeks if undamaged.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	81.3	299	0.6	-	2.7	0.4	-

## Fruit

**English:** Pawpaw, Papaya

**Local:** Cây đu đủ

**Scientific name:** *Carica papaya*

**Plant family:** CARICACEAE

**Description:** A well known tropical fruit that grows 3 - 5 m tall and only occasionally has branches. The stem is softly woody and has scars from fallen leaves along it. There is a clump of leaves at the top of the plant. The leaves are large (50 cm wide) deeply lobed and on leaf stalks up to 90 cm long. Trees can be male, female or bisexual. Male flowers are small and white and on long stalks. Female and bisexual flowers are on short stalks. These have no fruit, round fruit and long fruit respectively. There are three forms of long fruit. Seeds are black.



**Distribution:** A tropical plant that grows from sea level up to about 1700 m altitude in the equatorial tropics. In cooler regions they have to be planted but in humid tropical regions are commonly self-sown. Sunlight allows germination when forest is cleared. Cannot stand frost. Needs a night temperature above 12°C and don't tolerate water-logging. Plants die after 48 hours in standing water. It needs a pH between 5 - 8 and suits hardiness zones 11 - 12.

**Use:** Fruit can be eaten ripe and raw. Green fruit can be cooked as a vegetable. The young leaves can be eaten cooked, but are bitter. The flowers and the middle of the stem can be eaten. Papayas contain papain which is a meat tenderiser. The dried seeds can be used as a spice.

**Cultivation:** Seeds grow easily and plants grow quickly. Fresh seeds can be used. If dry seeds are used they should be soaked before planting. Seeds should be sown when temperatures are 24 - 30°C and need a reasonably fertile soil. Seeds can be sown directly or the seeds can be put in a nursery and the seedlings transplanted. Seeds in a nursery should be sown 1 - 2 cm deep. Seedlings can be transplanted when they are about 20 cm high. Plants should be about 3 m apart. Continuous fruit production depends on fertility, temperature and moisture being adequate to maintain active growth. Fruit is produced year round but the growth and development rate decreases with temperature. Size and quality of fruit declines at lower temperatures. Pollination is by wind and insects and is not normally limiting. Normally cross and self-pollination both occur.

**Production:** Seeds emerge in 2 - 3 weeks. Vegetative growth before flowering is 4 - 8 months. One or more fruit grow per leaf axil, about every 1 - 2 weeks under good growing conditions. With good growth, 100 fruit can be produced from one plant in a year. Pollination to maturity is about 2 - 3 months. On the coast in tropical equatorial regions, pawpaws start producing fruit after about 4 - 5 months, but in the highlands this may take 12 - 18 months. The first fruit are ready 6 - 11 months from planting. Tree life is about 2 - 3 years, although they may live for 10 - 12 years.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	75.4	378	8.0	-	140	0.77	-
fruit	88.0	163	0.5	290	54	0.4	0.18
fruit (unripe)	92.1	109	1.0	-	-	0.3	-

## Fruit

**English:** Pineapple

**Local:** Cây dứa, cây thom

**Scientific name:** *Ananas comosus*

**Plant family:** BROMELIACEAE

**Description:** A perennial herb with a rosette of long, thick, spiky leaves, up to 1 m high and spreading 1 - 1.5 m. The leaves are arranged in spirals. Some kinds have thorns along the edges of the leaves. The plant produces suckers, both near the base of the stem and also higher up the stem near the fruit. These are called slips, or suckers, and are broken off and used for planting. The main plant dies after producing a fruit, but the suckers keep growing. The plant produces a flower and fruit at the end. The fruit is made up of about 150 berry-like fruitlets that are almost fused together. There is a small crown of leaves on top of the fruit. The fruit can be 25 cm long and weigh 0.5 - 4 kg. There are two main types of pineapples. The rough-leaved variety has spines on the leaves and produces a smaller but sweeter fruit. The smooth leaf variety has spineless leaves and larger fruit.



**Distribution:** The plant has been taken to most tropical and subtropical places. It is a tropical plant. It grows up to 1800 m altitude near the equator. It can survive brief periods down to freezing, but cold retards growth, delays fruiting, and causes fruit to be more acid. A loose, well-drained soil with high organic matter is best. It can survive drought, but adequate soil moisture is necessary for good fruit production. Pineapples need an annual average temperature of 17.2 - 26.9°C. Growth ceases below 20°C. In the equatorial tropics, this is mostly between sea level and 1800 m altitude. Pineapples need well-drained and fertile soil. It suits an acid soil and can develop roots in soils where lime has been added. The soil acidity can be between pH 3.3 - 6.0. The best range is pH 4.5 - 5.5. Soils which are not sufficiently acid can be treated with sulphur. It suits hardiness zones 9 - 10.

**Use:** The fruit is eaten fresh or used for juice. Unripe fruit are also cooked and eaten. The young, heart-leaves can be eaten. It is an attractive and popular snack food.

**Cultivation:** The suckers, slips, and the top of the fruit, can be used for planting. The time to maturity is the fastest for the suckers near the bottom of the plant and slowest when the top of the fruit is planted. Therefore, use suckers that grow from the stem near the ground for earliest yield. Pineapple flowering hormone can be used for fruit production with thorny varieties and calcium carbide for smooth varieties. Fruiting is less seasonal in the highlands than in the lowlands. Pineapples can be planted with 35,000 - 43,000 plants/ha or 3 - 4 plants per square metre. If plants are spaced more widely, they produce more suckers. Fruits become more acid where plants are closely spaced. If too many suckers are left growing from the main plant, then smaller fruit will be produced. They can grow in partial shade and, in this situation, the plants are normally more green. The red colouring of pineapple leaves is due to a deficiency of the nutrient nitrogen. This shows up more quickly in plants in full sunlight. When the plant is sufficiently large, it responds to changes, such as less available nutrients or water, and starts to produce a flower, then a fruit. The number of hours of sunlight, as well as reducing temperature and reduced sunlight, also help the flowers form. The result of this is that flowering and fruiting is often seasonal. This can easily be changed by using a fruiting hormone which allows fruit to be produced at times to suit the grower. Pineapples can grow in semi-arid conditions because the leaves can store some water. They also tend to lose only small amounts of water from evaporation through their leaves, but they can grow well with

plenty of water. The roots are very sensitive to water-logging, so the soil must be well-drained. Pineapples do not cover the soil well, so it is good to use a mulch of plant material to help weed control, provide some nutrients and to stop soil erosion.

**Production:** Plants usually produce for about 4 years. Fruiting is less seasonal in the highlands than in the lowlands in the tropics. The growth rate for pineapples slows at cooler temperatures. Plants grown in the highlands, or at higher latitudes, take longer to mature. It takes 60 days from when the flower starts to form until the fruit appears, then a further 5 months until the fruit is ready for harvest. The time from planting to harvesting ranges from 11 - 32 months, depending on temperature. The fruit are smaller, poorer shape and more acid where the temperatures are lower or there is less sunlight.

**Food Value:** Per 100 g edible portion

<b>Edible part</b>	<b>Moisture %</b>	<b>Energy kJ</b>	<b>Protein g</b>	<b>proVit A µg</b>	<b>proVit C mg</b>	<b>Iron mg</b>	<b>Zinc mg</b>
fruit	84.3	194	0.5	60	25	0.4	0.1

## Fruit


**English:** Rambutan

**Local:** Cây chôm chôm

**Scientific name:** *Nephelium lappaceum*

**Plant family:** SAPINDACEAE

**Description:** A medium sized tree 12 - 25 m tall. It keeps its leaves throughout the year. Trees spread to 6 m wide. The leaves are alternate with leaflets opposite along the stalk. There are 5 - 7 pairs of leaflets. Each leaflet can be 20 cm long by 8 cm wide. The leaflet stalks are short and the leaflets are pointed at the tip and rounded at the base. The flowers are very numerous on compound flower arrangements. Seedling trees can produce only female or only male trees. Some kinds have both male and female flowers or both flower parts in the one flower. The dark red fruit has many prominent projections united at the base. The colour can vary between yellow, green, orange and bright red. Some trees are separately male and female, and sometimes seedless fruit is set. Male and female flowers are often separate on the same tree. The fruit is up to 5 cm long in loose clusters of 10 - 12. It normally contains one large seed. There are several named cultivated varieties.



**Distribution:** A tropical plant that suits the humid tropical lowlands mostly within 12° of the equator. It grows from sea level up to about 500 m in equatorial zones and will grow about 17° from the equator. It does best where the temperature is constant around 28°C or ranges from 22 - 30°C with a rainfall of 2,000 – 5,000 mm per year. Low humidity can cause leaf scorch. Excessive rainfall before flower formation reduces flowering. A dry period of about one month assists flower initiation and areas with two seasonal dry periods can get double crops. A well drained soil is needed.

**Use:** The fruit are eaten fresh. Sour fruit are eaten stewed. The seeds are sometimes roasted and eaten, and can be used for oil.

**Cultivation:** Trees can be grown from fresh seeds. The short lived seeds need to be planted fresh, after washing. Germination takes place in 7 - 10 days. Seedlings are not easy to transplant. Roots are sensitive to drying out. Selected trees can be propagated by budding or air-layering. Seedling tree quality varies considerably due to cross pollination. Trees are planted 10 m apart at densities of 80 - 120 per ha. About 1% male flowers are sufficient if enough pollinating insects are present. A mixture of varieties might be needed to get good fruit set. Organic manure is often used twice yearly.

**Production:** Seedling trees can bear after 6 years and vegetative trees after 4 years. Flowering occurs during the dry season. The main fruit season is November to March. Fruit mature about 110 days after flowering. Where trees with only female flowers occur, male pollinator trees are needed. Fruit can weigh 20 - 40 g and need to be harvested when ripe. A well maintained tree can produce 150 kg.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit (raw)	82.0	271	1.0	0	53	1.9	0.1

## Fruit


**English:** Watermelon

**Local:** Dưa hấu

**Scientific name:** *Citrullus lanatus*

**Plant family:** CUCURBITACEAE

**Description:** An annual climber, with deeply divided leaves and tendrils along the vine. It trails over the ground and has hairy, angular stems. The leaves are on long leaf stalks. The leaves are deeply divided along their length. These leaf lobes are rounded and can themselves be divided. The leaves are 5 - 20 cm long by 2 - 12 cm across. The tendrils are divided. The plant has separate male and female flowers on the same plant. The flowers are pale yellow and smaller than pumpkin flowers. The flowers occur in the axils of leaves. The male flowers appear first. Fruit are large and round or oval. They can be 60 cm long. Fruit have a hard smooth skin. Several fruit colours and shapes occur. They often have a dark green mottle, or blotches. The fruit has reddish, juicy flesh and black or red seeds. The seeds are oval-shaped and smooth.



**Distribution:** It grows in most tropical and subtropical countries. It grows best on the coast in the tropics, but will grow up to about 1000 m altitude. It will not stand water-logging and does well on sandy soils. Plants are frost-sensitive. Seed will not germinate below 21°C. Temperatures between 24 - 30°C are suitable. Fruit are sweeter in arid warm areas. It suits hardiness zones 10 - 12.

**Use:** The fruit is eaten raw when ripe. Small, unripe fruit can be cooked as a vegetable. The skin is sometimes candied in vinegar and eaten with fish. Seeds are also eaten. They are dried, soaked in salt water, then roasted. Oil is extracted from the seeds. Very young leaves are occasionally eaten. It is a popular fruit.

**Cultivation:** They are suitable mainly for the dry season. A spacing of 1.5 - 2 m is suitable. They grow easily from seed. They do best when fully exposed to the sun. Seed can be dried and stored. If too much vegetative growth occurs, picking out the tip to produce side branches will produce more fruit.

**Production:** Harvesting commences after 4 - 5 months. The main fruit season is November to January. The ripeness can be determined by tapping the fruit to get a dull sound. The part of the fruit on the ground changes from green to light yellow and the tendril near the base of the fruit becomes dry when ripe. Fruit yield can be 45 - 60 t/ha.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	94.0	92	0.4	20	5	0.3	0.1
seed	5.1	2330	28.3	0	0	7.3	10.2

## Vegetables

**English:** Bitter cucumber  
**Local:** Khổ qua, mướp đắng

**Scientific name:** *Momordica charantia*  
**Plant family:** CUCURBITACEAE

**Description:** A slender annual climber with flowers of both sexes on the one plant. Simple tendrils and vines can be 4 m long. Bright green lobed leaves 5 - 12 cm long on thin leaf stalks 3 - 10 cm long. Flowers have a sweet smell and 5 small, yellow petals. Fruit are green when young and orange when ripe and have a lumpy appearance, with ridges along its length and when fully ripe burst open. Bright red covering on the seeds inside., which are pale brown and 10 - 16 mm long and 7 - 10 mm wide. Considerable variation in the fruit occurs between varieties.



**Distribution:** A tropical plant that grows from sea level up to about 500 m and will probably grow to 1,000 m altitude in tropical regions. They require a well drained soil preferably rich in organic matter. Seeds do not germinate below 15°C. Plants grow best with temperatures of 18 - 35°C. A soil pH of 6.5 is best. It suits hardiness zones 9 - 12.

**Use:** The young bitter fruit are cooked and eaten. They are boiled, stuffed, fried or pickled. They are used in soups, stews and stir-fried dishes. The seed mass of the ripe fruit is used as a food flavouring. The leaves are also cooked and eaten as a flavouring. The tender shoots and leaves are sometimes eaten. **Caution:** The raw leaves are considered to cause diarrhoea and vomiting.

**Cultivation:** Plants are grown from seed. For large scale plantings, 6 - 7 kg of seed are required for planting one hectare. Seeds are planted at 50 cm spacing in the place where the plants are to grow and need a stick to climb up. Often plants are grown on raised beds 2 m apart with 0.5 m between plants. The seed has a hard seed coat and germinates slowly. Soaking seeds for 24 hours before sowing gives a quicker more even germination. Regular watering is required.

**Production:** Fruit are ready to harvest 45 - 55 days after planting. Fruit should be harvested when young and tender. Once fruit have begun to change colour to yellow they are past maturity for eating. Early removal of young fruit also ensures continuous fruit setting. This can allow 6 - 8 successive pickings of fruit. Fruit on the plant are sometimes wrapped in paper to prevent fruit fly damage. Seed well stored can remain viable for 4 - 5 years. The young bitter fruit are cooked and eaten. The fruit is blanched or soaked in salt water to reduce the bitter taste.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	8.6	2020	18.6	-	-	-	-
leaf tip (boiled)	88.7	146	3.6	173	57	1.0	0.3
fruit	93.6	105	1.2	-	-	0.2	-
pod (boiled)	94.0	79	0.8	11	33	0.4	0.8
pod (raw)	94.0	71	1.0	380	84	0.4	0.8

## Vegetables

**English:** Chinese radish, Daikon

**Scientific name:** *Raphanus sativus var. longipinnatus*

**Local:** Củ cải trắng

**Plant family:** BRASSICACEAE

**Description:** A cabbage family plant like a large radish. It has a large fleshy root and deeply divided leaves. The taproot is long and white. It takes one or two years to complete its life cycle. The stems can be 20 - 100 cm tall. The leaves near the base are long, divided and have coarse teeth. The leaves up the stem are simple and narrow. The flowers develop at the top. They are usually white with purple veins. The fruit are pods about 3 - 8 cm long and 1.25 cm across. They have a long tapering beak. There are 6 - 12 round seeds. These can be yellow or brown.



**Distribution:** A temperate plant. It is more common in lowland areas in the tropics. It needs a rich loose fertile soil.

**Use:** The roots are cooked and eaten. They are used in stir-fried dishes and soups. They can be pickled. The young leaves can be eaten cooked. The roots can also be shredded and eaten raw in salads. The young flowers are stir fried or added to soups. The young seedlings can be cooked like spinach. The peppery seeds can be used in salads and rice dishes. The pods can also be eaten.

**Cultivation:** Plants are grown from seeds. A spacing of 15 cm apart in rows is suitable. Often plants are sown more thickly and seedlings are thinned out and eaten. It is possible to save seed from plants.

**Production:** Plants are ready for harvesting about 50 days after planting.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg

## Vegetables

**English:** Okra, Lady's fingers

**Local:** Đậu bắp, đậu tây

**Scientific name:** *Abelmoschus esculentus*

**Plant family:** MALVACEAE

**Description:** A tropical annual herb that grows erect, often with hairy stems. It mostly grows about 1 m tall but can be 3.5 m tall. It becomes woody at the base. The leaves have long stalks up to 30 cm long. Leaves vary in shape but are roughly heart shaped with lobes and teeth along the edge. Upper leaves are more deeply divided than lower ones. The flowers are yellow with red hearts. The fruits are green, long and ribbed. The seeds are 4 - 5 mm across. They are round and dark green.



**Distribution:** A tropical plant that suits the hot humid tropical lowlands but is unsuited to the highlands. It is very sensitive to frost. It can grow in salty soils. It grows best where temperatures are 20 - 36°C. It can grow well in dry climates with irrigation. It suits hot humid environments. It does best on well drained well manured soils but will grow on many soils. A soil pH of 5.5 - 7.0 is best.

**Use:** Pods are eaten cooked. They are slimy, but less so if fried. Dried powdered seeds can be used in soups as a thickener. They can also be pickled. Young leaves can be eaten cooked. They can be dried and stored. Flowers can also be eaten. Okra is frozen and canned. The seeds are roasted and used as a coffee substitute.

**Cultivation:** They are grown from seeds, which are easy to collect. They need high temperatures for germination (over 20°C) and a sunny position. Often seeds are soaked for 24 hours before sowing to give quick germination. Seeds are sown 1.5 - 2.5 cm deep with 2 - 3 seeds per hole. Later these are thinned out to one plant. Seeds can be sown in nurseries and plants transplanted. Pinching out the tops of plants when 30 cm high encourages branching. A spacing of about 90 x 45 cm is suitable. About 8 - 10 kg of seed are required for one hectare. Most kinds respond to fertiliser. Seeds do not breed true and can cross with other kinds of okra growing nearby. This is not normally a problem but simply means plants and fruit are not all the same.

**Production:** Plants maintain production if the fruits are harvested regularly. Plants are ready to harvest 8 - 10 weeks after sowing. Seed yields of 500 - 800 kg per hectare are recorded. Pod yields of 4 - 6 tonnes per hectare occur. It takes 2 - 4 months from sowing to harvest of young pods. Pods develop 5 - 10 days after flowering. Pod harvests can continue for 1 - 2 months. Leaving pods on the plants stops new pods developing.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	9.2	1721	23.7	-	-	-	-
leaf	81.0	235	4.4	116	59	0.7	-
pod (fresh)	88.0	151	2.1	185	47	1.2	-
fruit (cooked)	90.0	134	1.9	58	16.3	0.5	0.6

## Vegetables


**English:** Chinese broccoli, chinese kale

**Scientific name:** *Brassica oleracea* var. *alboglabra*

**Local:** Cải làn, cải rồ

**Plant family:** BRASSICACEAE

**Description:** A cabbage plant with a single fleshy stem. Although it keeps growing from year to year it is normally grown as an annual. It grows 45 cm high and spreads 40 cm across. The leaves are dark green and rounded on long stems. Plants start to flower when 10 leaves are present. Flowers are white but there are varieties with yellow flowers. There are several named cultivars.



**Distribution:** It does best in a fertile soil. The soil needs to be well drained. It prefers a soil pH of 6 - 7. Temperatures during the day of 18 - 28°C are best. It can tolerate frost. It grows well in tropical regions but cool temperatures are necessary for flowering.

**Use:** The flower stalk, flower heads, buds and tender leaves are all eaten. The stems are steamed or braised and often served with oyster sauce. They are also used in soups.

**Cultivation:** Plants are grown from seed. Seed can be sown direct or put in a nursery then transplanted. Seed is sown about 0.5 cm deep and germinate in 3 - 10 days. A spacing of 15 cm is suitable. Wide spacing causes stems to become thick and tough. Because plants are shallow rooted, they need regular watering.

**Production:** Chinese broccoli is fast growing. Flower heads are harvested after about 9 weeks. Heads are harvested individually to allow others to form. Harvesting is done before buds start to open.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
flower (cooked)	93.5	92	1.1	164	28.2	0.6	0.4

Picture sourced from <http://thedahliafarm.blogspot.com.au/2010/06/gai-lohn-or-chinese-broccoli-or-chinese.html>

## Vegetables

**English:** Eggplant, Aubergine

**Local:** Cà tím, cà dái dê

**Scientific name:** *Solanum melongena*

**Plant family:** SOLANACEAE

**Description:** A perennial shrubby herb up to 1 m tall. It is often grown as an annual. It has a deep taproot and branched side roots. The stem is thick and covered with many woolly hairs. The plant has many branches. Often the plant is spiny. Leaves can be 20 cm long and wavy along the edge. Leaves are covered with hairs. Flowers are bluish red and 5 cm across. They are either solitary or in small groups opposite the leaves. They have 5 large woolly lobes which continue to surround the base of the fruit. Fruit are white, blue, green or purple. The fruit colour and shape vary. Sometimes the fruit is spiny. Often the fruit are 10 - 20 cm long and 5 - 8 cm wide. Numerous kidney shaped seeds are in the flesh of the berry. There are many cultivated varieties.



**Distribution:** A tropical plant. Plants grow from sea level up to 2,200 m altitude in the tropics. It suits wet climates but does well in dry climates with irrigation. It needs a long warm growing period. A daily mean temperature of 20 - 30°C is most suitable. They are frost tender. They need a rich, friable, well tilled soil. In the sub-tropics they can be grown as a summer crop.

**Use:** Fruit are mostly fried then eaten. They can also be grilled, baked, stuffed and stewed. They are used in curries. The fruit are also dried and stored. The leaves, although edible, are hairy and not good flavor.

**Cultivation:** Plants are grown from seeds. Seeds germinate slowly. At the best temperature, they germinate in 8 - 12 days. Seed are sown in nursery beds. Seedlings can be transplanted when about 8 cm tall or 4 - 6 weeks old. Plants need to be about 60 - 100 cm apart. Because some cross pollination can occur, seed crops need to have varieties planted 400 m apart.

**Production:** Fruit are ready for harvest after 3 months. They continue to yield for 3 - 4 months.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	91.8	117	0.83	6	1.3	0.4	0.2
fruit (fresh)	93.4	62	0.7	50	5	0.4	0.3

## Vegetables

**English:** Pumpkin, Winter squash

**Local:** Bí ngô, bí đỏ

**Scientific name:** *Cucurbita moschata*

**Plant family:** CUCURBITACEAE

**Description:** A pumpkin family plant. It is a creeping plant with long creeping stems and softly hairy but without prickly hairs. The stems are rounded or 5 angled and moderately hard. They can grow 15 - 20 m long. The leaves are large and shallowly lobed and divided like fingers on a hand. Occasionally the leaves have white blotches. They have rounded lobes. They are 20 cm by 30 cm. The leaf stalk is 12 - 30 cm long. The flowers have male and female flowers separately on the same plant. The fruit stalk is distinctly expanded where it joins the fruit. The fruit are not



hard shelled and are dull in colour. The flesh is yellow and often has fibres through it. The seeds are plump and white to brown. They separate easily from the pulp of the fruit. The edge of the seed is scalloped and irregular in outline. There are a large number of cultivated varieties.

**Use:** The fruit are eaten cooked. They are boiled, fried or baked. They can be mashed and used in pies, soups, bread and cakes. They can be dried, ground into flour and used for bread. The young leaves and flowers are edible. They can also be dried and stored. The seeds are eaten roasted. They can also be roasted in salt.

**Distribution:** A tropical plant that suits the wet tropics. It will thrive in humid as well as in very hot climates. A temperature of 18 - 30°C is best. It can tolerate some shade. It can grow in soils with a pH of 5.5 - 6.9. It suits hardiness zones 8 - 11.

**Cultivation:** Plants are grown from seed. Seeds can be put in a nursery and transplanted.

**Production:** Fruit mature in 70 - 180 days after sowing depending on variety.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	5.5	2331	23.4	-	-	2.8	-
leaf	93.6	88	3.0	95	10	2.1	-
fruit	95.0	35	0.7	-	14	0.4	-

## Vegetables

**English:** Wax Gourd

**Local:** Bí đao, bí xanh

**Scientific name:** *Benincasa hispida*

**Plant family:** CUCURBITACEAE

**Description:** A climbing, pumpkin family plant. The vine can grow to 3 m long. The plant re-grows from seed each year. The vines are thick, furrowed and hairy. The leaves are heart shaped with 5 - 7 lobes. They are rough to touch. Flowers are yellow. The immature fruit can have skin of various colours depending on variety. The fruit is up to 30 cm long and 20 cm across and green with a waxy covering when mature. This waxy layer enables the fruit to be stored for a long time. Fruit shape and size can vary with variety. The flesh is firm and white. The fruit are heavy, weighing from 8 - 45 kg.



**Distribution:** A tropical plant. It is suited to warm, lowland, tropical conditions. It does better in dry areas or drier seasons. They are reasonably drought tolerant. The best temperature for growing is 23 - 28°C. They need a well-drained soil. They grow best with a soil pH of 6.5.

**Use:** The white flesh is added to stir fried dishes. The seeds can be fried and eaten. Young leaves and flower buds can be eaten. The young fruit are used as a vegetable. The mature fruit are peeled, cut in pieces and candied.

**Cultivation:** It is grown from seeds. There are about 1,800 seeds per kg. Seeds are sown 3 - 5 cm deep with a spacing of 60 - 80 cm between plants. Seed can be sown in nurseries and transplanted when 15 - 20 cm tall. They are usually planted in mounds and allowed to grow over a strong trellis. They can be allowed to stay on the ground. If plants are going to be allowed to trail over the ground, a spacing of about 3 m is necessary. Decayed manure or compost is used where available. Plants are responsive to sulphate of ammonia. Flowering normally starts 60 - 80 days after planting. Flowers are open in the early morning. Hand pollination may assist fruit development. This becomes more important in colder areas. Thinning of fruit gives larger fruit. The growing tips of plants can be pruned out to encourage branching or to restrict growth.

**Production:** Fruit are ready 3 - 5 months after planting. The fruit keeps well when fully mature. They can be stored for 6 months at 13 - 15°C in a dry atmosphere. The pulp of wax gourds can be shredded and dried for later use.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	92.0	118	0.7	0	15	0.6	0.6
fruit (cooked)	96.6	54	0.4	0	10.5	0.4	0.6

## Vegetables

**English:** Chinese water chestnut

**Local:** Củ ấu, năn ngọt

**Scientific name:** *Eleocharis dulcis*

**Plant family:** CYPERACEAE

**Description:** A herb which grows in water. It is a tufted sedge with round green stems. The bases are covered with brown sheaths. The stem is about a metre high and 1 cm across. It grows 30 - 200 cm high. From the top of each planted corm, several slender horizontal rhizomes radiate out into the mud, each terminating in a corm. The edible part consists of a flattened corm. The rhizome is short. Under the ground there are stolons bearing tubers. The tubers are almost round and have 4 - 6 distinct rings. They are usually about 1 cm across but can be up to 4 cm across. They are dark brown. The stems are tufted and slender. There are fine lines along the stems. The purplish leaves are reduced to thin tube like sheaths. Each plant produces these long tubular leaves that project above the water surface. The flower spike is on the end of the plant. There are many flowers, 1.5 - 6 cm long by 3 - 6 mm wide. The fruit is a nut 1.5 - 2 mm long.



**Distribution:** It suits humid, monsoonal, tropical and subtropical locations. It is found in open wet places and shallow water. It grows in fresh water swampy grounds or in shallow water. It is also found in rice fields. It needs at least 220 frost free days. It needs a soil temperature above 15.5°C for germination of the corms. It needs a pH of 6.9 - 7.3. It can be grown up to 1,200 m altitude and suits plant hardiness zones 9 - 12.

**Use:** The tubers are cooked and eaten. The corms can be eaten raw, roasted or boiled after they have been peeled. Normally, they are cut into small slices and added to soup or to fish and meat dishes. They can be sweetened for desserts.

**Cultivation:** Plants can be grown by division or tubers. They are put in holes 20 - 30 cm deep. Fields are flooded after planting then allowed to drain. When top growth is 20 - 30 cm high fields are flooded to at least 10 - 12.5 cm. A spacing of 75 cm x 75 cm is suitable. 500 kg of corms per hectare are required for planting.

**Production:** Corms mature after 7 - 8 months. Yields of 20 - 40 t per ha are possible.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
corm	50.8	635	3.7	-	52	15	1.9
fruit	79.6	268	1.4	0	5	0.7	0.5

## *Nuts, seeds, herbs and other foods*

**English:** Cashew

**Local:** Đào lộn hột, cây điều

**Scientific name:** *Anacardium occidentale*

**Plant family:** ANACARDIACEAE

**Description:** An evergreen tree, with spreading branches, growing 7 - 14 m tall. The canopy can spread to 12 m. The roots grow deeply and spread widely. The shiny leaves are pale green and large. They are 10 - 15 cm long by 6 - 8 cm wide. They have fine veins. The flowers are produced on the ends of the branches. They are red in colour. The kidney-shaped nut is about 3 cm long and is borne below the "apple" which is really a fleshy stalk.



**Distribution:** It is a tropical plant that suits the lowland tropics but will grow up to about 1,200 m altitude. It only bears well in dry areas because of blight of the flowers. It grows with temperatures between 22 - 26°C. A rainfall of 1750 mm per year is considered suitable but good yields have been obtained with rainfall of 750 mm. It can grow on poor soils but needs good drainage.

**Use:** The fleshy "apple" is edible but acid until very ripe. It is used for jams and drinks. It is also candied, made into chutney and pickles. The nut is eaten after roasting. The young shoots and leaves are edible. They are picked during the rainy season and eaten fresh with hot and spicy dishes. **Caution:** The oil of the nut can blister the skin until roasted. The apple is used to make spirits.

**Cultivation:** It is usually grown from seeds. Seeds germinate poorly and slowly. Only nuts which sink in water (or a solution of 150 g of sugar in a litre of water) should be planted. Seeds are sun dried for 2 - 3 days to improve germination. Seeds can be sown in a nursery then transplanted, or more commonly, are sown directly. Trees are spaced 7 - 10 m apart. The crop is cross pollinated mostly by insects. For good production, complete fertiliser or appropriate organic material should be applied. Pruning to shape the tree is often undertaken in the first 2 - 3 years. Cashews are often planted scattered in gardens or amongst other trees. Clearing under the tree prevents fire and makes finding nuts easier. Allowing nuts to fall before harvesting ensures only ripe nuts are collected. Resin in the cashew nut shell can damage hands and discolour the nuts. Roasting the nuts before removing the kernel avoids this.

**Production:** Trees commence bearing after 3 years. Fruit production is seasonal, normally October - January. Mature nuts are produced in 2 - 3 months. Yields of 80 - 200 kg of nuts per hectare are normal. Trees reach maximum production after 10 years and trees last for about 100 years.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
nut	4.0	2478	17.5	-	-	2.8	4.8
leaf	69.9	418	5.2	-	-	-	-
fruit	84.7	213	0.8	0.12	265	1.0	0.2

## Nuts, seeds, herbs and other foods

**English:** Coastal almond

**Local:** Cây bàng

**Scientific name:** *Terminalia catappa*

**Plant family:** COMBRETACEAE

**Description:** A large tree, up to 25 - 40 m tall. It loses its leaves during the year. The trunk can be straight or twisted. There can be buttresses up to 3 m tall. The branches lie horizontally and come out in layers. The leaves are long, smooth and shiny, with an abrupt point at the tip and a rounded base. Leaves tend to be near the ends of branches. Leaves can be 17 - 29 cm long and 10 - 15 cm wide. Young leaves have soft hairs. The leaves turn red and fall off twice a year. Flowers are greenish-white and in a spike at the end of the branches. The lower flowers on a spike are female, and the others are male. The fruit is about 6 cm long by 3 - 4 cm wide, thick and flattened, with a flange around the edge. The fruit are green and turn red when ripe. The pulp is edible.



**Distribution:** It grows on beaches in almost all tropical countries in the world, including Solomon Islands. It is a tropical plant, and sometimes cultivated as a shade tree. The tree is common in lowland areas particularly on sandy or rocky beaches. Seeds are spread by bats and sea water, as well as being planted by people. It is common along streets in coastal towns. It will grow from sea level up to about 800 m altitude. Plants are frost-susceptible. It can tolerate drought. It suits hardiness zones 11 - 12.

**Use:** The kernel of the fruit is eaten raw. An edible oil can also be extracted.

**Cultivation:** Plants can be grown from seed. Seeds can be stored dry for a year or more. Seeds germinate freely and most seeds grow. Insects can badly damage the leaves of young seedlings.

**Production:** It is fast growing. Nut production is seasonal.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
nut (fresh)	31	1810	15.9	-	4	4.6	4.9
nut (dry)	4.2	2987	20.0	-	2	6.3	8.8

## Nuts, seeds, herbs and other foods

**English:** Peanut, Groundnut

**Local:** Đậu phộng, Lạc

**Scientific name:** *Arachis hypogea*

**Plant family:** FABACEAE

**Description:** Peanuts grow on spreading bushy plants up to about 40 cm high. The leaves are made up of 2 pairs of oppositely arranged leaflets. Flowers are produced in the axils of the leaves. Two main kinds of peanuts occur. The runner kind (Virginia peanut) has a vegetative or leafy branch between each fruiting branch and therefore produces a spreading bush. The bunch type (Spanish-Valencia peanuts) produces fruiting branches in a sequence one after the other along the branches. They grow as a more upright plant and grow more quickly. Pods are produced on long stalks which extend under the ground and they contain between 2 - 6 seeds. The stalk or peg from the flower grows down into the soil and then produces the pod and seed under the ground. The flowers need to be no more than 18 cm from the soil surface for the seed pod to develop underground.



**Distribution:** Peanuts grow in tropical and subtropical areas. They grow well from sea level up to about 1,650 metres in the equatorial tropics. They require temperatures of 24 - 33°C. Plants are killed by frost. They need a well drained soil and cannot stand water-logging and often require raised garden beds. Peanuts need 300 - 500 mm of rain during the growing season. Dry weather is needed near harvest.

**Use:** The seeds can be eaten raw or cooked. They are boiled, steamed, roasted, salted or made into peanut butter or flour. The young leaves and unripe pods are edible after cooking. Sprouted seeds can be eaten. An edible oil is extracted from the seeds. The remaining meal can also be eaten.

**Cultivation:** Peanuts require soil with good levels of calcium and boron or they produce empty pods. Peanuts have nitrogen fixing root nodule bacteria and therefore can give good yields in soils where nitrogen is low. The nuts are normally removed from the shell before planting and are sown 2 - 3 cm deep, with 10 cm between plants and 60 - 80 cm between rows. The soil needs to be weeded and loose by the time the flowers are produced to allow the peg for the seed pods to penetrate the soil.

**Production:** Flowering can commence in 30 days and it takes 3.5 - 5 months until maturity. Peanuts are harvested by pulling out the plant when the top of the plants die down. After harvesting, they should be left to dry in the sun for 3 - 4 days. Virginia peanuts have a longer growing season and the seeds need to be stored for 30 days before they will start to re-grow.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	4.5	2364	24.3	-	-	2.0	3.0
seed (fresh)	45	1394	15	-	10	1.5	-
leaf	78.5	228	4.4	-	-	4.2	-

## Nuts, seeds, herbs and other foods

**English:** Safflower

**Local:** Cây rum, hồng hoa

**Scientific name:** *Carthamus tinctorius*

**Plant family:** ASTERACEAE

**Description:** An erect, annual herb that grows to 60 - 150 cm tall. It has many branches. It has spines but the numbers vary. The stems are white, stiff and round with fine grooves along their length. The types with more spiny leaves are better for oil production. The leaves are arranged in spirals around the stem. They do not have leaf stalks. The leaves are dark green and glossy. They are 10 - 15 cm long and 2 - 4 cm wide. The flower head is made up of many small flowers that are 13 mm long and like tubes. They are yellow to orange in colour. The fruit is 4 angled and has a hard hull and a single white or grey seed. The seed is oblong.



**Distribution:** It grows in both tropical and temperate zones. It does better in drier regions. It cannot tolerate waterlogging. It does not suit the low, wet tropics. It needs a good dry season for drying. It is resistant to drought. It can stand some wind and salinity. High temperatures can result in poor seed set. It does best where temperatures are 17 - 20°C on average. At the equator it can grow at 1,600 – 2,000 m altitude but most commonly in other regions it grows below 900 m altitude. A soil pH of 5 - 8 is suitable. It can grow in arid places.

**Use:** The young shoots and leaves are eaten cooked or raw. They can be seasoned with soy sauce. The seeds are hulled and roasted. They are eaten as snacks. They are also used in chutneys. The seed oil is used in cooking and as a salad oil. (This can be done by boiling the seeds and floating off the oil). The dried, edible petals are used to colour foods. They can give red or yellow dyes. The slightly bitter petals can be cooked with rice.

**Cultivation:** Plants are grown from seed. A fine seed bed is required and seed are broadcast or drilled. It is best sown about 2 - 3 cm deep. Seeds germinate in 4 - 7 days and a soil temperature of 15°C is best. Plants should be topped as soon as the first buds appear to increase the number of flower heads. A spacing of 15 - 30 cm between plants is suitable. Wider spacing gives more heads per plant and closer spacing gives higher yields per area. A seeding rate of 20 - 30 kg per hectare is required. Crops respond to fertiliser if there is sufficient moisture. In very dry weather, harvesting in the more moist morning or evening avoids seed shattering. Plants are uprooted and heaped for a few days before threshing.

**Production:** Plants take 120 days to maturity. Seeds are ripe about 35 - 40 days after maximum flowering. Plants are harvested when leaves turn brown.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	5.6	2163	16.2	5	0	4.9	5.5

## Nuts, seeds, herbs and other foods

**English:** Sesame, Oil-seed sesame

**Local:** Mè, vùng

**Scientific name:** *Sesamum indicum*

**Plant family:** PEDALIACEAE

**Description:** A small, upright annual plant. It is erect and very branched and grows 1 - 2 m tall. The stem is stout, 4 sided and furrowed along its length. It is densely covered with fine, downy, glandular hairs that vary in shape. The lower leaves have long stalks and are spear shaped, often with lobes or a toothed edge. The leaf stalks are 3 - 11 cm long. The leaf blade is 4 - 20 cm long by 2 - 10 cm wide. Upper leaves are narrow and oblong. They are 0.5 - 2.5 cm wide. The flowers are pink and white. They occur in the axils of upper leaves, either on their own, or in



groups of 2 or 3. They can be white, pink, purplish and with yellow spots and stripes. The fruit can be smooth or rough and there are 2 chambers in the capsule. The fruit are brown or purple. They are oblong and deeply grooved. The seeds are small and oval. They are 3 mm by 1.5 mm and vary in colour from white, yellow, grey, red, brown or black. The fully ripe pods burst open.

**Distribution:** A tropical plant that suits the hot, dry, semi-arid tropics and sub-tropics. It can tolerate short periods of drought once established. It needs a temperature of 20 - 24°C in early growth then 27°C for ripening. It grows from sea level to about 1,200 m in areas with an annual rainfall of 400 - 1000 mm. Soils need to be well drained. It is very intolerant of water-logging. It cannot stand high humidity and needs frost free conditions. It needs a dry period for seed drying. It does not like acid soils. It grows in open sunny places. It can grow in arid places.

**Use:** The seeds are eaten. They are used in soups or fried or boiled. They are used in tahini and hummus. Seeds are eaten in the form of sweetmeats. Roasted seeds are used in pickles. They are also put on bread. Oil from the seeds is used in cooking and on salads. The refuse from the seed after the oil has been extracted is boiled in water and made into soup.

**Cultivation:** Plants are grown from seed. Seed will not germinate below 21°C. Seeds are broadcast on well prepared land and then harrowed in using feet or a light harrow. Plants can be thinned or weeded during early growth to produce a better crop. Seeding rates of 9 - 11 kg/ha are used. Plants are spaced 2 - 15 cm apart and in rows placed at 20 - 45 cm apart. Some varieties shatter easily.

**Production:** Yields of 340 - 500 kg/ha are average. Plants reach maturity in 80 - 180 days. Crops are harvested as the leaves begin to drop. Plants are cut and stooked or dried in racks. The hull is removed by soaking in water overnight, then partly dried and rubbed against a rough surface.

**Food Value:** Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	4.7	2397	17.7	1	0	14.6	7.8
leaf	85.5	188	3.4	-	-	-	-

## *Nutritional values of food plants by plant Family*

Plant Family	Scientific name	Common name	Edible part	Moisture %	Energy kJ	Protein g	Vit A µg	Vit C mg	Iron mg	Zinc mg	Page
Amaranthaceae	<i>Amaranthus tricolor</i>	Amarants	leaf	91.7	96	2.5	292	43.3	2.3	2.3	26
Anacardiaceae	<i>Mangifera indica</i>	Mango	fruit	83	253	0.5	1200	30	0.5	0.5	35
Anacardiaceae	<i>Anacardium occidentale</i>	Cashew	nut	4	2478	17.5	-	-	2.8	2.8	51
Araceae	<i>Amorphopallus paeonifolius</i> var. <i>campanulatus</i>	Elephant foot yam	tuber	78	340	2	0	6	2.4	2.4	13
Araceae	<i>Colocasia esculenta</i>	Taro	root	66.8	1231	1.96	3	5	0.68	0.68	17
Asteraceae	<i>Carthamus tinctorius</i>	Safflower	seed	5.6	2163	16.18	5	0	4.9	4.9	54
Basellaceae	<i>Basella alba</i>	Indian spinach	leaf	85	202	5	56	100	4	-	29
Brassicaceae	<i>Raphanus sativus</i> var. <i>longipinnatus</i>	Daikon									44
Brassicaceae	<i>Brassica juncea</i>	Indian mustard	leaf	92	108	2.4	31	73	2.7	-	28
Brassicaceae	<i>Brassica oleracea</i> var. <i>alboglabra</i>	Chinese broccoli	flower (cooked)	93.5	92	1.1	164	28.2	0.6	0.6	46
Brassicaceae	<i>Brassica oleracea</i> var. <i>acephala</i>	Kale	leaf (raw)	84	210	3.3	70	120	1.7	0.3	30
Bromeliaceae	<i>Ananas comosus</i>	Pineapple	fruit	84.3	194	0.5	60	25	0.4	0.4	39
Cactaceae	<i>Hylocereus undatus</i>	Dragon fruit	fruit	89.4		0.5		25	0.4		33
Caricaceae	<i>Carica papaya</i>	Pawpaw	fruit	88	163	0.5	209	54	0.4	0.18	38
Clusiaceae	<i>Garcinia mangostana</i>	Mangosteen	fruit	81.3	299	0.6	-	2.7	0.4	-	37
Combretaceae	<i>Terminalia catappa</i>	Coastal almond	nut (dried)	4.2	2987	20	2	2	6.3	6.3	52
Convolvulaceae	<i>Ipomoea batatas</i>	Sweet potato	tuber (baked)	72.9	431	1.7	2182	24.6	0.5	0.3	15
Convolvulaceae	<i>Ipomoea aquatica</i>	Kangkong	leaf (boiled)	92.9	84	2.1	520	16	1.3	1.3	31
Cucurbitaceae	<i>Benincasa hispida</i>	Wax gourd	fruit	92	118	0.7	0	15	0.6	0.6	49
Cucurbitaceae	<i>Momordica charantia</i>	Bitter cucumber	fruit	93.6	105	1.2			0.2		43
Cucurbitaceae	<i>Cucurbita moschata</i>	Pumpkin	seed	5.5	2331	23.4	-	-	2.8	-	48
Cucurbitaceae	<i>Citrullus lanatus</i>	Watermelon	seed	5.1	2330	28.3	0	0	7.3	7.3	42
Cyperaceae	<i>Eleocharis dulcis</i>	Chinese water chestnut	fruit	79.6	268	1.4	0	5	0.7	0.5	50
Dioscoreaceae	<i>Dioscorea bulbifera</i>	Potato yam	tuber	70.8	357	2.7	-	78	3.1	3.1	14
Durionaceae	<i>Durio zibethinus</i>	Durian	fruit	61.1	602	2.5	5	24	0.7	0.7	34
Euphorbiaceae	<i>Manihot esculenta</i>	Cassava	tuber	62.8	625	1.4	30	15	0.23	0.48	11
Fabaceae	<i>Acacia farnesiana</i>	Sweet Acacia	seed (dry)	8.1	1522	36.6	-	-	6	6	21
Fabaceae	<i>Canavalia ensiformis</i>	Jack bean	seed	10	1423	20.4	160	0	4.9	-	20
Fabaceae	<i>Vigna radiata</i>	Mung bean	seed	11	1432	22.9	55	4	7.1	-	19
Fabaceae	<i>Cajanus cajan</i>	Pigeon pea	seed	10	1449	19.5	55	-	15	-	22
Fabaceae	<i>Glycine max</i>	Soybean	seed	9	1701	33.7	55	-	6.1	-	23
Fabaceae	<i>Psophocarpus tetragonolobus</i>	Winged bean	seed	8.5	1764	41	0	-	15	15	24
Fabaceae	<i>Arachis hypogea</i>	Peanut	seed (dry)	4.5	2364	24.3	0	-	2	2	53

Plant Family	Scientific name	Common name	Edible part	Moisture %	Energy kJ	Protein g	Vit A µg	Vit C mg	Iron mg	Zinc mg	Page
Fabaceae	<i>Vigna unguiculata</i> <i>subsp. sesquipedalis</i>	Yard long bean	mature seed (raw)	8.4	1453	24.3	18.5	1.6	8.61	3.5	25
Malvaceae	<i>Abelmoshus esculentus</i>	Okra	pod (fresh)	88	151	2.1	185	47	1.2	-	45
Moringaceae	<i>Moringa oleifera</i>	Horseradish tree	leaf	76.4	302	5	197	165	3.6	-	27
Musaceae	<i>Musa sp (A &amp;/or B genome) cv.</i>	Banana	fruit	65.3	510	2	113	18.4	06.	0.1	32
Pedaliaceae	<i>Sesamum indicum</i>	Sesame	seed	4.7	2397	17.7	1	0	14.6	14.6	55
Sapindaceae	<i>Nephelium lappaceum</i>	Rambutan	fruit	82	271	1	0	53	1.9	1.9	41
Solanaceae	<i>Solanum melongena</i>	Eggplant	fruit (fresh)	93.4	62	0.7	50	5	0.4	0.4	47