

# Potentially Important Food Plants of Liberia



FOOD PLANT SOLUTIONS  
ROTARY ACTION GROUP  
Solutions to Malnutrition and Food Security

A project of the Rotary Club of  
Devonport North and District 9830

[www.foodplantsolutions.org](http://www.foodplantsolutions.org)



# Potentially Important Food Plants of Liberia

## **Dedication**

This book is dedicated to the 3 billion hard working farmers and families around the world who cultivate these, and other, food plants for their own subsistence, and who help conserve them in their rich diversity for other people to enjoy.

## Preface

This guide is based on information from the Food Plants International (FPI) database developed by Tasmanian agricultural scientist Bruce French. The source material and guidance for the preparation of the book has been made possible through the support of Food Plants International, the Rotary Clubs of District 9830, particularly the Rotary Club of Devonport North who founded Food Plant Solutions, (previously the Learn2Grow project), and many volunteers who have assisted in various ways.

The selection of plants included in this guide has been developed by Lyndie Kite working in a voluntary capacity using the selection criteria developed by Food Plant Solutions. These selection criteria focus on the local plants from each of the main food groups with the highest levels of nutrients important to human nutrition and alleviation of malnutrition. It is intended as a **Draft Guide only** to indicate some important food plants that serve as examples for this purpose. Other important nutritious plants may be equally useful, and it is recommended that the FPI database be used to source information on the full range of plants known to occur in Liberia. This guide has been developed with the best intention to create interest and improve understanding of the important local food plants of Liberia, and on the understanding that it will be further edited and augmented by local specialists with appropriate knowledge and understanding of local food plants.

Food Plant Solutions was initiated by the Rotary Club of Devonport North to assist in creating awareness of the edible plant database developed by Food Plants International, and its potential in addressing malnutrition and food security in any country of the world. In June 2007, Food Plant Solutions was established as a project of Rotary District 9830, the Rotary Club of Devonport North and Food Plants International. The primary objective of the project is to increase awareness and understanding of the vast food resource that exists in the form of local plants, well adapted to the prevailing conditions where they naturally occur, and how this resource may be used to address hunger, malnutrition and food security. For more information, visit the website [www.foodplantsolutions.org](http://www.foodplantsolutions.org). More detailed or specific information on plants, including references to material by other authors, is available on DVD on request.

**Disclaimer:** This Field Guide has been produced using information from the “Edible Plants of the World” database compiled by Bruce French of Food Plants International. Although great care has been taken by Food Plants International and Food Plant Solutions, neither organisation, or the people involved in the compilation of the database or this Field Guide:

- makes any expressed or implied representation as to the accuracy of the information contained in the database or the Field Guide, and cannot be held legally responsible or accept liability for any errors or omissions
- can be held responsible for claims arising from the mistaken identity of plants or their inappropriate use
- assume responsibility for sickness, death or other harmful effects resulting from eating or using any plant described in the database or this Field Guide

Always be sure you have the correct plant, and undertake proper preparation methods, by consulting with specialist scientists or local users of the plant. The Food Plants International database, from which the information in this Field Guide is drawn, is a work in progress and is regularly being amended and updated.

## Contents

|   |    |
|---|----|
| INTRODUCTION .....                                      | 1  |
| STARCHY STAPLES.....                                    | 11 |
| LEGUMES.....  | 21 |
| LEAFY GREENS .....                                      | 29 |
| FRUIT.....  | 36 |
| VEGETABLES .....  | 47 |
| NUTS, SEEDS, HERBS AND OTHER FOODS .....                | 55 |
| NUTRITIONAL VALUES OF FOOD PLANTS BY PLANT FAMILY ..... | 60 |



## **Introduction**

Potentially Important Food Plants of Liberia has been produced to provide information on approximately 40 edible plants that are known to grow in Liberia. These plants come from all the major food groups and have been chosen because of their high nutritional value. Many of the plants in this book may be neglected and under-utilised plants. This means they may not be well known. However, because they are high in many beneficial nutrients, and they are already adapted to the environment, and therefore likely to require minimal inputs, they could be important food plants that are likely to be superior to imported foods and plants. Commercially grown plants may also be included in the book, but only if they are significant foods for household consumption. It is hoped people will become confident and informed about how to grow and use these plants as many local food plants provide very good quality food.

## **Growing food**

Growing food to feed a family is, without doubt, one of the most important things anyone can do. The more interest you take in your garden and the more you learn about plants and how to grow them well, the more interesting and fun food gardening becomes.

## **A country with very special plants**

The local food plants of most countries have not been promoted and highlighted in the way they deserve. Visiting a local food market will quickly show what a rich variety of food plants can be grown in this country. Good information about these plants is often still in the minds and experience of local farmers, and has not been written down in books. This can make it hard for the next generation of young people to find out how to grow them.

In many countries, some of the traditional food plants are only harvested from the wild and others are only known in small areas. Others have hundreds of varieties and are the main food for people in different regions. Information on all these plants, their food value and the pest and diseases that damage them is available in the Food Plants International database.

## **Getting to know plants**

People who spend time in gardens and with their food plants get to know them very well. It is a good idea to learn from someone who grows plants well. Each plant grows best in certain conditions and there are often special techniques in getting it to grow well. For example, sweet potato will not form tubers if the soil is too wet, but it may still grow lots of green leaves. Taro will grow in light shade, but sweet potato will not. Ginger can grow in fairly heavy shade. Pruning the tips of betel leaf or pepper vines will cause more side branches to grow and therefore, produce more fruit. Stored yam tubers need special treatment if you want them to put out shoots early. There are lots of unique things about every plant and learning about these helps a good gardener produce more food.


## **Naming of plants**

Many food plants have local names, as well as a common English name. Every type of plant also has its own scientific name. Although the scientific name might not be widely recognised, this is the link by which people in different countries and with different languages can recognise the same plant. We know that many plants are grown in many different countries, but relying on local or common names, we might not recognise the same plant grown in different places. By using scientific names to accurately identify plants, we can get useful information from people in other countries. Wherever possible, plants in this book are named by their common English name and their scientific name.

### Local food plants are often very good

People sometimes think that local food plants are not very special and that any food plant that is new or comes from another country must be a lot better. This is often not true. Many of the newer or introduced food plants, such as the round or ballhead cabbages, have very little food value. Many traditional tropical green, leafy vegetables and ferns have 10 times or more food value as ballhead cabbage or lettuce. It is important to find out more information about the food value of different foods if we want to eat well. Citrus fruit, such as lemons and oranges, are often grown for vitamin C that helps keep people healthy. These fruits do not grow well in the tropics-the common guava fruit has three times as much vitamin C and is loved by children. This is just one example that there are often much better choices of local foods with higher levels of important nutrients.

Our bodies need a variety of food plants to enable us to grow, stay healthy and have enough energy to work. Different foods are needed to provide energy, protein, vitamins and minerals. The following diagram highlights the iron content value of some traditional edible, tropical plant leaves, compared with cabbage. Iron is a nutrient that is very important for our bodies and especially our blood. People who are short of iron become anaemic and lack energy.



The relative iron content of some edible leaves

### A healthy balanced diet

Good nutrition, or eating a healthy balanced diet, is really very simple. If people eat a wide range of food plants, their bodies will normally get a balanced amount of all the different nutrients they require. If a nutrient is lacking in one food plant, then they are likely to get it from another plant if they are eating a range of food plants. For this reason, everybody should eat a range of different food plants every day. The food group that is especially important for young people is the dark green leaves. Everyone should eat a good serving of dark green leaves every day. They have many vitamins and minerals, as well as protein. There are many spices or flavouring plants that can improve the taste of foods, but taste should be considered separately from food value.

### Learning to cook well

Even though some nutrients in food can lose some of their value during cooking, it is normally much safer to cook all food plants, at least for a short time. Bacteria, which cause diarrhoea, can occur in gardens and on food plants. These are killed during cooking. Many plants in the tropics develop cyanide, a chemical that makes them bitter and poisonous. This happens often with cassava (tapioca, manioc) and beans, but can also occur in many other plants. Boiling the food for two



minutes normally destroys cyanide and makes the food safe to eat. Some of the nutrients our bodies need (such as vitamin A for good eyesight) only become available when food is cooked in oil.

### **Learning to grow “wild” food plants**

Many plants grow wild in the bush and are not cultivated by people. We can normally find someone who has taken an interest in them and has learned to grow them. This may be people from a different language group. It may be that in their area they have found better types than the ones that simply grow wild.

### **Saving better types of plants**

If we simply allow plants to grow from seed, the improvements that have been made in finding sweeter or better types may get lost. Some fruit trees are like this and the fruit produced may not be sweet at all. It is often necessary to take cuttings from a tree to be sure the new plant is exactly the same as the old one. If the plants won't easily grow from cuttings simply by sticking a piece of the branch in the ground, there are other ways of helping these plants to form roots and start to grow. One good way is to make a small cut in the bark of a young branch and then wrap soil around the cut and cover it with plastic. With plants like guava, new roots will start to grow from this cut and grow into the soil wrapped around the branch. It can then be cut off and planted. This is called air-layering. A similar method is used with the roots of breadfruit. A shallow root is uncovered and a small cut made from which a new sucker will start to grow. This can be cut off and replanted.

### **Growing from cuttings and suckers**

Many food plants are grown from cuttings and suckers. This is very important, as it allows all the different kinds of yams, taros, bananas, sweet potato and sugarcane to be continually grown and ensures the varieties are preserved. Each plant has its own special propagation method. It is important to use healthy planting material, as diseases can be spread in planting material.

### **Saving seed**

Some food plants are grown from seed. Sometimes this is very easy as the seeds are large, store well, grow easily and grow the same as the original plant. It is more difficult with other plants. Many large fleshy seeds, such as breadfruit, need to be planted while still fresh as they do not store easily. Other seeds do not “breed true” or do not grow into new plants that are the same as the original plants. For example, the fruit may not be as large or sweet or have the same colour or taste. With many of these plants, it may be necessary to find ways of growing them from cuttings or other methods such as grafting. Some plants “inbreed” and get smaller or poorer. This happens when a plant self-pollinates or receives pollen from a close relative. Corn grown in small plots normally does this and the plants grown from seed grown in this situation get smaller and smaller each year. The seed needs to be saved from several different plants with different history and then mixed together before sowing. All the seeds on one cob are related and will inbreed. Some seeds develop a hard seed coat and need to be scratched, soaked in water, or even put into hot water, before they will start to grow. Saving local seeds is often a good idea as they are already adapted to local conditions. For example, seed saved from pumpkins grown locally will produce plants with less pest and disease damage than those grown from imported seed. *If you can't get seeds or planting material from local gardens – it is probably not a suitable local plant!*

### **Growing a garden of mixed plants**

In nature, one variety of one plant never grows alone. There are always lots of different plants of different kinds and sizes, all growing together. Anyone who has ever walked into a tropical jungle will know this very well. The reason people all over the world want to save the rainforest is because it has so many different kinds of plants all growing together. Growing plants in a food garden in a

way similar to how they grow in nature, as a mixed group of plants, is very good agriculture. Mixing plants in a garden usually gives more reliable food production, as any disease from one plant will wash off in the rain onto a different plant, where it cannot survive. Small plants fill the gaps and reduce the need for weeding.

### **Different types of plants for food security**

There is another reason for growing a range of food plants in a local garden or around a village. If something goes wrong, like extreme insect damage to plants, some disease occurring in the garden, or a poor growing season, some plants will be more damaged than others. With a variety of plants, there will still be some food to eat until the other plants recover and grow again. Also, a wide variety of plants will mean that different ones will be maturing at different times, which helps ensure a continuous supply of food. There are shrubs that can be planted as edible hedges around houses, and fruit and nut trees that need to be planted as a gift for your children, several years before they will be able to enjoy them. Some nuts can be stored and eaten when other foods are not available. Most yams will store well for a few months.

### **Looking after the soil**

Gardeners in traditional tropical agriculture usually move their gardens often by shifting to a new piece of land. There are usually three reasons for this:

- In the tropical lowlands, weeds can become a very big problem. There are usually a lot fewer weeds in the first year or two after clearing and burning the land, but weeds increase in the following years.
- Some of the nutrients in the soil are used each year and the soil becomes poorer and plants do not grow as well. There are ways of reducing this loss of nutrients.
- Very small worms called nematodes build up in the soil after a few years and get into the roots, especially of annual vegetable plants, and stop their roots working properly. For example, root knot nematode will cause the roots of plants like tomatoes and beans to become twisted resulting in poor growth of the plant.

### **Building up the soil**

When a new garden has been cleared, it has lots of leaf mulch and other old plant material. This provides plant nutrients for new plants to grow. There is a simple rule for growing plants and improving the soil—"If it has lived once, it can live again." Any old plant material can provide nutrients for new plants to grow, but it must be allowed to rot into mulch or compost for this to happen. If this plant material is burnt, some nutrients, especially phosphorus and potassium ("potash"), get left behind in the ashes for new plants to use, although it also allows these important nutrients to be lost by being washed away by rain. But with burning other important nutrients, such as nitrogen and sulphur, get lost in the smoke and disappear from the garden and soil. These last two plant nutrients are especially important for growing green leaves and when their levels are low, plants grow small or pale green. When nitrogen is lacking, the old leaves of the plant go pale and fall off early, and when sulphur is lacking, the young leaves go pale. Wherever possible, old plant material should be covered with some soil to allow it to rot down and not simply dry out or get burnt.

### **Poor soils where crops won't grow**

When soils are very acid (or sour), plants cannot get the necessary nutrients. Natural chemicals in the soil that are toxic to plants when present at higher levels become soluble, get into plants, and stop them growing. Adding limestone to these soils can improve them. Using compost will not make them less acid, but will keep the plant nutrients in the soil in a more readily available form that plants can use.

## **Soil nutrients**

Plants need 16 different kinds of plant food or nutrients in different amounts to grow properly. A plant that has already been growing will have these nutrients in them and probably even have them in a balanced amount. That is why composting old plant material is so important. Plants usually show some signs or symptoms if any of these nutrients is running out.

One of the most common and important nutrients for plant growth is nitrogen, which actually comes from the air, but gets into plants through the soil. When plants are short of nitrogen, their older leaves often become yellow or pale. When grass family plants, like sugarcane and corn, are short of nitrogen, the centre of the oldest (lowest) leaves starts to develop a dry or dead V-shape. The plant cannot find enough nitrogen in the soil so it gets it from an old leaf to grow a new leaf. This causes the old leaf to die, forming a characteristic V-shape in the centre of the leaf. The plant does not get any bigger as an old leaf dies each time a new leaf is produced. Village farmers often walk through grassland before they clear it for gardens, looking to see if the grass leaves are dry and dead, because they know gardens on this soil won't grow well. It is necessary to use compost or legumes (such as beans) to put nitrogen back into the soil. Growing plants from the bean family (legumes) is the most efficient way to increase the level of nitrogen in the soil.

Corn is a good plant for indicating which nutrients are running short in the soil. If the older leaves go dry along the edges, the soil is running out of potash. If leaves that are normally green develop a bluish colour, the soil is short of phosphorus. Generally, leafy crops need lots of nitrogen, and root crops need lots of potash.

## **Making compost**

Compost is old plant material that has been allowed to rot down into a fine, sweet smelling mulch that is full of nutrients that can be put back on the soil to grow new plants. Making good compost is very simple. A simple heap of plant material can be made in the corner of a garden or near a house. The composting process is carried out by small bacteria that live in the soil and feed on decaying plants. They break down old plant material into compost. These bacteria are living, so they need air, water and food. A good compost heap must have air, so don't cover it with plastic or put it in a container. This makes a foul smelling compost, as different bacteria that don't need air turn it into an acid mixture that preserves it. Good compost must have moisture, so keep the heap damp, but not too wet. The compost bacteria like a balanced diet, which means that both green material and dried material is needed to balance the carbon and nitrogen in the compost pile. If the compost material gets too dry and brown, it will not break down, and if it gets too green, it will go slimy. Using a little bit of compost from an old heap will make sure the right bacteria are there to start the whole process off. As soon as the plant material is broken down to a fine mulch it can be put onto the garden. It is best if it is dug in, but if it is regularly put onto the surface of the garden, worms will mix it into the soil.

## **Pests**


There are a large number of insects that enjoy sharing our food with us! We should not try to kill all these insects as they have an important role to play in keeping everything in nature in balance. What we need to do is to learn to manage these insects so we can all get some food to eat! Some insects are attracted to lights, and if the garden is near village lights some insects can cause a lot of damage. If large areas of one particular crop are planted, insects can breed more quickly and cause a lot of damage. As an example, insects called armyworms can breed up in large numbers on the shade trees of cacao and then move "like an army" into gardens. Some insects are large and breed slowly and can be picked off and removed. The large, green grubs with pointy tips that hide under taro leaves are best controlled by simply picking them off. Some insects, like taro beetles, can be a

serious problem, but the young curl grubs of this insect are tasty if you catch and cook them. Some insects do not like sunlight. The very small moth that damages banana fruit is like this. Simply pulling off the leafy bracts over the banana fruit reduces the damage, as this lets sunlight in and the insect flies away. The best rule for reducing pest damage is to grow healthy plants, as they suffer less damage.


### **Diseases**

The living organisms that cause disease are much smaller than insects. These disease organisms can often only be seen with a microscope. There are three main kinds of disease organisms-fungi, bacteria and viruses. Fungi are like the mushrooms we eat, only very much smaller. They usually make distinct dry spots on leaves and other plant parts. Fungi have spores that often blow in the wind. Bacteria are often smaller and live in damp places. They usually make plants go soft and squashy, and they may cause a smell. Bacteria are mostly spread with rain and in water. Viruses are very, very small and usually make irregular stripes and patterns on leaves and other plant parts. Viruses usually spread in planting material or in the mouths of small sucking insects. One common fungus disease on sweet potato causes the leaves to become wrinkled and twisted. It usually gets worse in old gardens and where soils are running out of nutrients. It doesn't affect all kinds of sweet potato to the same extent. The answer is not to stop the disease, but to improve the soil. The general rule is that healthy plants that are growing well will suffer less damage from disease.


## Food value charts for a selection of plants from Liberia




Protein helps the body repair cells and make new ones. Protein is also important for growth and development in children, teens, and pregnant women. Symptoms of protein deficiency include wasting and shrinkage of muscle tissue, and slow growth (in children).




Vitamin A is very important for eyesight and fighting disease, particularly in infants, young children and pregnant women. People who are short of Vitamin A have trouble seeing at night.



Vitamin C helps us avoid sickness, heal wounds, prevent infections and absorb iron from food. Severe vitamin C deficiency increases the risk of scurvy with symptoms such as inflammation of the gums, scaly skin, nosebleed and painful joints.



Iron is important because it helps red blood cells carry oxygen from the lungs to the rest of the body. Low levels of iron cause anaemia, which makes us feel fatigued. Iron is also important to maintain healthy cells, skin, hair and nails. Iron is more available when Vitamin C is also present.



Zinc is particularly important for the health of young children and teenagers, and to help recovery from illness. It is needed for the body's immune system to work properly. It plays a role in cell division, cell growth, wound healing, and the breakdown of carbohydrates. Zinc is also needed for the senses of smell and taste. Zinc deficiency is characterized by stunted growth, loss of appetite, and impaired immune function.

**Note regarding plant selection:** In compiling these field guides, we acknowledge that some staple foods and commercial crops which are grown widely in the target country may be omitted. Such foods are often in the starchy staple category (e.g. rice, corn). This does not mean that they are not useful, but merely reflects a desire for the Food Plant Solutions project to concentrate on plants that are less well known and/or underutilised.





## Starchy staples

**English:** Floating rice

**Local:**

**Scientific name:** *Oryza glaberrima*

**Plant family:** POACEAE

**Description:** An annual grass. Most varieties have a reddish colour. It grows to 1.5 m tall but can be up to 5 m in some floating kinds. Dryland types often form tufts and floating rice often branches. The leaves are alternate and simple. The leaf sheath is 25 cm long. The leaf blade is 20-25 cm long and 6-9 mm wide. The flowers are in a compact group 25 cm long at the top of the plant. The fruit is a grain 9 mm long by 3 mm wide.

**Distribution:** It is a tropical plant that grows in swamps and on flood plains of savanna regions. It does best with temperatures of 30-35°C. It grows from sea level to 1700 m altitude and can tolerate low soil fertility.

**Use:** The grain can be cooked and eaten, or ground into flour. This is sweetened with rice flour and honey for bread. It is fermented for beer.



**Cultivation:** Plants are grown from seed. Usually seeds remain dormant for a few months after harvesting. Seed usually emerge after 4-5 days. Seed are usually broadcast without using a nursery. The juvenile stage lasts for 3 weeks then tillering occurs for 3-4 weeks.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed        | 11.3       | 1538      | 7.4       | -           | -           | 3.4     | -       |

Image sourced from: <https://i.pinimg.com/474x/77/48/1f/77481fb8134243595c851b56ac63ea4c--natural-resources-mali.jpg>

## Starchy staples

**English:** Sorghum

**Local:**

**Scientific name:** *Sorghum bicolor*

**Plant family:** POACEAE

**Description:** Sorghum is a millet grass. A mature sorghum plant resembles maize in its stature. Plants vary in height from 45 cm to 4 m. It is an annual grass with erect solid stems. The stems can be 3 cm across at the base. Prop roots occur at the base of the plant. There are numerous sorghum varieties. Some have one main stem while others produce multiple tillers. More tillers are produced when plants are widely spaced. The nodes on the stem are slightly thickened. Short types have up to 7 leaves while tall late varieties may have up to 24 leaves. The leaf blade can be 30-135 cm long. Leaves are bluish green and waxy. They have a prominent midrib. The large flower panicle can be 20-40 cm long. The flower occurs at the top of the plant. It can stick upright or bend over. The flower can be open or compact. Over 1000 cultivated varieties occur in China.



**Distribution:** Sorghum is a tropical plant. It suits the savannah zones in the tropics and can tolerate heat and drought. It can recover from drought even as a seedling. It can tolerate water-logging. It can be grown on heavy or light soils. Sorghum requires short day lengths to flower. Many kinds are adapted to specific day length and rainfall patterns. It suits hardiness zones 9-12.

**Use:** Sorghum seeds are eaten as a cereal. Flour can be made from the grain and then used for porridge or other dishes. It is used for dumplings, fried cakes and drinks. It cannot be used for bread as it contains no gluten. The stems of some kinds are sweet and can be chewed. The grains can be popped and eaten. The sprouted seeds can also be eaten.

**Cultivation:** Sorghum seeds will germinate soon after harvest. The seeds also store well if kept dry and protected from insects.

**Production:** Grain is ready for harvest 4-8 weeks after flowering.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed        | -          | 1459      | 11.1      | -           | -           | -       | -       |

## Starchy staples

**English:** Taro

**Local:**

**Scientific name:** *Colocasia esculenta*

**Plant family:** ARACEAE

**Description:** This plant has large flat leaves on the end of upright leaf stalks. It grows up to 1 m high. The leaf stalk or petiole joins the leaf towards the centre of the leaf. The leaves are 20-50 cm long. Near the ground a thickened rounded corm is produced. Around this plant there is normally a ring of small plants called suckers. Many different varieties occur. If left to maturity, a lily type flower is produced in the centre of the plant. It has a spathe 15-30 cm long which is rolled inwards. The flowers are yellow and fused along the stalk. There are many named cultivated varieties. Taro comes in two basic forms. The Dasheen type *Colocasia esculenta* var. *esculenta* and *Colocasia esculenta* var. *antiquorum* or the Eddoe type. The basic difference is the adaptation of the Eddoe type to storage and survival in seasonally dry places, while the dasheen type needs to be maintained in a more or less continuously growing vegetative stage.



**Distribution:** It is a tropical plant. Taro grows from sea level up to about 2300 m altitude in the tropics. It grows well in humid places. It can stand damp soil and grow under light shade. It suits hardiness zones 9-12.

**Use:** The corms, petioles and leaves are all edible after cooking. The leaves are also dried and stored. Fresh leaves can be stored for 4-5 days. **Caution:** Some varieties burn the throat due to oxalate crystals.

**Cultivation:** Taro can be planted from cormels or from the top of the central corm. Other sections of the corm could also be used but this is not commonly done. Flowering of taro and seed production can lead to new cultivars. Flowering can be promoted by the use of gibberellic acid. The general growth pattern is for an increase in top growth, in terms of leaf number, leaf area and petiole length, to continue for about 6 months under tropical lowland conditions then for each of these to decrease and tuber storage to continue to increase. Corm weight increases significantly from 5-11 months. Starch content also increases with time but protein content declines over the corm development period.

Taro can be grown under flooded conditions but root rots develop if the water becomes stagnant. For flooded cultivation, the land is cleared, ploughed, cultivated and puddled. The aim is to get a field that is flat with embankments allowing the impounding of water. Planting is done into 2-5 cm of standing water.

For dryland taro, the soil is prepared by digging, unless a fresh bush fallow is used where the natural friability of the soil allows plants to be put into the undug soil in a small hole that is prepared. Plants are put into a hole 5-7 cm deep or deeper. Mulching to conserve moisture and reduce weed growth is beneficial. Setts from corms normally give higher yield than that from cormels. The greater leaf area and root production may be responsible for this. Setts of about 150 g are optimum.

The time of planting is primarily determined by the availability of moisture. Planting is done shortly after the rainfall has become regular, if seasonally distinct wet and dry occur. Higher rainfall, higher temperatures, and higher hours of sunlight, enhance production and determine seasonality of production.

Evapotranspiration for flooded taro averages about 4 mm per day, ranging from 1.5-7.2 mm, with a total of about 1200 mm for the crop. Intermittent moisture can result in irregular shaped corms. Flooding has been found to be more effective than sprinkler irrigation, or furrow irrigation. Increased suckering, giving greater leaf area, seems to be the reason for this.

Taro is sensitive to weed competition throughout most of its growth, but it is more critical during early growth up to 3-4 months. About 7-9 weeding are required, to keep the crop clean under tropical lowland conditions, where flooding is not used. Due to the decrease in height and leaf area towards the end of the growth cycle when starch accumulation in the corms is maximum, weed competition and weed control are again significant. Mechanical weeding needs to be shallow to avoid damaging the superficial taro roots. A range of herbicides have been recommended in various situations.

Taro produces the highest dry matter yield under full sunlight, but it can still grow under moderate shade. Under shaded conditions it grows more slowly and develops fewer cormels. They require good moisture conditions and have little tolerance for drought. Taro residue has an allelopathic factor which can reduce the germination and growth of other plants, for example, beans.

Taro tends to demand high fertility, and is responsive to additional NPK fertiliser. Higher doses of K increases starch content and higher doses of N increases protein content. Both N and K applications increase oxalic acid content of the tubers.

Spacing affects total yield, and marketable, harvestable yield, of corms. Close spacing increases the corm yield per area, and the shoot yield per area, but decreases the corm yield per plant, and the contribution of sucker corms, to the yield. Where spacings of 30 cm x 30 cm are used, giving about 110,000 plants per hectare, a very large amount of planting material is required, which reduces the net return per unit of planting material. A spacing of 60 cm x 60 cm is more common. Wider spacings of 90 cm x 90 cm reduces overall yield.

**Production:** Crops mature in 6-18 months. Yields of 5-15 tonnes per hectare are probably average.

**Food Value:** Per 100 g edible portion

| Edible part   | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|---------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| root          | 66.8       | 1231      | 1.96      | 3           | 5           | 0.68    | 3.2     |
| leaf          | 85.0       | 210       | 5.0       | 57          | 90          | 0.62    | 0.7     |
| leaf stalk    | 93.0       | 101       | 0.5       | 180         | 13          | 0.9     | -       |
| leaf (cooked) | 92.2       | 100       | 2.7       | 424         | 35.5        | 1.2     | 0.2     |

## Starchy staples

**English:** Chinese water chestnut

**Local:**

**Scientific name:** *Eleocharis dulcis*

**Plant family:** CYPERACEAE

**Description:** A herb which grows in water. It is a tufted sedge with round green stems. The bases are covered with brown sheaths. The stem is about a metre high and 1 cm across. It grows 30-200 cm high. From the top of each planted corm, several slender horizontal rhizomes radiate out into the mud, each terminating in a corm. The edible part consists of a flattened corm. The rhizome is short. Under the ground there are stolons bearing tubers. The tubers are almost round and have 4-6 distinct rings. They are usually about 1 cm across but can be up to 4 cm across. They are dark brown. The stems are tufted and slender. There are fine lines along the stems. The purplish leaves are reduced to thin tube like sheaths. Each plant produces these long tubular leaves that project above the water surface. The flower spike is on the end of the plant. There are many flowers, 1.5-6 cm long by 3-6 mm wide. The fruit is a nut 1.5-2 mm long.



**Distribution:** It suits humid, monsoonal, tropical and subtropical locations. It is found in open wet places and shallow water. It grows in fresh water swampy grounds or in shallow water. It is also found in rice fields. It needs at least 220 frost free days. It needs a soil temperature above 15.5°C for germination of the corms. It needs a pH of 6.9-7.3. It can be grown up to 1200 m altitude and suits plant hardiness zones 9-12.

**Use:** The tubers are cooked and eaten. The corms can be eaten raw, roasted or boiled after they have been peeled. Normally, they are cut into small slices and added to soup or to fish and meat dishes. They can be sweetened for desserts.

**Cultivation:** Plants can be grown by division or tubers. They are put in holes 20-30 cm deep. Fields are flooded after planting then allowed to drain. When top growth is 20-30 cm high fields are flooded to at least 10-12.5 cm. A spacing of 75 cm x 75 cm is suitable. 500 kg of corms per hectare are required for planting.

**Production:** Corms mature after 7-8 months. Yields of 20-40 t per ha are possible.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| corm        | 50.8       | 635       | 3.7       | -           | 52          | 15      | 1.9     |
| fruit       | 79.6       | 268       | 1.4       | 0           | 5           | 0.7     | 0.5     |



## Starchy staples

**English:** Cassava

**Local:**

**Scientific name:** *Manihot esculenta*

**Plant family:** EUPHORBIACEAE

**Description:** A plant which can re-grow year after year from the thickened roots. It has several stems. The stems are woody and have some branches. Plants grow up to 3 metres tall. Stalks have distinct scars where leaves have fallen. The leaves tend to be near the ends of branches. The leaves are divided like the fingers on a hand. The leaves have long leaf stalks. The leaves have 3-7 long lobes which can be 20 cm long. These are widest about 1/3 of the distance from the tip and taper towards the base. The colour varies. It produces several long tubers. These can be 50 cm long by 10 cm across. The flowers are on short stalks around a central stalk. They are produced near the ends of branches. The female flowers are near the base of the flower stalk and the male flowers higher up.



**Distribution:** A tropical plant. Plants grow from sea level up to about 1650 m. They can grow in poor soil and can survive drought. It grows between 25°N and 25°S and needs a rainfall above 750 mm. It suits hardiness zones 10-12.

**Use:** The tubers are eaten after thorough cooking. They are boiled, roasted or made into flour. The starch is used in puddings, soups and dumplings. Young leaves are edible after cooking. They are also sometimes dried and stored. Seeds are also eaten. **Caution:** Bitter kinds of cassava contain poison but this is destroyed on heating. This kind of cassava should be cooked, sun dried, soaked and cooked again.

**Cultivation:** Cassava is planted from sections of the stalk. Sections about 15-20 cm long of the more mature woody stem are cut and stuck into the ground. They can be completely buried or put at almost any angle and it affects the growth little. Soon roots form and leaves start to sprout from the stalk. Cassava seeds need a soil temperature of 30°C for their germination. Flower and fruit production is more common under lower temperatures such as in highland or less equatorial conditions.

It is not necessary to dig a hole to plant cassava and on many soils where the soil is loose it can be planted without digging the soil first. Cassava does not suit waterlogged soils and preferably they should not be too shallow or stony.

Cassava can be planted at any time of the year but to get started it needs moisture so is often planted near the beginning of the wet season. The crop once established can survive for several months without rain. The ability to tolerate drought varies significantly with cultivar. During drought less and smaller leaves are produced and leaves die off more quickly but storage roots can be increased in the short term.

Because cassava can still grow satisfactorily in poorer soils it is often put last in a rotation after others crops have already been grown on the piece of land. Cassava is more responsive to nitrogen and potassium than phosphorus under many field situations. Nitrogen can increase cyanide levels. Under very acid conditions with high soluble aluminium levels, cassava has been able to achieve and

maintain top growth but with significantly reduce root yields. When drainage is good and soil moisture is adequate, cassava stalks can be planted at any orientation from horizontal to vertical, but in very sandy soils horizontal planting is best and in heavy clay soils vertical planting is best.

Because of the slow growth in early establishment stages, soil loss from erosion with heavy rains can be significant. To avoid this planting should be timed so that the maximum vegetative growth is occurring during the heaviest rains. A leaf area index between 2.5-3.5 is optimal for cassava yield. The critical period for weed control is the time from 2-8 weeks after planting. Cassava tuber bulking is delayed under shaded conditions. Yields are also reduced. In mixed cropping situations using crops which mature early, allowing the cassava time to recover, is one possible strategy. For optimum production shading should be avoided.

Cassava takes about 10-12 months to produce mature tubers in the lowlands tropics although some varieties produce a smaller yield earlier. Yields in the range of 20-45 t/ha have been recorded for 12-14 month crops. The plants can be left growing and the tubers stored in the soil for considerable time. Crops of 24 months duration occur. Once the tubers have been dug they do not keep for more than a few days. Pre-harvest pruning of plants increases the storage time of tubers after harvest.

Spacing and plant density varies with soil climatic conditions and variety. Plant densities from 10000-30000 plants per hectare are used. Plants from the higher density crops have been shown to have quick post-harvest deterioration. Mulching has given significant yield increases in some conditions. It also reduces the incidence and damage of some root boring insects.

**Production:** Plants can be harvested after 10 months in the lowlands. There are some faster growing varieties. Yields in the range of 20-45 t/ha have been recorded for 12-14 month crops.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| tuber       | 62.8       | 625       | 1.4       | 30          | 15          | 0.23    | 0.48    |
| leaf        | 82.0       | 382       | 7.1       | 57          | 275         | 7.6     | -       |

## Starchy staples

**English:** Potato yam

**Local:**

**Scientific name:** *Dioscorea bulbifera*

**Plant family:** DIOSCOREACEAE

**Description:** A yam with a long smooth stemmed vine, round in cross section and without spines. The vine winds to the left, can climb into trees and grow to long lengths. The large leaves (14-30 cm across and slightly longer than wide) have pointed tips and round bases. About 7 veins arise from the tip of the leaf stalk. It produces often flattened bulbils (potatoes) in the leaf angles along the vine. They can be grey brown or purple. The smaller tuber underground is normally covered with roots. The flowers are large. The male flowers are in spikes up to 20 cm long. The female spikes are usually in pairs. The winged fruit are about 2.5 cm long by 1.5 cm across. The seeds have wings. The bulbils normally have few fibres through the tissue compared to some yam tubers. Many varieties have yellow flesh.



**Distribution:** An annual tropical plant. It will grow from the coast up to about 1700 m altitude in equatorial zones. It is common near the edge of grassland and forest at mid altitudes. Both wild and cultivated forms occur. It is common near secondary forest at low and medium altitudes.

**Use:** The cooked tubers aerial bulbils are eaten. Some kinds are bitter and inedible or at least require special processing and cooking. Some varieties are poisonous.

**Cultivation:** Either the vine bulbils or the underground tubers are planted. The long vines can be trained up trees. The bulbils need a set storage time before sprouting. The leaves die off for 1-4 months each year before re-sprouting from the tuber. Bulbils only grow shoots from one end unless the bulbil is cut into pieces. If larger bulbils are cut, the cut surfaces should be dried and healed in a shady place for 2-3 days before planting. Bulbils are planted 8-12 cm below ground at a spacing of about 100 cm by 100 cm. Nitrogen and potassium fertilisers give greater responses than phosphorus. Friable well drained soils are most suitable. Often little cultivation or mounding is done. A high level of organic matter improves yield. Strong staking is required with branched 2 m stakes, or with trees or living stakes. Vine and tuber growth can be extensive and heavy.

**Production:** Bulbils or aerial yams are produced as leaves begin to unfold, continue until plant maturity, and often fall. Harvesting can start 3 months after planting but immature tubers have less starch. Underground tubers are normally not harvested until leaf die back. Wounds and damage to the tubers normally heal naturally given dry aerated conditions. Some varieties have seasonally dormant aerial tubers which only grow after an extended storage period. Others germinate quickly.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| tuber       | 70.8       | 357       | 2.7       | -           | 78          | 3.1     | 0.4     |
| bulbil      | 79.4       | 326       | 1.4       | -           | -           | 2.0     | -       |



## Starchy staples

**English:** Sweet potato

**Local:**

**Scientific name:** *Ipomoea batatas*

**Plant family:** CONVOLVULACEAE

**Description:** This is a root crop which produces long creeping vines. The leaves are carried singly along the vine. Leaves can vary considerably from divided like fingers on a hand, to being entire and rounded or heart shaped. Purple trumpet shaped flowers grow at the end of the vine. Fattened tubers are produced under the ground. There are a large number of varieties which vary in leaf shape and colour, tuber shape, colour, texture and in several other ways.



**Distribution:** A tropical and subtropical plant. They grow from sea level up to about 2700 m altitude in the tropics. Plants can grow with a wide range of rainfall patterns and in different soils. Plants are killed by frost and can't stand water-logging. Plants grow well with temperatures between 21-26°C. It can grow with a pH between 5.2-6.8. Sweet potato are not tolerant to shading. It suits hardiness zones 9-12.

**Use:** Tubers are boiled or baked. They can be steamed, fried, mashed or dried. They can be fermented into alcoholic drinks. They can also be used in pies, cakes, puddings and candies and jams. They can be used in noodles. The chopped and dried tubers can be boiled with rice or ground into flour and mixed with wheat flour to make cakes or bread. The young leaves are edible.

**Cultivation:** Vine cuttings are used for planting. In grassland soils it is grown in mounds, ridges or other raised beds. In bush fallow, it is mostly planted in undug loose soils. It needs a sunny position. Tubers won't form if the ground is waterlogged when tubers start to develop. Sweet potato is grown by cuttings of the vine. About 33000 cuttings are required per hectare. These weigh about 500 kg. Vine lengths of about 30 cm are optimum. As long as the vine is adequately inserted in the soil, the length of vine inserted does not significantly affect yield. Fresh sweet potato seeds germinate relatively easily and lead to continuous production of new cultivars under tropical conditions. Excess nitrogen restricts storage root initiation and therefore excess leaves are produced without significant tuber yield. Dry matter percentage increases with increasing age of the crop. Higher dry matter tubers are normally preferred.

Sweet potato are not tolerant to shading. Under shaded conditions, both foliage growth and storage root production are decreased. Some cultivars can be selected for increased production under mild shade but not heavy shade. The survival of cuttings at planting is also reduced under shaded conditions. Under shaded conditions, plant become more climbing and with fewer, larger leaves. With increasing shade, fewer tubers are produced and these grow more slowly. Sweet potato tends to be responsive to potassium fertiliser. Cultivars are often selected for yield under low fertility conditions.

Under lowland conditions in the tropics sweet potato tubers undergo active tuber enlargement from 6-16 weeks. Weed control is essential especially during early stages of growth. The rate of ground coverage by foliage varies greatly with growing conditions and cultivar, but once ground coverage has occurred, weed control is less of a problem. Sweet potato tuber initiation is subject to

aeration in the soil. Either heavy clay soils, waterlogged conditions or other factors reducing aeration can result in poor tuber production. For this reason, sweet potatoes are often grown on mounded beds. In well drained or high organic matter soils, digging or mounding is not as essential. Leaf scab (*Elsinoe batatas*) can significantly reduce yield especially in sites where leaf production is low due to low soil fertility. To reduce sweet potato weevil damage, plants need to be hilled or have the tubers well covered with soil. Cracking soils can allow the weevil access to tubers.

**Production:** The time to maturity ranges from 5 months to 12 months depending on the variety planted and the altitude at which it is being grown. Yields range from 6-23 t/ha.

**Food Value:** Per 100 g edible portion

| Edible part    | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|----------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| tuber (baked)  | 72.9       | 431       | 1.7       | 961         | 24.6        | 0.5     | 0.3     |
| tuber (raw)    | 70.0       | 387       | 1.2       | 709         | 25          | 0.7     | 0.4     |
| tuber (boiled) | 72.0       | 363       | 1.1       | 787         | 15          | 0.6     | 0.3     |
| leaf           | 86.3       | 168       | 3.9       | 105         | 58          | 2.9     | -       |

## Legumes

**English:** Sweet acacia

**Local:**

**Scientific name:** *Acacia farnesiana*

**Plant family:** FABACEAE

**Description:** An evergreen shrub. It grows 5-7 m tall and 3 m across. The stem is slender and erect. The crown is open. It is a spreading, densely branched shrub. The bark is smooth and brown. The leaves are branched and green. There are 4-6 pairs of larger leaves and 10-20 pairs of small leaflets. They have tiny leaflets and thorns up to 2 cm long, occur in pairs. The leaf stalk has a gland at or above the middle. The flowers are large orange balls. They are strongly perfumed. (The oil is used as a perfume in France.) The pods are long and dark brown to black. They



are 5-8 cm long by 0.5-1 cm wide. They are inflated and sausage like. Often they are curved. They are marked with narrow lines. The pods have hard grey seeds imbedded in a pithy substance. The pods do not split open at maturity. The seeds are chestnut brown and 7-8 mm long by 5.5 mm wide.

**Distribution:** It is a tropical plant. This tree occurs naturally in Australia, Asia and Africa. It will grow on most soils. It is drought and frost resistant. It most commonly grows naturally on clay soils. It grows in areas with an annual rainfall between 400-4000 mm. It can grow in acid or alkaline soils. It can grow in arid places. It suits hardiness zones 11-12.

**Use:** The pods have been recorded as eaten after cooking. The gum is eaten. The ground up seeds are eaten. The germinated seeds are claimed to be eaten. The gum is used to prepare sweets. The young leaves are used in India as a substitute for tamarind in chutneys.

**Cultivation:** It is grown from seed.

**Production:** It is fast growing. Flowering can occur almost continuously if watering is regular. In northern Australia, flowering is normally May to July, with pods available from September to November.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)  | 8.1        | 1522      | 36.6      | -           | -           | 6.0     | 0.6     |

## Legumes

**English:** Soybean

**Local:**

**Scientific name:** *Glycine max*

**Plant family:** FABACEAE

**Description:** A small erect bean growing up to 60 cm tall. It grows each year from seed. Straggling kinds can occur. Stems, leaves and pods are softly hairy. The leaves have 3 leaflets. The leaflets have stalks. Flowers are small and white or blue. They occur in groups in the axils of leaves. The pods are broad, flat and hairy. Pods have 2-4 seeds. The seeds can be yellow to black.



**Distribution:** It is a temperate plant that suits lowland areas. It can be grown from sea level to 2000 m altitude. Many varieties will not flower in the tropics (short days). It needs fertile soil. The best soil acidity is pH 5.5-7.0. It is damaged by frost.

**Use:** The young pods and ripe seeds are eaten. They are used for flour. The dried seeds are boiled or baked and used in soups, stews and casseroles. The seeds are used for oil. Toasted seeds are eaten like a snack. Strongly roasted seeds are used for coffee. Soy flour is used for noodles, and confectionary. The beans are fermented and used in a range of foods. Sometimes the young leaves are eaten. The seeds are also used for sprouts and for making cooking oil and soya sauce etc. Because soybean contains a trypsin inhibitor they should be cooked and even the sprouts should be lightly cooked.

**Cultivation:** It is grown from seed. Seeds need to be inoculated with bacteria before planting. Plants need to be about 20 cm apart.

**Production:** Plants flower about 8 weeks after sowing and pods mature about 16 weeks after sowing. Often plants are pulled up and hung up before threshing out the seed.

**Food Value:** Per 100 g edible portion

| Edible part     | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-----------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed            | 9.0        | 1701      | 33.7      | 55          | -           | 6.1     | -       |
| seed (immature) | 68.0       | 584       | 13.0      | 16          | 27          | 3.8     | 0.9     |
| sprout          | 79.5       | 339       | 8.5       | 1.0         | 8.3         | 1.3     | 1.0     |

## Legumes

**English:** Egyptian sesban

**Local:**

**Scientific name:** *Sesbania sesban*

**Plant family:** FABACEAE

**Description:** A shrub that grows to 6 m tall. The bark is reddish-brown. The leaves are made up of 10-25 pairs of opposite oblong leaflets. They can be 15 mm long by 3 mm wide. The flowers are yellow and pea shaped. The standard petal is often speckled with finely veined dark maroon. They occur in many flowered sprays that are up to 15 cm long. The fruit are long slender pods, 30 cm long by 0.3 cm wide. They are often slightly curved.



**Distribution:** A tropical plant that grows in low lying areas, usually near water. It can survive water-logging and salty soils. It grows in areas 350-1500 m above sea level. It grows in savannah woodland and can grow in arid places.

**Use:** The leaves and young flowers are eaten. They are often fried or pounded with rice and beans. The seeds are used for food in times of scarcity. (They have a protein inhibitor preventing the protein being well used.) The seeds are also fermented into a flavouring paste.

**Cultivation:** It can be grown as a hedge. It can be cut back and will re-grow.

**Production:** It is fast growing and only lives for short time.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed        | 9.2        | 1446      | 32.0      | -           | -           | -       | -       |



## Legumes

**English:** Cowpea

**Local:**

**Scientific name:** *Vigna unguiculata*

**Plant family:** FABACEAE

**Description:** A creeping bean type plant with straight firm pods. There is a deep tap root and many branches occur from it in the surface of the soil. The root nodules are large and round. The leaves have 3 leaflets. The end leaflet can be 12-16 cm long. The side leaflets are asymmetrical. The stipules at the base of the leaf are large and with spurs at their base. Flowers occur often in pairs on the end of long flowering shoots. Only 2-4 flowers in each stalk produce pods. Flowers are white, yellow or blue. They are large and showy. The pods are about 15 cm long. The seeds are white except for a dark scar.



**Distribution:** It grows in tropical and subtropical climates. It grows from sea level to 1800 metres altitude in the tropics. Plants can stand high temperatures. Some kinds can tolerate drought. They are sensitive to cold and killed by frost. Plants germinate with a temperature between 11.5-15.5°C. The best growth occurs between 20-35°C. They can grow on a range of soils providing they are well drained. They are a short day plant. They do well in the semiarid tropics. It will not tolerate acid or alkaline soils. It grows in areas with an annual rainfall between 280-410 mm. It can grow in arid places.

**Use:** The young leaves, young pods and ripe seeds are all eaten. They can be steamed, boiled, stir-fried etc. The leaves can be dried and stored. The dried seeds are used in soups and stews. They are ground into flour or fermented. The seeds are also used for bean sprouts. Roasted seeds are used as a coffee substitute.

**Cultivation:** It is grown from seeds. Seeds remain viable for several years if carefully stored. A seeding rate of about 20 kg per ha is suitable and seed are sometimes broadcast then thinned.

**Food Value:** Per 100 g edible portion

| Edible part               | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|---------------------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)                | 11.2       | 1189      | 23.5      | -           | 1.5         | 6.4     | -       |
| seed (young, boiled)      | 75.5       | 406       | 3.2       | 79          | 2.2         | 1.1     | 1.0     |
| leaf                      | 88.4       | 143       | 4.2       | 36          | 35          | 4.7     | 0.3     |
| young pod + seed (boiled) | 89.5       | 142       | 2.6       | 45          | 17.0        | 0.7     | 0.2     |
| leaf (boiled)             | 91.3       | 92        | 4.7       | 29          | 18          | 1.1     | 0.2     |

## Legumes

**English:** Lablab bean

**Local:**

**Scientific name:** *Lablab purpureus*

**Plant family:** FABACEAE

**Description:** A climbing bean which can have vines 1-5 m long. It keeps growing from year to year. The stems can be smooth or hairy. Leaves are made up of 3 almost triangular leaflets. The leaflets are 5-15 cm long and 3-14 cm wide. The side leaflets are somewhat asymmetrical. Often the plants are flushed purple. The flowering clusters are 5-20 cm long. Flowers are often white but can vary from red to blue. The pods are flattened, pointed and up to 12 cm long and 2 cm wide. They can be green, purple or white. Inside there are 3-5 white or dark seeds. Seed pods have a wavy margin. The seeds are 0.5-1.5 cm long. (This bean is similar to Lima bean but the keel of the flower is not spirally twisted, the pod ends more bluntly with a long thin style at the end and the hilum on the seed is longer.)



**Distribution:** It is a tropical and subtropical plant. It mostly grows between 750 and 2175 m altitude in the tropics. It is drought resistant and can grow in quite low rainfall areas. Some varieties are short day and some are long day kinds. It suits hardiness zones 9-12.

**Use:** The young pods, ripe seeds and young leaves are edible, cooked. Flowers can be eaten raw, steamed or added to soups and stews. Dried seeds can be cooked as a vegetable. The seeds can also be sprouted then crushed and cooked. The large starchy root is edible. **Caution:** Many types can be poisonous. They should be boiled and the cooking water thrown away.

**Cultivation:** Seeds are sown at 30 x 60 cm spacing near stakes or trees. About 20 kg of seed per hectare are required. Fertilising with nitrogen and potash until flowering is recommended.

**Production:** Young pods are ready 4-6 months after planting and seeds 6-8 months. Pods are often harvested over 2 or 3 years. Pollination and seed setting are reduced in cold weather.

**Food Value:** Per 100 g edible portion

| Edible part  | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|--------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)   | 10.0       | 1428      | 22.8      | -           | -           | 9.0     | -       |
| seed (young) | 86.9       | 209       | 3.0       | 14          | 5.1         | 0.8     | 0.4     |
| pod (fresh)  | 86.7       | 203       | 3.9       | -           | 1.0         | 2.4     | -       |

## Legumes

**English:** Jack bean

**Local:**

**Scientific name:** *Canavalia ensiformis*

**Plant family:** FABACEAE

**Description:** A perennial climber, although short kinds do occur. Often it is a more bushy plant than the sword bean. Plants grow up to 1.5 m long. Stems can be hairy. Leaves have 3 leaflets. The leaflets are oval and 5.7-20 cm long by 3.2-11.5 cm wide. The leaf tends to be wedge shaped at the base. The leaf stalks are 2.5-11 cm long. Flowers are red/purple. They occur on flower clusters 5-12 cm long and with flower cluster stalks which are 10-34 cm long. The individual flower stalks are 2-5 mm long. Pods are long and sword shaped. Pods can be 15-35 cm long. Seeds are white with a light brown hilum half as long as the seed. Seeds are 2 cm long, by 1 cm across.

**Distribution:** It grows in tropical and subtropical places. It requires a fairly high temperature (15°-30°C). It will possibly grow up to 900 m altitude. It is fairly drought resistant and also has some resistance to water-logging and salt in the soil. It can tolerate shade. It can tolerate pH from 4.5-8.0 but does best at about 6.1. The optimum mean annual temperature is 14.4°-27.8°C. Seed germinate between 24-27.5°C. It is a short day plant growing well with a day length of 10-12 hours of sunlight. It can grow in arid places.



**Use:** The leaves and top shoots are eaten. The very young pods are boiled and eaten. The flowers can be eaten. The young seeds are eaten boiled, roasted, or peeled and cooked. The seeds are also fermented. The ripe seeds are roasted and used as a coffee substitute. **Caution:** The ripe seeds can contain poison and need to be well cooked and the water changed before eating. They are also often left under running water or fermented.

**Cultivation:** It is grown from seeds. Seeds need to be 2 cm deep. A spacing of about 60 cm is suitable. Plants preferably need a support to climb over. It benefits from a fertile soil but adding nitrogen depresses yield.

**Production:** Green pods are produced in 3-4 months, but ripe seeds need 6-9 months. Yield of seeds can range from 700-5400 kg/ha.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed        | 10.0       | 1423      | 20.4      | 160         | -           | 4.9     | -       |
| pod (fresh) | 88.0       | 155       | 2.4       | -           | -           | -       | -       |



## Legumes

**English:** Rice bean

**Local:**

**Scientific name:** *Vigna umbellata*

**Plant family:** FABACEAE

**Description:** An annual twining, climbing bean plant with a slender hairy vine. It grows from seed each year. It grows to 1.5-3 m long. Leaves have 3 leaflets which can vary in shape. They are mostly oval and 3-13 cm long by 1.5-7 cm wide. They taper towards the tip and are rounded at the base. Usually they are hairy. The leaf stalks are 3-16 cm long. Flowers are about 1.5 cm long in dense cone shaped clusters. These flowering stalks can be 3-10 cm long. The flowers are yellow. The fruit are straight pods about 10 cm long and 5 mm wide. Seeds are small (5-8 mm long) and yellow to brown. The pods split open easily. The seeds can be yellow, green, brown, red, black or mottled.



**Distribution:** A tropical plant that grows to 1800 m altitude in the tropics. It suits wet climates. It occasionally becomes self sown in coastal grasslands. It needs a sunny protected position and is drought and frost tender. It can grow in arid places.

**Use:** The young pods and ripe seeds are eaten cooked. The dried seeds are boiled and served with rice or used in soups and stews. The young leaves can be eaten. The seeds are used in bean sprouts. Seeds should be cooked or crushed if fed to pigs.

**Cultivation:** It is grown from seeds. Seed collection is easy. Seeds often have a hard skin which must be broken (e.g. by scraping) to help germination.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed        | 13.0       | 1373      | 20.9      | -           | -           | 10.9    | -       |

## Legumes

**English:** Pigeon pea

**Local:**

**Scientific name:** *Cajanus cajan*

**Plant family:** FABACEAE

**Description:** An upright perennial shrubby legume that can live for 3-4 years. They can grow up to 4 m tall and spread to 1.5 m wide. It has a bushy appearance and a strong deep taproot. The root nodules are round and sometimes lobed. The leaf consists of 3 narrow, green leaflets which are silvery-green underneath. The end leaflet is larger with a longer leaf stalk. The pea shaped flowers are red and yellow and occur on branched flower stalks which stick upwards in the axils of leaves. Pods are long, straight and narrow, often with 4-8 seeds. Seeds vary in shape, size and colour. The pods are slightly hairy. Pods are often 4-8 cm long and have a beak at the end. Pods are constricted between the seeds. Many varieties of pigeon pea occur. Some are dwarf and day length neutral.



**Distribution:** A tropical plant that requires a tropical or subtropical climate. Plants grow from sea level up to about 1800 m in the tropics. They can tolerate drought and are suited to a drier climate. They can grow in places with less than 600 mm rainfall per year. They do less well in the wet tropics. They suffer in waterlogged soils and are damaged by frost. It can also tolerate heat. It will grow on poor soils cannot grow on salty soils. It can grow in arid places and suits hardiness zones 10-12.

**Use:** Young leaves, shoots and pods are eaten. The pods can be used in curries. The leaves and shoots as potherbs. Young seeds are cooked and eaten like peas. Ripe seeds are also cooked and eaten in soups and curries. Bean sprouts can be produced and eaten. Preparation of the seeds for dahl is somewhat complicated.

**Cultivation:** They are grown from seeds. It is best to sow seeds where the plants are to grow. Seeds normally germinate easily and well. Before sowing seed it helps to soak them in cold water for one day. Seeds store well if kept cool and dry. A spacing of 1.5 m x 1.5 m is suitable. Plants can be cut back and allowed to re-grow. Plants can also be grown from cuttings.

**Production:** Plants are fast growing. Pods are ready after 5 months. Mature seeds take about 8 months. Plants will often live for 3-4 years. Plants are cross pollinated by insects, or self pollinated.

**Food Value:** Per 100 g edible portion

| Edible part          | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|----------------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed                 | 10.0       | 1449      | 19.5      | 55          | -           | 15.0    | -       |
| pod (young)          | 64.4       | 477       | 8.7       | -           | -           | 2.0     | -       |
| seed (young, boiled) | 71.8       | 464       | 6.0       | 2           | 28.1        | 1.6     | 0.8     |

## Leafy greens

**English:** Goat's horns

**Local:**

**Scientific name:** *Sida cordifolia*

**Plant family:** MALVACEAE

**Description:** An erect, woody shrub that grows about 0.4-1 m high. It keeps growing from year to year. It is covered with short and long hairs that make the plant feel soft. The leaf stalk is 1-2.5 cm long. The leaves are one after the other and heart shaped at the base. They are toothed at the edge and 1.5-4.5 cm long. The flowers are yellow and occur in the axils of the leaves. The fruit are about 6-8 mm across and have 20 fine bristles on the top.



**Distribution:** A tropical plant that grows in open waste places in the tropics and sub-tropics. It is common and widely distributed in the Philippines. It grows in hot arid places with a marked dry season. It grows in places with an annual rainfall below 520 mm. It grows in dry sandy soils and can grow in salty soils. It grows below 1100 m altitude. It can tolerate shade and can grow in arid places.

**Use:** The leaves are edible when cooked.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| leaf        | 6.6        | 1296      | 24.2      | -           | -           | 79.8    | -       |

Image accessed from

[http://upload.wikimedia.org/wikipedia/commons/f/f4/Sida\\_cordifolia \(Bala\) in Hyderabad, AP W IMG 9420.jpg](http://upload.wikimedia.org/wikipedia/commons/f/f4/Sida_cordifolia_(Bala)_in_Hyderabad,_AP_W_IMG_9420.jpg)

## Leafy greens

**English:** Sticky cleome

**Local:**

**Scientific name:** *Cleome viscosa*

**Plant family:** CLEOMACEAE

**Description:** An erect annual herb about 0.3 to 1 m tall. It is sticky and has a rank smell. The leaves are made up of 3-5 leaflets each 1-3 cm long. The flowers are in leafy groups at the end of branches. The flower stalks are less than 1 cm long. The petals are yellow and 7-8 mm long. The fruit is a narrow capsule and gradually tapers near the tip. The stems and seed pods are hairy. The seeds are round, black and 1 mm across.



**Distribution:** It is a tropical plant found in waste places at low and medium altitudes. It is damaged by drought and frost. It can grow in arid places. It restricts the germination and growth of Pearl millet.

**Use:** The leaves are edible when cooked. The young fruit are eaten candied. Roasted seeds are used in curries and pickles. Seed oil is used for cooking. The leaves are soaked, fermented and used as a spice.

**Cultivation:** Plants are grown from seed.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| leaf        | 80.4       | -         | 5.6       | -           | -           | 24      | -       |

Image accessed from: [https://live.staticflickr.com/2232/2205182171\\_3fece152a5\\_b.jpg](https://live.staticflickr.com/2232/2205182171_3fece152a5_b.jpg)

## Leafy greens

**English:** Nettle tree

**Local:**

**Scientific name:** *Celtis integrifolia*

**Plant family:** ULMACEAE

**Description:** A large tree that grows to 25 m tall. The trunk is 1.5 m across and can have short buttresses. The branches occur low down. The bark is grey and smooth. The oval leaves are alternate and oblique, and taper to the tip. The twigs are green with white hairs. The flowers occur in clusters in the axils of leaves on one year old shoots. The fruit are oval, about 1 cm long, with one white seed.



**Distribution:** A tropical plant that grows in the Sahel in West Africa. It needs rainfall of 500-700 mm per year. It often grows in depressions and near waterholes. It can grow in arid places.

**Use:** The leaves are used in soups. Sometimes they are used with baobab leaves. Young leaves can be eaten in salads. The fresh fruit are eaten.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| leaf (dry)  | 10.8       | 1058      | 8.0       | -           | -           | 19.7    | -       |

Image sourced from:

[http://www.westafricanplants.senckenberg.de/images/pictures/Celtis\\_integrifolia\\_MS\\_1566\\_1790\\_39f3a2.JPG](http://www.westafricanplants.senckenberg.de/images/pictures/Celtis_integrifolia_MS_1566_1790_39f3a2.JPG)



## Leafy greens

**Common name:** Lotus-seed herb

**Local:**

**Scientific name:** *Alternanthera sessilis*

**Plant family:** AMARANTHACEAE

**Description:** A low lying and spreading plant which has many branches. It continues to grow from year to year. It has a strong taproot. The stem and branches are up to 60-100 cm long and near the ends there are 2 lines of hairs along the stem. The leaves are smooth and attached to the stem without a stalk. They are opposite. The leaves are 1-10 cm long and 0.2-2 cm wide. The flowers heads are white and 5-7 mm long. They grow along the plant and do not have flower stalks. It flowers all year round. The fruit are oval and compressed on the side. The seed is about 1.5 cm across. When plants are growing in water the stems become hollow and the plants float.



**Distribution:** A tropical plant. It grows in the lowlands and the highlands. It occurs in most tropical places. It is common in waste land at low and medium altitudes in the Philippines. It grows in open moist places from sea level to 2000 m in Papua New Guinea. In Fiji it grows from sea level to 500 m. In Nepal it grows to 2400 m altitude. It can grow in arid places. It is best in alkaline soil. It can grow in seasonally water logged soils and near rivers and ditches.

**Use:** The leaves and tender tips are cooked and eaten. They are used in soups. It is also used to prepare a cool drink. The harvested leaves can only be stored for 2-3 days.

**Cultivation:** It can be grown by dividing the underground stem. It can also be grown from sections which root at the nodes. It can be grown by seed.

**Production:** The first harvest of leaves can be taken 50-60 days after planting.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| leaf        | 89.3       | 109       | 4.5       | 57          | 77          | -       | -       |
| shoot       | -          | -         | 5.0       | 577         | -           | 16.7    | -       |

Image sourced from: [https://commons.wikimedia.org/wiki/File:Alternanthera\\_sessilis.jpg](https://commons.wikimedia.org/wiki/File:Alternanthera_sessilis.jpg)

## Leafy greens

**English:** Vegetable kenaf

**Local:**

**Scientific name:** *Hibiscus cannabinus*

**Plant family:** MALVACEAE

**Description:** A herb that can grow from seed each year, or keep growing from year to year. It grows up to 3.5 m tall. It has a few sharp spines. The leaf stalk is 6-20 cm long. The leaf blade has 2 forms. The leaves lower on the stem are heart shaped and those higher on the stem have 4-7 lobes arranged like fingers on a hand. These lobes are sword shaped and 2-12 cm long by 0.6-2 cm wide. They have teeth around the edge and taper at the tip. The flowers are yellow, white or ivory and red at the base. They occur singly in the axils of leaves. They are large and up to 10 cm across. They have very short stalks. The fruit is a capsule about 1.5 cm across. The seeds are kidney shaped.



**Distribution:** A tropical plant. It can grow in well-drained sandy soils and in dry but seasonally waterlogged places. It grows from 1500-2100 m above sea level. It grows in areas with an annual rainfall of 500-635 mm. It can grow in arid places and suits hardiness zones 10-12. It grows in many African and Asian countries.

**Use:** The leaves are eaten cooked as a vegetable. They are also used as a substitute for tamarind for curries. They are used in soups. The leaves are cooked with the aid of potashes. The seeds are roasted and eaten. They are also fermented. The seeds yield an edible oil. The flowers are eaten cooked as a vegetable. The bark is sweet and is chewed by children.

**Cultivation:** It is usually grown from seeds but can be grown from cuttings. Seeds will last for about 8 months. Seeds germinate best at 35°C.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)  | 8.1        | 1785      | 20.2      | -           | -           | -       | -       |
| leaf        | 79.0       | 280       | 5.5       | 34          | -           | 12.1    | -       |

## Leafy greens

**English:** Purple amaranth

**Local:**

**Scientific name:** *Amaranthus cruentus*

**Plant family:** AMARANTHACEAE

**Description:** An annual erect plant. It grows to 2 m high. The stems are angular. It often branches in the upper section. It is smooth but may be hairy on younger plant parts. The young parts can be tinged purple. The leaves are oval to sword shaped and can be 10-15 cm long by 3-6 cm wide. They have a leaf stalk 1-7.5 cm long. The leaves often narrow towards the tip. They can also become thinner towards the base. There may be hairs on the midrib. The leaf may be tinged purple underneath. The flowers clusters are often branched and on side branches. The stiff branched flower arrangement at the top can be 15-25 cm long. The fruit is oval and the seed can be 1-1.3 mm across. The seed is dark brown but pale brown forms are used as grain in Central America.



**Distribution:** It grows in the tropics and more temperate regions. In the tropics it grows mainly in the highlands. It needs a night temperature above 15°C and preferably a day temperatures above 25°C. It grows best in fertile, well drained soil and suits hardiness zones 8-11.

**Use:** The leaves and young plant are eaten cooked. They are also dried and stored. The seeds are ground into flour and used to make bread. **Caution:** This plant can accumulate poisonous nitrates if grown with high nitrogen inorganic fertilisers.

**Cultivation:** Plants can be grown from seed if the soil is warm. Seeds are small and grow easily. They can be put in a nursery and then transplanted after 2-3 weeks. Cuttings of growing plants root easily.

**Production:** Yields of 800-1500 kg per hectare are achieved. Plants can be harvested by pulling up the entire plant or by removing leaves over several harvests.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed        | 87.7       | 2006      | 14.7      | -           | 0           | 3.8     | -       |
| leaf        | 84.0       | 176       | 4.6       | -           | 64          | 8.9     | -       |



## Leafy greens

**English:** Indian spinach

**Local:**

**Scientific name:** *Basella alba*

**Plant family:** BASELLACEAE

**Description:** An annual or perennial climbing herb with thick fleshy leaves. The vine is smooth and juicy and can be 10 m long. It branches freely. The vine and leaves can be red or green. The leaves are fleshy and pointed at the tip. They can be 8-18 cm long and 8-10 cm across. They are carried alternately along the vine. Leaves can be heart shaped or oval. It has white, pink or red flowers in short spikes which are in the axils of the leaves. The fruit are round and soft. They can be red, white or black and are 6-8 mm across. The seeds are round and black.



They are 3 mm across. (Often the ones with heart shaped leaves are called *Basella cordifolia*, the ones with a red stem *Basella rubra* and the short day flowering dark green kind *Basella alba*.)

**Distribution:** A tropical plant. It occurs mostly in the tropical lowlands and is best below 500 m but will grow up to about 1600 m. in the equatorial tropics. It will grow quite well in the temperature range 15-35°C. It does not like water-logging but can survive 4-12 weeks drought once well established. It requires adequate water during the growing season. The best pH is 5.5-7.0. It cannot tolerate salty conditions. Flowering does not occur when day lengths are over 13 hours.

**Use:** The young shoots and leaves are eaten cooked. They are somewhat slimy. In soups and stews the mucilage can be used as thickening. The purple colour of fruit is harmless and is used to colour vegetables and agar-agar. Some lemon juice added to the dye enhances the colour. The leaves can be eaten raw in salads or cooked like a vegetable. The leaves are used to make tea and can also be dried and stored. The seeds can be crushed to use as an edible dye for jellies.

**Cultivation:** It can be sown from seeds or cuttings. Seeds germinate in a few days. Sticks can be provided for support, or it can grow over fences and stumps. If seeds are used, 3 kg of seed will sow one hectare. They are best sown in a nursery and transplanted. A spacing of 1 m is suitable. Plants grown from seed are more productive than those grown from cuttings. When cuttings are used, 20-25 cm long cuttings are suitable. Where the plant grows over light soil it can root at the nodes and continue growing. Partial shade, rich fertile soil and adequate moisture favour abundant leaf production. It is responsive to nitrogen fertiliser. Light shade gives bigger leaves. It requires a trellis to climb over. Frequently picking of the bud encourages branching.

**Production:** It is 4-6 weeks until the first harvest. It grows reasonably well on poor soils and is fairly resistant to pest and disease. Leaves will only store for one day at 20-30°C. Yields of 40 kg of leaves from a 10 metre square bed is possible over 75 days.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| leaf        | 85.0       | 202       | 5.0       | 56          | 100         | 4.0     | -       |

## Fruit

**English:** Boabab

**Local:**

**Scientific name:** *Adansonia digitata*

**Plant family:** BOMBACACEAE

**Description:** A large tree. It grows up to 25 m tall. It loses its leaves during the year. The branches are thick, angular and spread out wide. The trunk is short and stout and can be 10-14 m around. Often the trunk has deep grooves or is fluted. The bark is smooth and grey but can be rough and wrinkled. The leaves spread out like fingers on a hand. There are 5-9 leaflets. Often the leaves are crowded near the ends of branches. The flowers are large and 12-15 cm across. The petals are white and the stamens are purple. The fruit hangs singly on a long stalk. The fruit has a woody shell. This can be 20-30 cm long and 10 cm across. Inside the fruit are hard brown seeds. They are about 15 mm long. The seeds are in a yellow white floury pulp. The pulp is edible. The thick roots end in fattened tubers.



**Distribution:** It is a tropical plant that grows in the lowlands. It grows in the hot dry regions of tropical Africa, such as the Sahel. It survives well in dry climates. It grows where rainfall is 100-1000 mm a year. It can tolerate fire. It grows where the annual temperatures are 20-30°C. In most places it grows below 900 m altitude but occasionally grows to 1500 m altitude. It requires good drainage. It can grow in arid places and suits hardiness zones 11-12.

**Use:** The young leaves are eaten as a cooked vegetable. The dried leaves are also used to thicken soups. The fruit pulp is eaten raw. It is also used for a drink. The flowers are eaten raw or cooked. The seeds can be eaten fresh or dried and ground into flour then added to soups. They yield a cooking oil. The shoots of germinating seeds are eaten. The young tender roots are eaten. The fattened root tubers are cooked and eaten. The bark is eaten and the dried leaves are used as flavouring.

**Cultivation:** Trees are grown from seed. The seed remain viable for several years but before planting the seeds must be treated to break the hard seed coat, by soaking the seeds in hot water for several minutes or by cutting the seed coat. Seeds that float in water should not be used. Seeds can be planted in nurseries in plastic bags then transplanted after 6 months. Plants can also be grown from cuttings.

**Production:** Trees grow quickly reaching 2 m in 2 years. Trees produce fruit after 2-15 years. The plant is pollinated by bats, insects and winds. Trees can last 600 or more years. Fruit can be stored for about a year.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| nut (dry)   | 7.8        | 1832      | 33.7      | -           |             | 13.9    | -       |
| fruit       | 16.0       | 1212      | 2.2       | -           | 360         | 7.4     | 6.7     |
| leaf        | 77.0       | 290       | 3.8       | -           | 50          | -       | -       |

## Fruit


**English:** Mango

**Local:**

**Scientific name:** *Mangifera indica*

**Plant family:** ANACARDIACEAE

**Description:** An erect, branched evergreen tree. It can grow to 10-40 m high and is long lived. (Trees grown by vegetative means are smaller and more compact.) Trees spread to 15 m across. It has strong deep roots. The trunk is thick. The bark is greyish-brown. The leaves are simple and shaped like a spear. Some kinds of mangoes have leaves with a wavy edge. They can be 10-30 cm long and 2-10 cm wide. They are arranged in spirals. The leaf stalk is 1-10 cm long and flattened. Leaves are often brightly coloured and brownish-red when young. These tender leaves which are produced in flushes become stiff and dark-green when mature. The flower stalks are at the ends of branches. They are 10-50 cm long and branching. Up to 6000 flowers can occur on a stalk. Most of these are male and up to 35% have both male and female flower parts. Fruit are green, yellow or red and 2.5-30 cm long. The fruit hang down on long stalks. The outside layer of the seed is hard and fibrous and there is one seed inside. Several embryos can develop from one seed by asexual reproduction. The fruit shape and colour vary as well as the amount of fibre and the flavour. India has many varieties and they cannot tolerate humidity.



Leaves are often brightly coloured and brownish-red when young. These tender leaves which are produced in flushes become stiff and dark-green when mature. The flower stalks are at the ends of branches. They are 10-50 cm long and branching. Up to 6000 flowers can occur on a stalk. Most of these are male and up to 35% have both male and female flower parts. Fruit are green, yellow or red and 2.5-30 cm long. The fruit hang down on long stalks. The outside layer of the seed is hard and fibrous and there is one seed inside. Several embryos can develop from one seed by asexual reproduction. The fruit shape and colour vary as well as the amount of fibre and the flavour. India has many varieties and they cannot tolerate humidity.

**Distribution:** A tropical and subtropical plant. It grows in the lowlands. It grows from sea level up to 1300 m altitude in the tropics. It does best in areas below 700 m and with a dry season. Rain and high humidity at flowering reduces fruit set. It thrives best where temperatures are about 25°C but will grow with temperatures from 10-42°C. Temperatures of 0°C will damage young trees and flowers. Low temperatures (10-20°C) at flowering time will reduce fruiting. As temperatures get lower due to latitude or altitude, fruit maturity is later and trees become more likely to only have good crops every second year. Mangoes can grow on a range of soils. In wetter areas soils with less clay are better. They can withstand occasional flooding. A soil pH of 5.5-6.5 is best. Soils with pH above 7.5 cause plants to develop iron deficiency. It grows in the Sahel. It can grow in arid places. It suits hardiness zones 11-12.

**Use:** Ripe fruit are eaten raw. Unripe fruit is pickled. Seeds can be eaten cooked. They are boiled or roasted. They are made into meal by powdering. Young leaves can be eaten raw or cooked. Amchur is made from the dried unripe fruit. This is used in curries, and pickles and chutneys. The seed kernels are used for famine food in India. They are boiled, roasted or soaked to remove the bitterness. **Caution:** The sap from the tree or fruit can cause skin problems with some people.

**Cultivation:** Trees are grown by planting fresh seed and they can be transplanted. Mangoes vary in their ability to breed true from seed. When more than one seedling emerges from the seed some of these are asexual and breed true. Clean seed germinate best if they are treated at 50°C for 20 minutes, then planted on their edge with the round bulge upwards and near the soil surface. The husk around the seed should be removed. Seeds germinate in 3-6 weeks. The strongest growing seedlings from this seed are used and the others thrown away. The seedlings from the folds of the seed are vegetative while the seedling from the centre of the seedling near the stalk end may be sexual and show variation from type. Other seeds only produce one seedling and these normally

vary and can be different from the parent tree. Plants can be propagated by budding, or by grafting using in-arching. This is not easy and care is required. In wetter places, flowers need to be protected with fungicides to enable fruit to form. If organic manure is used this should not be directly in the planting hole nor immediately against the new plant. Young transplanted seedlings need regular watering. A spacing of 6-12 m between plants is used. Wind protection is advisable to prevent fruit rubbing and getting damaged. Trees should only ever be lightly pruned as fruit develop on new growth and heavy pruning can reduce flowering. Flowering can be brought about by foliar sprays of potassium nitrate.

**Production:** Seeds germinate after about 20 days. Seedling trees produce after 4-6 years and increase in production up to 20 years. Trees often bear better each second year. Rain at flowering reduces fruit setting. Fruiting is at the end of the year. Fruit take 4-5 months to mature. Fruit vary in weight from 200-1000 g. Trees can produce one million flowers but only 500 fruit. Trees last for many years.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| fruit       | 83.0       | 253       | 0.5       | 54          | 30          | 0.5     | 0.04    |
| leaf        | 82.1       | 226       | 3.9       | -           | 60          | 2.8     | -       |



## Fruit

**Common name:** Soursop

**Local:**

**Scientific name:** *Annona muricata*

**Plant family:** ANNONACEAE

**Description:** A low, bushy tree growing 8-10 m tall. The leaves are long (14 cm), narrow (4 cm) and thick and slightly shiny on top. The flowers are large (2-3 cm), rounded and produced on short stems on the branches. They occur singly or in groups of three. The flowers have two layers of thick, fleshy petals. The fruit are 10-30 cm long and spiny. The flesh is juicy. Many black seeds are embedded in the white flesh. Fruit are often mis-shaped, due to only some of the ovules, or young undeveloped seeds, being fertilised. Beetles are normally thought to do the pollinating. This means fruit end up heart-shaped when unevenly pollinated. Several types occur with different sweetness, shape and juiciness.



**Distribution:** A tropical plant that has been taken to most tropical countries and grows in tropical lowland areas below 1200 m altitude. It can tolerate poor soils and a humid climate. It cannot tolerate frost. The trees can withstand temperatures down to freezing (0°C) for a short time but salt-laden winds from the sea can kill the trees. It needs a well-drained soil and cannot tolerate waterlogging. Trees continue to grow and produce satisfactorily in fairly poor, compacted soil, but improving the fertility increases the amount of fruit. It can grow well in hot humid areas, but a fungus disease called Blossom blight can cause flowers to fall off. It suits hardiness zones 10-12.

**Use:** Fruit can be eaten fresh, or used in ice-cream and for drinks. Young fruit can be cooked as a vegetable. Leaves are edible when cooked. The trees are fairly common, but the fruit may not be widely used. **Caution:** The seeds are toxic, so should be removed before processing.

**Cultivation:** Trees are grown either as seedling trees or grafted plants. They can be grown from cuttings or air-layering (part of the plant cut and wrapped in dirt so it produces new shoots). Trees are easy to grow and maintain. Plants can easily be grown from seeds. Seeds can be planted fresh or stored. Seeds grow in about 15-20 days. Trees grown from seeds vary in the quality of the fruit. Seedlings are transferred to plastic bags when 15 cm tall. Trees can also be grown from cuttings or by grafting. This allows better trees to be selected and produced. Seedlings are suitable for grafting after 6 months. Trees need to be about 5 m apart. Flowers are pollinated by insects. Hand-pollination of flowers can increase the number of fruit that are produced. Fruit are soft and fleshy, and difficult to transport.

**Production:** Trees grow quickly and commence bearing by the third year. It bears fruit almost continually throughout the year, but there is normally one season when more fruit are getting ripe. A tree can produce 12-24 fruit in a year, weighing 4-5 kg each and with 11-14% sugars.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| fruit       | 82.4       | 294       | 0.88      | -           | 16          | 0.3     | 0.1     |

## Fruit

**English:** Milk plum

**Local:**

**Scientific name:** *Englerophytum magalismsontanum*

**Plant family:** SAPOTACEAE

**Description:** A shrub or small tree that usually grows up to 15 m tall, although can occasionally grow up to 33 m. Often the roots are twisted around a rock. It has a leafy rounded crown. The trunk is thin and has flutes along it. The stem is mostly short and thick and the branches can touch the ground. Often the trunk has wart-like growths. The bark is grey and rough. The firm, leathery leaves are simple and alternate and are often crowded towards the ends of branches. The leaves are oval and dark green and shiny on top but brown and hairy underneath. The



young shoots and twigs have dense rusty hairs. The upper leaf surface often has a bloom. The under surface of the leaves has silver or red hairs. The leaves are usually 5-10 cm long but can be 30 cm long. The leaf stalks are 0.6-2.5 cm long. Flowers and fruit are produced on old wood. The flower buds occur in clusters, often on knobs on the branches and trunk. The fruit are 2.5 cm long. They are bright red when ripe. There is a small point at the end. The fruit have a leathery skin and 2 stones inside. The flesh is purple and they have a sticky latex. The seeds are oval and 1.7 x 1.4 cm.

**Distribution:** A tropical plant that occurs near quartzite and granite rocks. It can grow in the lowlands and up to 1800 m in South Africa. They mostly grow in coastal areas in deciduous woodland. They are somewhat drought resistant and can tolerate frost. They grow on a range of soils. They do best on dry well drained soils. It can grow in arid places.

**Use:** The fruit can be eaten raw when immature or when ripe. They are also used for beer, wine and jams.

**Cultivation:** They can be grown from seed or cuttings. Fresh seed germinate quickly. Cuttings from old wood can be used. Fruit are ready to harvest 4-5 months after flowering.

**Production:** Trees grow fairly quickly. In Tanzania, fruit are collected from April to June.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| fruit       | 74.3       | 404       | 0.9       | -           | 13.1        | 0.3     |         |

## Fruit

**English:** Pawpaw

**Local:**

**Scientific name:** *Carica papaya*

**Plant family:** CARICACEAE

**Description:** Pawpaw is a tropical fruit that grows 3-5 m tall and only occasionally has branches. The stem is softly woody with scars from fallen leaves along it. There is a clump of leaves at the top of the plant. The leaves are large (50 cm wide) deeply lobed and on leaf stalks up to 90 cm long. Trees can be male, female or bisexual. Male flowers are small and white and on long stalks. Female and bisexual flowers are on short stalks. These have no fruit, round fruit and long fruit respectively. There are three forms of long fruit. The seeds are black.



**Distribution:** It is a tropical plant that grows from sea level up to about 1700 m altitude in the equatorial tropics. In cooler regions they have to be planted but in humid tropical regions are commonly self-sown. Sunlight allows germination when forest is cleared. It cannot stand frost. It needs a night temperature above 12°C and don't tolerate water-logging. Plants die after 48 hours in standing water. It needs a pH between 5-8 and suits hardiness zones 11-12.

**Use:** Fruit can be eaten ripe and raw. Green fruit can be cooked as a vegetable. The young leaves can be eaten cooked, but are bitter. The flowers and the middle of the stem can be eaten. Papayas contain papain which is a meat tenderiser. The dried seeds can be used as a spice.

**Cultivation:** Pawpaw seeds grow easily and plants grow quickly. Fresh seeds can be used. If dry seeds are used they should be soaked before planting. Seeds should be sown when temperatures are 24-30°C. They need a reasonably fertile soil. Seeds can be sown directly or the seeds can be put in a nursery and the seedlings transplanted. Seeds in a nursery should be sown 1-2 cm deep. Seedlings can be transplanted when they are about 20 cm high. Plants should be about 3 m apart. Continuous fruit production depends on fertility, temperature and moisture being adequate to maintain active growth. The fruit is produced year round but the growth and development rate decreases with temperature. The size and quality of fruit declines at lower temperatures. Pollination is by wind and insects and is not normally limiting. Normally cross and self-pollination both occur. Seeds are dispersed by birds, bats and people and remain viable for a few months.

**Production:** Seeds emerge in 2-3 weeks. Vegetative growth before flowering is 4-8 months. One or more fruit grow per leaf axil, about every 1-2 weeks under good growing conditions. With good growth, 100 fruit can be produced from one plant in a year. Pollination to maturity is about 2-3 months. On the coast in tropical equatorial regions, pawpaws start producing fruit after about 4-5 months, but in the highlands this may take 12-18 months. The first fruit are ready 6-11 months from planting. Tree life is about 2-3 years, although they may live for 10-12 years.

**Food Value:** Per 100 g edible portion

| <b>Edible part</b> | <b>Moisture<br/>%</b> | <b>Energy<br/>kJ</b> | <b>Protein<br/>g</b> | <b>proVit A<br/>µg</b> | <b>proVit C<br/>mg</b> | <b>Iron<br/>mg</b> | <b>Zinc<br/>mg</b> |
|--------------------|-----------------------|----------------------|----------------------|------------------------|------------------------|--------------------|--------------------|
| leaf               | 75.4                  | 378                  | 8.0                  | -                      | 140                    | 0.77               | -                  |
| fruit              | 88.0                  | 163                  | 0.5                  | 290                    | 54                     | 0.4                | 0.18               |
| fruit (unripe)     | 92.1                  | 109                  | 1.0                  | -                      | -                      | 0.3                | -                  |



## Fruit


**English:** Watermelon

**Local:**

**Scientific name:** *Citrullus lanatus*

**Plant family:** CUCURBITACEAE

**Description:** An annual climber, with deeply divided leaves and tendrils along the vine. It trails over the ground and has hairy, angular stems. The leaves are on long leaf stalks. The leaves are deeply divided along their length. These leaf lobes are rounded and can themselves be divided. The leaves are 5-20 cm long by 2-12 cm across. The tendrils are divided. The plant has separate male and female flowers on the same plant. The flowers are pale yellow and smaller than pumpkin flowers. The flowers occur in the axils of leaves. The male flowers appear first.



Fruit are large and round or oval. They can be 60 cm long. Fruit have a hard smooth skin. Several fruit colours and shapes occur. They often have a dark green mottle, or blotches. The fruit has reddish, juicy flesh and black or red seeds. The seeds are oval-shaped and smooth.

**Distribution:** It grows in most tropical and subtropical countries. It grows best on the coast in the tropics, but will grow up to about 1000 m altitude. It will not stand water-logging and does well on sandy soils. Plants are frost-sensitive. Seed will not germinate below 21°C. Temperatures between 24-30°C are suitable. Fruit are sweeter in arid warm areas. It suits hardiness zones 10-12.

**Use:** The fruit is eaten raw when ripe. Small, unripe fruit can be cooked as a vegetable. The skin is sometimes candied in vinegar and eaten with fish. Seeds are also eaten. They are dried, soaked in salt water, then roasted. Oil is extracted from the seeds. Very young leaves are occasionally eaten. It is a popular fruit.

**Cultivation:** They are suitable mainly for the dry season. A spacing of 1.5-2 m is suitable. They grow easily from seed. They do best when fully exposed to the sun. Seed can be dried and stored. If too much vegetative growth occurs, picking out the tip to produce side branches will produce more fruit.

**Production:** Harvesting commences after 4-5 months. The main fruit season is November to January. The ripeness can be determined by tapping the fruit to get a dull sound. The part of the fruit on the ground changes from green to light yellow and the tendril near the base of the fruit becomes dry when ripe. Fruit yield can be 45-60 t/ha.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| fruit       | 94.0       | 92        | 0.4       | 20          | 5           | 0.3     | 0.1     |
| seed        | 5.1        | 2330      | 28.3      | 0           | 0           | 7.3     | 10.2    |

## Fruit

**English:** Rambutan

**Local:**

**Scientific name:** *Nephelium lappaceum*

**Plant family:** SAPINDACEAE

**Description:** A medium sized tree 12-25 m tall. It keeps its leaves throughout the year. Trees spread to 6 m wide. The leaves are alternate with leaflets opposite along the stalk. There are 5-7 pairs of leaflets. Each leaflet can be 20 cm long by 8 cm wide. The leaflet stalks are short and the leaflets are pointed at the tip and rounded at the base. The flowers are very numerous on compound flower arrangements. Seedling trees can produce only female or only male trees. Some kinds have both male and female



flowers or both flower parts in the one flower. The dark red fruit has many prominent projections united at the base. The colour can vary between yellow, green, orange and bright red. Some trees are separately male and female, and sometimes seedless fruit is set. Male and female flowers are often separate on the same tree. The fruit is up to 5 cm long in loose clusters of 10-12. It normally contains one large seed. There are several named cultivated varieties.

**Distribution:** A tropical plant that suits the humid tropical lowlands mostly within 12° of the equator. It grows from sea level up to about 500 m in equatorial zones and will grow about 17° from the equator. It does best where the temperature is constant around 28°C or ranges from 22-30°C with a rainfall of 2000-5000 mm per year. Low humidity can cause leaf scorch. Excessive rainfall before flower formation reduces flowering. A dry period of about one month assists flower initiation and areas with two seasonal dry periods can get double crops. A well drained soil is needed.

**Use:** The fruit are eaten fresh. Sour fruit are eaten stewed. The seeds are sometimes roasted and eaten, and can be used for oil.

**Cultivation:** Trees can be grown from fresh seeds. The short lived seeds need to be planted fresh, after washing. Germination takes place in 7-10 days. Seedlings are not easy to transplant. Roots are sensitive to drying out. Selected trees can be propagated by budding or air-layering. Seedling tree quality varies considerably due to cross pollination. Trees are planted 10 m apart at densities of 80-120 per ha. About 1% male flowers are sufficient if enough pollinating insects are present. A mixture of varieties might be needed to get good fruit set. Organic manure is often used twice yearly.

**Production:** Seedling trees can bear after 6 years and vegetative trees after 4 years. Flowering occurs during the dry season. The main fruit season is November to March. Fruit mature about 110 days after flowering. Where trees with only female flowers occur, male pollinator trees are needed. Fruit can weigh 20-40 g and need to be harvested when ripe. A well maintained tree can produce 150 kg.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| fruit (raw) | 82.0       | 271       | 1.0       | 0           | 53          | 1.9     | 0.1     |

## Fruit

**English:** Avocado

**Local:**

**Scientific name:** *Persea americana*

**Plant family:** LAURACEAE

**Description:** A small to medium sized tree that normally grows 8-10 m tall, but can reach 25 m. The leaf stalk is 1.5-5 cm long. Leaves are entire, oval and 5-40 cm long. Flowers are greenish, small and on the ends of branches. Clusters of flowers may contain 200-300 flowers. Normally only 1-3 fruit develop from each cluster. The fruit is round or pear shaped, and 7-20 cm long. The fruit are greenish-yellow with some red coloration. The fruit has greenish-yellow flesh and a large round seed. There are 3 named races-West Indian, Guatemalan and Mexican.



**Distribution:** A subtropical plant that grows from sea level up to 2,250 m in the tropics. It cannot stand water-logging. Branches are easily damaged by wind. It needs a frost free location or where frosts are rare. West Indian varieties thrive in humid, tropical climates, freeze at or near 0°C and can stand some salinity. Mexican types come from dry subtropical plateaus and thrive in a Mediterranean climate. They are hardy to -7° C. They are salt sensitive, have the smallest fruits and the thinnest skin. The best daytime temperature is 25-33°C. Guatemalan types come from cool, high-altitude tropics and are hardy to -3° C. It does best with neutral or slightly acid and well aerated soil. Growth is disrupted when soil temperature is below 13°C. It needs high humidity at flowering and fruit set. It can grow in arid places.

**Use:** The fruit pulp is eaten raw or cooked. It is used in salads, soups, sandwiches, spreads, ice cream, and also in tortillas and wine. The fruit is mixed with sugar and water to make a drink. Oil is extracted from the flesh and is used in salad dressing. The leaves can be used for tea sweetened with sugarcane juice. Toasted leaves are used to season stews and bean dishes. **Caution:** Some people are allergic to avocado.

**Cultivation:** Plants are often grown from seed. Seeds remain viable for 2-3 weeks. Fresh seed held at 25°C day to 15°C night will germinate in 3 weeks. It is best to propagate vegetatively. Tip cuttings, layers and grafts can be used. Because different types have pollen at different times of day, a mixture of trees which have pollen and flowers receptive at different times gives best fruit set. Although trees will grow in shade, they need sun for fruiting. The leaves do not rot easily and can accumulate under trees. Other plants cannot be grown under avocado trees.

**Production:** Seedlings grow quickly and continuously in warm, moist conditions. Seedlings bear after 5-8 years. Grafted trees can fruit in 1-2 years. A good tree produces 400-600 fruit each year. A fruit can weigh 50 g-1 kg. In the subtropics, trees often produce 2 main flushes of fruit per year. From fruit set to maturity can take 6-12 months. Fruit ripen off the tree in 4-14 days. For the Mexican types, the fruit weigh less than 250 g and they ripen 6-8 months after flowering.

**Food Value:** Per 100 g edible portion

| <b>Edible part</b> | <b>Moisture<br/>%</b> | <b>Energy<br/>kJ</b> | <b>Protein<br/>g</b> | <b>proVit A<br/>µg</b> | <b>proVit C<br/>mg</b> | <b>Iron<br/>mg</b> | <b>Zinc<br/>mg</b> |
|--------------------|-----------------------|----------------------|----------------------|------------------------|------------------------|--------------------|--------------------|
| fruit              | 74.4                  | 805                  | 1.8                  | 480                    | 11                     | 0.7                | 0.4                |

Image accessed from:

[https://upload.wikimedia.org/wikipedia/commons/7/7d/Avocados\\_\(Persea\\_americana\)\\_ \(18159574242\).jpg](https://upload.wikimedia.org/wikipedia/commons/7/7d/Avocados_(Persea_americana)_ (18159574242).jpg)

## Vegetables

**English:** Horseradish tree

**Local:**

**Scientific name:** *Moringa oleifera*

**Plant family:** MORINGACEAE

**Description:** A small, soft-wooded tree that grows 9-12 m tall. The tree loses its leaves during the year. The bark is grey, thick, corky and peels off in patches. The leaves are pale green and the leaf is divided 3 times. The whole leaf is 30-60 cm long and the leaflets are usually oval and 1-2 cm long. The leaflets are jointed with a gland near the joint. The flowers are pale yellow. They occur in long sprays 30 cm long. Each flower has 5 petals and of these one is erect and 4 are bent backwards. The fruit is a long capsule 30-100 cm long by 2 cm wide. The seed capsules are up to 45 cm long. They are roughly triangular in shape. The seeds have 3 wings. Often the fruiting kinds are grown as annual plants.



**Distribution:** A tropical and subtropical plant. They suit the dry lowland areas and grow up to 1,350 m altitude in the tropics. They are not hardy to frost. They cannot tolerate water-logging. A pH of 6-7.5 is suitable. It can grow in arid places. It suits hardiness zones 9-12.

**Use:** The young tops and leaves are eaten cooked. They are eaten as potherbs or used in soups and curries. They can be dried and stored for later use. The very young long pods are eaten cooked, especially in curries and soup. They are also pickled. The young seeds are eaten roasted or fried. Sometimes the roots are used as a horseradish substitute. A gum from the bark is used as seasoning. The bark is used for tea. The roots, leaves, flowers and fruits are eaten cooked in water and mixed with salt and chili peppers. The oil expressed from the seeds is used in salads.

**Cultivation:** It is best to grow plants from 1 metre long cuttings but they can be grown from seed. They can be used as a hedge and pruned regularly to produce more leaves. Properly dried seed can be stored for a long time in sealed containers in a cool place. Normally perennial types are grown from cuttings and annual types are grown from seed.

**Production:** Trees are fast growing. They can be pruned or topped. With one variety the tree flowers and fruits continuously while with the other variety there are flowers and fruit once per year. The fruit ripens 3 months after flowering. Annual types produce fruit 6 months after planting. Leaves are best dried in the shade to retain more of their Vitamin A.

**Food Value:** Per 100 g edible portion

| Edible part   | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|---------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| leaf          | 76.4       | 302       | 5.0       | 197         | 165         | 3.6     | -       |
| flower        | 84.2       | 205       | 3.3       | -           | -           | 5.2     | -       |
| leaf (boiled) | 87         | 189       | 4.7       | 40          | 31.0        | 2.0     | 0.2     |
| pod (raw)     | 88.2       | 155       | 2.1       | 7           | 141         | 0.4     | 0.5     |
| seed          | 6.5        | -         | 46.6      | -           | -           | -       | -       |



## Vegetables

**English:** Marrow

**Local:**

**Scientific name:** *Cucurbita pepo*

**Plant family:** CUCURBITACEAE

**Description:** A bristly hairy annual vine in the pumpkin family. It has branched tendrils. The stems are angular and prickly. The leaves are roughly triangular. The leaves have 5 lobes which are pointed at the end and are toothed around the edge. Male and female plants are separate on the same plant. Male flowers are carried on long grooved flower stalks. Female flowers are borne on shorter more angular stalks. The fruit stalks have furrows along them but are not fattened near the stalk. The fruit vary in shape, size and colour. Often they are oval and yellow and 20 cm long by 15 cm wide. The seeds are smaller than pumpkin and easy to separate from the tissue. The scar at their tip is rounded or horizontal, not oblique. There are a large number of cultivated varieties.



**Distribution:** A subtropical plant. They are more suited to drier areas. They are frost sensitive, and grow best with day temperatures between 24-29°C and night temperatures of 16-24°C. It suits tropical highland regions. It suits hardiness zones 8-11.

**Use:** The young fruit are cooked and eaten. They can be steamed, boiled or fried. They are used in pies, soups, stews and cakes. The young leaves and the ripe seeds can also be eaten cooked. The seeds are dried, salted and toasted and eaten as a snack food. The seeds can also be pressed to produce oil. The sprouted seeds are used in salads. Flowers and flower buds can be eaten boiled. They can be dried for later use.

**Cultivation:** They are grown from seeds. The seeds germinate after one week. They can be grown from cuttings. They are best planted on mounds. A spacing of 2-3 m between plants is needed. Hand pollination assists fruit setting. Plants can also be grown from cuttings as plants root at the nodes.

**Production:** The first usable immature fruit are ready 7-8 weeks after planting.

**Food Value:** Per 100 g edible portion

| Edible part          | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|----------------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)           | 3.7        | 2266      | 29.4      | -           | -           | 7.3     | -       |
| leaf                 | 89.0       | 113       | 4.0       | 180         | 80          | 0.8     | -       |
| fruit (mature)       | 92.0       | 105       | 1.6       | 17          | 16          | 2.4     | -       |
| fruit                | 91.3       | 102       | 1.1       | -           | 12          | 0.8     | 0.2     |
| yellow fruit         | 92.0       | 97        | 1.0       | 180         | 8           | 1.4     | -       |
| immature fruit (raw) | 92.0       | 92        | 1.5       | -           | 9           | 0.4     | 0.1     |

## Vegetables

**Common name:** Snake gourd

**Local:**

**Scientific name:** *Trichosanthes cucumerina*

**Plant family:** CURCUBITACEAE

**Description:** Snake gourd is a member of the pumpkin family. It is a climber with tendrils. It grows up to 6 m high and spreads 1.5-3 m wide. The vine has furrows along the stem. The leaves have 3-7 lobes and a toothed edge. The flowers are white. The female flowers are produced singly on long stalks. The fruit tend to curve and can be 1-2 m long. When ripe they turn orange or red, but are grey and green when young.



**Distribution:** Snake gourd is a tropical plant, common in the humid tropical lowlands up to 500 m. It does not tolerate dry soil and requires a good moisture reserve in the soil, but is sensitive to water-logging. The optimum temperature for growth is 30-35°C with a minimum of 20°C.

**Use:** The long immature fruit of snake gourd are eaten once cooked. Sometimes a bitter taste occurs but this disappears with boiling. They can be baked, stuffed or used in curries and stews. The fruit are inedible when mature. The young leaves can be eaten once cooked.

**Cultivation:** Snake gourd is grown from seed. Seeds germinate after 10 days. Seed can be sown in a nursery and transplanted at the 2 true leaf stage. More commonly plants are sown where they are to grow. A spacing of 60-100 cm is suitable. Plants need supports to climb over. In home gardens, plants can be trained up a house wall. Plants respond to fertiliser but excessive nitrogen can produce too much vegetative growth. Pruning can improve the female to male flower ratio.

**Production:** Flowering starts 5 weeks after planting. Male flowers appear first then female flowers appear 3 days later. Pollination is normally by insects. Harvesting of fruit starts 6-7 weeks after planting and continues for 1-2 months. Fruit are picked 12-20 days after fruit set when they are 30-60 cm long. Fruit do not store well but can be stored for 10-14 days at a temperature of 16-17°C and relative humidity of 85-90%. By the time the fruit turn orange they are too mature to eat.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| fruit       | 95         | 76        | 0.9       | 810         | 6           | 1.0     | -       |
| seed        | 97         | -         | 0.2       | -           | -           | 0.2     | 0.1     |

Image accessed from: <https://tse3.mm.bing.net/th?id=OIP.Qi54QN2ROhlj60BD9KXeNgHaFN&pid=Api>



## Vegetables

**English:** Common bean

**Local:**

**Scientific name:** *Phaseolus vulgaris*

**Plant family:** FABACEAE

**Description:** There are many bush and climbing varieties of this bean. Climbing forms can be 2-3 m tall. Bush types are 20-60 cm tall. The leaves have three leaflets, one after another along the stem. The leaf stalk has a groove on the top. The side leaflets are unequal in shape, and can be 8-15 cm by 5-10 cm. The flowers are in the axils of leaves (where the leaves join the stem) and occur in a loose form. Flowers are white to purple. Pods are smooth, slender and 8-20 cm long by 1-1.5 cm wide. They are straight or slightly curved with a beak at the end and often have 10-12 coloured, kidney-shaped seeds.



**Distribution:** It is a temperate plant that grows in many temperate and subtropical countries, including Solomon Islands. It mostly grows from 700-2000 m altitude in the tropics. It suffers from pest and disease damage in the lowlands, but can be grown to sea level. It is not suited to the wet tropics. It is shallow-rooted and damaged by excess moisture near the roots. A crop lifecycle needs about 350 mm of water. It is sensitive to frost and high temperatures. Flowers will not form below 9.5°C. Night temperatures above 37°C cause flowers to drop. The best temperature range is 15-21°C. It does not suit very acid soils. It suits hardiness zones 8-11.

**Use:** The young pods, leaves and mature seeds are edible. Dry seeds are soaked in water and boiled until soft.

**Cultivation:** Plants are grown from seed, preferably sown in raised beds. Seeds remain viable for 2 years. Germination is normally good if seed has been well stored. Climbing types need stakes. Plants are self-fertilised. These beans are intercropped with other plants in many places. If grown on their own, bush types can be spaced at 25 cm x 25 cm. They can be sown closer together in rows wider apart to make weeding and harvesting easier. For dried beans, once the pods are mature and turning yellow, the whole plants are pulled, then dried and threshed. About 50-75 kg of seed will sow a hectare. Flowering in most French bean varieties is not affected by day length.

**Food Value:** Per 100 g edible portion

| Edible part  | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|--------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)   | 10.0       | 1386      | 25.0      | 10          | 1           | 8.0     | 2.8     |
| seed (young) | 92.0       | 142       | 3.0       | -           | 20          | 0.8     | 0.2     |
| pod          | 88.0       | 151       | 2.5       | 750         | 27          | 1.4     | 0.2     |
| sprout       | 90.7       | 121       | 4.2       | -           | 38.7        | 0.8     | 0.4     |

## Vegetables

**English:** Pumpkin

**Local:**

**Scientific name:** *Cucurbita maxima*

**Plant family:** CUCURBITACEAE

**Description:** A pumpkin family plant. It is a creeping vine with tendrils. It is an annual plant. The stems are soft and round in cross section. The leaves are large and hang loose. They are dark green and kidney shaped. The edges of the leaves are entire. There are large nodes at the base of the leaf. The tendrils are fairly stout and are divided half way along their length into many branches. Male flowers are carried on long upright stalks. The 5 petals are united into a long yellow tube. The female flowers are larger



than the male and are fewer in number and carried on shorter stalks. The fruit varies in size, colour and patterns on the skin. They can be round, oval or flattened, with yellow, orange or green skin. The surface can be smooth or rough and warty. The flesh is yellow and edible. The seeds are in the centre. The seeds are white or brown. They are flattened but plump and have a slanting scar at the top. The seeds are edible. (*C. moschata* does not have hairy stems but has fruit with a thickened stalk near where it joins the fruit.) There are a large number of cultivated varieties.

**Distribution:** A subtropical plant that grows from sea level to 2400 m altitude. They need a fertile soil. *C. moschata* is better suited to coastal areas. They are frost sensitive but better suited to cooler areas than *C. moschata*. It can grow in arid places. It suits hardiness zones 8-11.

**Use:** The young leaf tips are eaten cooked. They can also be dried and stored. The fruit can be eaten cooked. They are baked, boiled, fried, steamed or mashed. They are used in pies and cakes. The seeds are edible, raw or roasted. They are also ground into a meal. The male flowers are eaten after removing the stamen and calyx.

**Cultivation:** They are grown from seed. Usually 2 or 3 seeds are planted together in a mound. The distance apart depends on the cultivar. Some kinds are better for leaf tips. It is good to save seed of adapted varieties.

**Production:** Fruit are ready for harvest after about 3-4 months. Seed can be saved from fruit for re-sowing, but as pumpkins cross-pollinate, different types become mixed.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)  | 6.9        | 2264      | 24.5      | 38          | 1.9         | 14.9    | 7.5     |
| fruit       | 69.6       | 439       | 1.4       | -           | -           | -       | -       |
| leaf        | 88.0       | 160       | 4.9       | 260         | 28          | 2.5     | 0.9     |
| flower      | 88.7       | 107       | 1.4       | 173         | 14          | 0.8     | 0.1     |

## Vegetables

**English:** Bitter cucumber

**Local:**

**Scientific name:** *Momordica charantia*

**Plant family:** CUCURBITACEAE

**Description:** A pumpkin family plant. It is a slender annual climber with flowers of both sexes on the one plant. It has simple tendrils and vines can be 4 m long. It has bright green lobed leaves 5-12 cm long on thin leaf stalks 3-10 cm long. The flowers have a sweet smell and 5 small, yellow petals. Fruit are green when young and orange when ripe. The fruit have a lumpy appearance, with ridges along its length and when fully ripe burst open. It has bright red covering on the seeds inside. The seeds are pale brown and 10-16 mm long and 7-10 mm wide. Considerable variation in the fruit occurs between varieties.



**Distribution:** A tropical plant that grows from sea level up to about 500 m and will probably grow to 1000 m altitude in tropical regions. They require a well-drained soil preferably rich in organic matter. Seeds do not germinate below 15°C. Plants grow best with temperatures of 18-35°C. A soil pH of 6.5 is best. It suits hardiness zones 9-12.

**Use:** The young bitter fruit are cooked and eaten. They are boiled, stuffed, fried or pickled. They are used in soups, stews and stir-fried dishes. The seed mass of the ripe fruit is used as a food flavouring. The leaves are also cooked and eaten as a flavouring. The tender shoots and leaves are sometimes eaten. **Caution:** The leaves are considered to cause diarrhoea and vomiting.

**Cultivation:** Plants are grown from seed. For large scale plantings, 6-7 kg of seed are required for planting one hectare. Seeds are planted at 50 cm spacing in the place where the plants are to grow and need a stick to climb up. Often plants are grown on raised beds 2 m apart with 0.5 m between plants. The seed has a hard seed coat and germinates slowly. Soaking seeds for 24 hours before sowing gives a quicker more even germination. Regular watering is required.

**Production:** Fruit are ready to harvest 45-55 days after planting. Fruit should be harvested when young and tender. Once fruit have begun to change colour to yellow they are past maturity for eating. Early removal of young fruit also ensures continuous fruit setting. This can allow 6-8 successive pickings of fruit. Fruit on the plant are sometimes wrapped in paper to prevent fruit fly damage. Seed well stored can remain viable for 4-5 years. The young bitter fruit are cooked and eaten. The fruit is blanched or soaked in salt water to reduce the bitter taste.

**Food Value:** Per 100 g edible portion

| Edible part       | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed              | 8.6        | 2020      | 18.6      | -           | -           | -       | -       |
| leaf (raw)        | 84.7       | 252       | 5.0       | 44          | 170         | 7.1     | 0.3     |
| leaf tip (boiled) | 88.7       | 146       | 3.6       | 173         | 57          | 1.0     | 0.3     |
| fruit             | 93.6       | 105       | 1.2       | -           | -           | 0.2     | -       |
| pod (boiled)      | 94.0       | 79        | 0.8       | 11          | 33          | 0.4     | 0.8     |
| pod (raw)         | 94.0       | 71        | 1.0       | 380         | 84          | 0.4     | 0.8     |

## Vegetables

**English:** African horned cucumber

**Local:**

**Scientific name:** *Cucumis metuliferus*

**Plant family:** CUCURBITACEAE

**Description:** An annual pumpkin family plant. It grows to 0.5 m tall and spreads to 1.5 m wide. The stems are trailing and hairy. The tendrils are curled and do not branch. The leaves have 3 lobes and are heart shaped. The edges of the leaves have teeth. The flowers are funnel shaped and yellow. They open into five lobes. Male and female flowers occur on the same plant. The female flower is above a prickly green ovary which enlarges to become the fruit. The fruit are oblong and spiny and change from green to orange as they ripen. They are 12 cm long and 6 cm across. The fleshy pulp surrounding the seeds is bright green. The seeds are white. It tastes and smells like a cucumber. The skin is not eaten.



**Distribution:** A tropical plant. It requires conditions similar to cucumbers. It prefers light well drained soil. It needs a protected sunny position. It is drought and frost tender. It grows at low and medium elevations in Zimbabwe. It grows from 210-1800 m above sea level. It can grow in arid places.

**Use:** The peeled fruit are eaten raw. They can also be cut open and sun dried for storage and use in preserves. The young leaves are stripped and cooked then eaten. The seeds are pounded and used for flavouring. **Caution:** If the plants are bitter they need to be used with caution.

**Cultivation:** Plants are grown from seed. They take 8-16 days to germinate in warm soil.

**Production:** Plants grow rapidly. They flower and set fruit in 6-8 weeks. It is then 7 weeks to maturity.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| fruit       | 91.0       | 136       | 1.1       | -           | 18.6        | 0.5     | 0.3     |



## Vegetables

**English:** Jute

**Local:**

**Scientific name:** *Corchorus olitorius*

**Plant family:** MALVACEAE

**Description:** An annual plant. It is upright, branching, and slightly woody. Plants vary in height, shape, leafiness and hairiness. Plants grown for leaves are usually only 30 cm tall. They also have many branches. Leaves are shiny and have leaf stalks. The leaves have teeth along the edge. The tips of the lowest leaves in each side, have long bristle like structures. Small clusters of yellow flowers grow in the axils of the leaves. The fruit are ridged capsules. They can be 7 cm long. These have partitions across them between the seeds. A ripe capsules contains 180-230



seeds. The seeds are dull grey and with four faces and one long point. Each seed has one pale line along it.

**Distribution:** A tropical plant. It is mostly coastal, below 250 m altitude. Temperatures of 22-35°C are suitable. It can stand both drought (2-3 weeks) and water-logging, except when young. A well-drained soil is best. They require humus-rich soils. A soil pH of 5.5-7.0 is best, but they can grow in soils with pH up to 8.5. They also need adequate moisture for good leaf production. A rainfall of 1000 mm is suitable. A high relative humidity (80-90%) is best. It produces seeds when day lengths are short. It grows in most African and Asian countries.

**Use:** The young leaves and stem tops are eaten cooked. They are slimy unless fried. They are also used to make a thick soup. Leaves can be sun dried, pounded to flour, then stored for a long time.

**Cultivation:** Plants grow from seed, and they can be transplanted. Seeds are often broadcast into fine seed beds at the beginning of the wet season. Mixing the small seeds with sand makes it easier to sow them evenly. Often seeds are slow to start growing. This can be overcome by soaking them in hot water. A spacing of 20-30 cm between plants is suitable. For vigorous varieties this could be increased to 45-50 cm. Seeds are saved from pods for re-sowing.

**Production:** First leaves can be harvested after 5-6 weeks. Tips about 20-30 cm long are picked. Production of edible green tips, is not large. 7-8 kg of leaf tips can be harvested from 3-8 pickings over 3-4 months. Seeds can be collected after 13-15 weeks. If seeds of a particular variety are desired, it is necessary to grow these plants 16 m away from other plants, to avoid cross pollination. Seeds can be stored for 8-12 months in well-sealed jars.

**Food Value:** Per 100 g edible portion

| Edible part   | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|---------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| leaf (raw)    | 80.4       | 244       | 4.5       | 574         | 80          | 7.2     | -       |
| leaf (cooked) | 87.2       | 155       | 3.4       | 156         | 33.0        | 3.1     | 0.8     |

## Nuts, seeds, herbs and other foods

**English:** Peanut

**Local:**

**Scientific name:** *Arachis hypogea*

**Plant family:** FABACEAE

**Description:** Peanuts grow on spreading bushy plants up to about 40 cm high. The leaves are made up of 2 pairs of oppositely arranged leaflets. Flowers are produced in the axils of the leaves. Two main kinds of peanuts occur. The runner kind (Virginia peanut) has a vegetative or leafy branch between each fruiting branch and therefore produces a spreading bush. The bunch type (Spanish-Valencia peanuts) produces fruiting branches in a sequence one after the other along the branches. They grow as a more upright plant and grow more quickly. Pods are produced on long stalks which extend under the ground and they contain between 2-6 seeds. The stalk or peg from the flower grows down into the soil and then produces the pod and seed under the ground. The flowers need to be no more than 18 cm from the soil surface for the seed pod to develop underground.



**Distribution:** Peanuts grow in tropical and subtropical areas. They grow well from sea level up to about 1650 metres in the equatorial tropics. They require temperatures of 24-33°C. Plants are killed by frost. They need a well-drained soil and cannot stand water-logging and often require raised garden beds. Peanuts need 300-500 mm of rain during the growing season. Dry weather is needed near harvest.

**Use:** The seeds can be eaten raw, cooked or sprouted. They are boiled, steamed, roasted, salted or made into peanut butter or flour. The young leaves and unripe pods are edible after cooking. An edible oil is extracted from the seeds. The remaining meal can also be eaten.

**Cultivation:** Peanuts require soil with good levels of calcium and boron or they produce empty pods. Peanuts have nitrogen fixing root nodule bacteria and therefore can give good yields in soils where nitrogen is low. The nuts are normally removed from the shell before planting and are sown 2-3 cm deep, with 10 cm between plants and 60-80 cm between rows. The soil needs to be weeded and loose by the time the flowers are produced to allow the peg for the seed pods to penetrate the soil.

**Production:** Flowering can commence in 30 days and it takes 3.5-5 months until maturity. Peanuts are harvested by pulling out the plant when the top of the plants die down. After harvesting, they should be left to dry in the sun for 3-4 days. Virginia peanuts have a longer growing season and the seeds need to be stored for 30 days before they will start to re-grow.

**Food Value:** Per 100 g edible portion

| Edible part  | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|--------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)   | 4.5        | 2364      | 24.3      | -           | -           | 2.0     | 3.0     |
| seed (fresh) | 45         | 1394      | 15        | -           | 10          | 1.5     | -       |
| leaf         | 78.5       | 228       | 4.4       | -           | -           | 4.2     | -       |



## Nuts, seeds, herbs and other foods

**English:** Doum palm

**Local:**

**Scientific name:** *Hyphaene thebaica*

**Plant family:** ARECACEAE

**Description:** A tall, branched palm that grows 15-20 m tall. There are usually 4 crowns, but occasionally 8 or 16. Each branch gives rise to a crown. Each crown has 8-20 fan shaped leaves. The stem is smooth but it has the scars of the old leaf bases. The leaves are broad (130-180 cm) and with 20-40 long, thin segments. The leaves are grey-green. The leaf stalk is 90-140 cm long and spiny with curved hooks along the edges. The male flower is 1.3 m long and the female flower is a similar length, but is densely furry in the axils at the base of the fruit. The fruit is a dry fruit with a stone inside. They are about 7.5 x 5 cm and usually an irregular shape. The shiny outer layer encloses a middle layer which smells of gingerbread. This surrounds the hard stone which contains the seed kernel. The seeds are very hard and with a white kernel and a hollow centre. About 20-50 seeds weigh one kilogram. The mealy flesh of the fruit is edible.



**Distribution:** A tropical plant that grows in warm temperate, subtropical and tropical regions. It will not tolerate frost. It grows in coastal regions and can grow in arid places. It is found in the drier parts of West Africa and in the Sahel. It often grows on soil left by river floods. It forms dense stands in hot dry valleys. They grow where the temperature is 20-40°C and up to 600 m altitude. They are common in areas with a rainfall between 150 mm and 600 mm per year. A soil pH of 6.5-7.6 is preferred. They can grow on saline soils and suit hardiness zones 10-12. They are pollinated by wind.

**Use:** The spongy middle layer of the fruit is used for flour. The kernel is eaten. The seed kernels are sprouted in soil then eaten. The shoot from the developing seed before leaves emerge can be eaten. The young fruit are boiled and eaten. The palm cabbage is edible. The sap from the growing point can be drunk fresh or made into wine. The trunk contains a kind of sago starch which is edible.

**Cultivation:** Plants are grown from seed. Sometimes suckers are used. Seed should either be sown in a deep container or direct in the field. The seeds take a long time to germinate.

**Production:** The first harvest occurs after 6-8 years. Fruit ripen in 8-12 months. A tree lasts for about 60 years. Tapping the tree for sap kills the palm.

**Food Value:** Per 100 g edible portion

| Edible part  | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|--------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| nuts (dry)   | 6.0        | 1651      | 3.9       | -           | -           | -       | -       |
| fruit starch | 10.7       | 1237      | 2.6       | -           | -           | -       | -       |

## Nuts, seeds, herbs and other foods

**English:** African wild mango

**Local:**

**Scientific name:** *Irvingia gabonensis*

**Plant family:** IRVINGIACEAE

**Description:** A tree that grows to 40 m high. The trunk grows to 1 m across. It has narrow buttresses. The bark is light grey and smooth. The leaves are simple and alternate, and are 3.5-16 cm long and 2-8 cm wide. The yellowish-green scented flowers are small and occur among the leaves. The fruit are 10-13 cm long and 3-4 cm wide. The fruit has a thick covering and one seed.

**Distribution:** A tropical plant that grows in the humid forest zone in central Africa. It grows below 1000 m altitude. It grows in areas with a rainfall of 1500-3000 mm per year and temperatures of 25-32°C.



**Use:** The seed provides oil used in cooking. It is used to make Gabon chocolate or Dika bread. The kernels are ground and eaten in dishes of mixed vegetables. The kernels are extracted from the stones then roasted. They are then pounded and poured into a mould. This cheese is then scraped and added to boiling meat or vegetables. It is like a relish, especially for plantain bananas. The pulp of the fruit is eaten fresh.

**Cultivation:** Plants are grown from seed that germinate in about 14 days. It can be grown from stem cuttings under mist. Plants can also be budded.

**Production:** Young trees are slow growing. Fruit are usually harvested from the ground.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| nut         | 4.0        | 2918      | 8.5       | -           | -           | 3.4     | -       |
| fruit       | 81.4       | 255       | 0.9       | -           | -           | 3.4     | -       |

Image sourced from:

<https://www.researchgate.net/profile/Ebimieowei-Etebu/publication/270721748/figure/fig1/AS:295085292965893@1447365224580/Figure-1-Unripe-Irvingia-fruits-on-the-day-of-harvest.png>

## Nuts, seeds, herbs and other foods

**English:** Coastal almond

**Local:**

**Scientific name:** *Terminalia catappa*

**Plant family:** COMBRETACEAE

**Description:** A large tree, up to 25-40 m tall. It loses its leaves during the year. The trunk can be straight or twisted. There can be buttresses up to 3 m tall. The branches lie horizontally and come out in layers. The leaves are long, smooth and shiny, with an abrupt point at the tip and a rounded base. Leaves tend to be near the ends of branches. Leaves can be 17-29 cm long and 10-15 cm wide. Young leaves have soft hairs. The leaves turn red and fall off twice a year. Flowers are greenish-white and in a spike at the end of the branches. The lower flowers on a spike are female, and the others are male. The fruit is about 6 cm long by 3-4 cm wide, thick and flattened, with a flange around the edge. The fruit are green and turn red when ripe. The pulp is edible.



**Distribution:** It grows on beaches in almost all tropical countries in the world, including Solomon Islands. It is a tropical plant, and sometimes cultivated as a shade tree. The tree is common in lowland areas particularly on sandy or rocky beaches. Seeds are spread by bats and sea water, as well as being planted by people. It is common along streets in coastal towns. It will grow from sea level up to about 800 m altitude. Plants are frost-susceptible. It can tolerate drought. It suits hardiness zones 11-12.

**Use:** The kernel of the fruit is eaten raw. An edible oil can also be extracted.

**Cultivation:** Plants can be grown from seed. Seeds can be stored dry for a year or more. Seeds germinate freely and most seeds grow. Insects can badly damage the leaves of young seedlings.

**Production:** It is fast growing. Nut production is seasonal.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| nut (fresh) | 31         | 1810      | 15.9      | -           | 4           | 4.6     | 4.9     |
| nut (dry)   | 4.2        | 2987      | 20.0      | -           | 2           | 6.3     | 8.8     |

## Nuts, seeds, herbs and other foods

**English:** African breadfruit

**Local:**

**Scientific name:** *Treculia africana*

**Plant family:** MORACEAE

**Description:** An evergreen tree. It grows to 15-30 m tall. It can grow up to 50 m tall. It has a dense spreading crown. The trunk is fluted. The bark is dark grey and smooth. It is thick and produces a white latex when cut. This later turns rusty red. The leaves are simple and alternate. They are very large. Leaves can be 30 cm by 14 cm or larger. They are dark green and smooth above but paler and slightly hairy underneath. The leaves are tough. They have 10-18 pairs of clear veins. The leaf stalk is 1.5 cm long and the leaf tip is pointed. Young leaves are red or yellow. The flower heads are rounded and a yellow-brown. They are 2.5-10 cm across. Male and female flowers are usually separate. Flowers can grow in the axils of leaves or on older wood down to the trunk. The fruit is a compound fruit. It is rounded and very large. It can be 30-45 cm across. It grows on the trunk and main branches. Inside there are many orange seeds about 1 cm across. They are in a spongy pulp. The outer fruit surface is covered with pointy growths.



**Distribution:** A tropical plant. It suits hot, tropical lowland climates. It grows in forests near rivers. It can grow in swampy areas. It grows from sea level up to 1500 m in Uganda or 1200 m in Tanzania.

**Use:** The seeds can be dried, fried and eaten. They are also boiled, roasted or ground into flour. The flour is used in soups and nut milk. An edible oil can be extracted from the seeds.

**Cultivation:** Plants are grown from seed. Seed can be planted in pots then transplanted or they can be sown direct. There are about 5,000 seeds per kg. Seeds will only store for a few weeks but seed treatment is not needed before sowing.

**Production:** The tree is fairly fast growing. A fruit can weigh 12 kg.

**Food Value:** Per 100 g edible portion

| Edible part | Moisture % | Energy kJ | Protein g | proVit A µg | proVit C mg | Iron mg | Zinc mg |
|-------------|------------|-----------|-----------|-------------|-------------|---------|---------|
| seed (dry)  | 9.2        | 1555      | 12.6      | -           | -           | 320     | -       |

## Nutritional values of food plants by plant Family

| Plant Family   | Scientific name                             | Common name             | Edible part          | Moisture % | Energy kJ | Protein g | Vit A µg | Vit C mg | Iron mg | Zinc mg | Page |
|----------------|---|-------------------------|----------------------|------------|-----------|-----------|----------|----------|---------|---------|------|
| AMARANTHACEAE  | <i>Alternanthera sessilis</i>               | Lotus-seed herb         | shoot                | -          | -         | 5.0       | 577      | -        | 16.7    | -       | 32   |
| AMARANTHACEAE  | <i>Amaranthus cruentus</i>                  | Purple amaranth         | leaf                 | 84.0       | 176       | 4.6       | -        | 64       | 8.9     | -       | 34   |
| ANACARDIACEAE  | <i>Mangifera indica</i>                     | Mango                   | fruit                | 83.0       | 253       | 0.5       | 54       | 30       | 0.5     | 0.04    | 37   |
| ANNONACEAE     | <i>Annona muricata</i>                      | Soursop                 | fruit                | 82.4       | 294       | 0.88      |          | 16       | 0.3     | 0.1     | 39   |
| ARACEAE        | <i>Colocasia esculenta</i>                  | Taro                    | root                 | 66.8       | 1231      | 1.96      | 3        | 5        | 0.68    | 3.2     | 13   |
| ARECACEAE      | <i>Hyphaene thebaica</i>                    | Doum palm               | fruit starch         | 10.7       | 1237      | 2.6       | -        | -        | -       | -       | 56   |
| BASELLACEAE    | <i>Basella alba</i>                         | Indian spinach          | leaf                 | 85.0       | 202       | 5.0       | 56       | 100      | 4.0     | -       | 35   |
| BOMBACACEAE    | <i>Adansonia digitata</i>                   | Boabab                  | fruit                | 16.0       | 1212      | 2.2       | -        | 360      | 7.4     | 6.7     | 36   |
| CARICACEAE     | <i>Carica papaya</i>                        | Pawpaw                  | fruit                | 88.0       | 163       | 0.5       | 290      | 54       | 0.4     | 0.18    | 41   |
| CLEOMACEAE     | <i>Cleome viscosa</i>                       | Sticky cleome           | leaf                 | 80.4       | -         | 5.6       | -        | -        | 24      | -       | 30   |
| COMBRETACEAE   | <i>Terminalia catappa</i>                   | Coastal almond          | nut (fresh)          | 31         | 1810      | 15.9      | -        | 4        | 4.6     | 4.9     | 58   |
| CONVOLVULACEAE | <i>Ipomoea batatas</i>                      | Sweet potato            | tuber (baked)        | 72.9       | 431       | 1.7       | 961      | 24.6     | 0.5     | 0.3     | 19   |
| CUCURBITACEAE  | <i>Citrullus lanatus</i>                    | Watermelon              | fruit                | 94.0       | 92        | 0.4       | 20       | 5        | 0.3     | 0.1     | 43   |
| CUCURBITACEAE  | <i>Cucurbita pepo</i>                       | Marrow                  | fruit (mature)       | 92.0       | 105       | 1.6       | 17       | 16       | 2.4     | -       | 48   |
| CUCURBITACEAE  | <i>Cucurbita maxima</i>                     | Pumpkin                 | leaf                 | 88.0       | 160       | 4.9       | 260      | 28       | 2.5     | 0.9     | 51   |
| CUCURBITACEAE  | <i>Momordica charantia</i>                  | Bitter cucumber         | pod (boiled)         | 94.0       | 79        | 0.8       | 11       | 33       | 0.4     | 0.8     | 52   |
| CUCURBITACEAE  | <i>Cucumis metuliferus</i>                  | African horned cucumber | fruit                | 91.0       | 136       | 1.1       | -        | 18.6     | 0.5     | 0.3     | 53   |
| CUCURBITACEAE  | <i>Trichosanthes cucurmerina</i>            | Snake gourd             | fruit                | 95         | 76        | 0.9       | -        | 6        | 1.0     | -       | 49   |
| CYPERACEAE     | <i>Eleocharis dulcis</i>                    | Chinese water chestnut  | corm                 | 50.8       | 635       | 3.7       | -        | 52       | 15      | 1.9     | 15   |
| DIOSCOREACEAE  | <i>Dioscorea bulbifera</i>                  | Potato yam              | tuber                | 70.8       | 357       | 2.7       | -        | 78       | 3.1     | 0.4     | 18   |
| EUPHORBIACEAE  | <i>Manihot esculenta</i>                    | Cassava                 | tuber                | 62.8       | 625       | 1.4       | 30       | 15       | 0.23    | 0.48    | 16   |
| FABACEAE       | <i>Acacia farnesiana</i>                    | Sweet acacia            | seed (dry)           | 8.1        | 1522      | 36.6      | -        | -        | 6.0     | 0.6     | 21   |
| FABACEAE       | <i>Glycine max</i>                          | Soybean                 | seed                 | 9.0        | 1701      | 33.7      | 55       | -        | 6.1     | -       | 22   |
| FABACEAE       | <i>Sesbania sesban</i>                      | Egyptian sesban         | seed                 | 9.2        | 1446      | 32.0      | -        | -        | -       | -       | 23   |
| FABACEAE       | <i>Vigna unguiculata subsp. unguiculata</i> | Cowpea                  | seed (young, boiled) | 75.5       | 406       | 3.2       | 79       | 2.2      | 1.1     | 1.0     | 24   |
| FABACEAE       | <i>Lablab purpureus</i>                     | Lablab bean             | seed (young)         | 86.9       | 209       | 3.0       | 14       | 5.1      | 0.8     | 0.4     | 25   |
| FABACEAE       | <i>Canavalia ensiformis</i>                 | Jack bean               | pod (fresh)          | 88.0       | 155       | 2.4       | -        | -        | -       | -       | 26   |
| FABACEAE       | <i>Vigna umbellata</i>                      | Rice bean               | seed                 | 13.0       | 1373      | 20.9      | -        | -        | 10.9    | -       | 27   |
| FABACEAE       | <i>Cajanus cajan</i>                        | Pigeon pea              | seed (young, boiled) | 71.8       | 464       | 6.0       | 2        | 28.1     | 1.6     | 0.8     | 28   |
| FABACEAE       | <i>Phaseolus vulgaris</i>                   | Common bean             | pod                  | 88.0       | 151       | 2.5       | 750      | 27       | 1.4     | 0.2     | 50   |
| FABACEAE       | <i>Arachis hypogea</i>                      | Peanut                  | seed (dry)           | 4.5        | 2364      | 24.3      | -        | -        | 2.0     | 3.0     | 55   |
| IRVINGIACEAE   | <i>Irvingia gabonensis</i>                  | African wild mango      | nut                  | 4.0        | 2918      | 8.5       | -        | -        | 3.4     | -       | 57   |
| LAURACEAE      | <i>Persea americana</i>                     | Avocado                 | fruit                | 74.4       | 805       | 1.8       | 480      | 11       | 0.7     | 0.4     | 45   |
| MALVACEAE      | <i>Sida cordifolia</i>                      | Goat's horns            | leaf                 | 6.6        | 1296      | 24.2      | -        | -        | 79.8    | -       | 29   |
| MALVACEAE      | <i>Hibiscus cannabinus</i>                  | Vegetable kenaf         | leaf                 | 79.0       | 280       | 5.5       | 34       | -        | 12.1    | -       | 33   |
| MALVACEAE      | <i>Corchorus olitorius</i>                  | Jute                    | leaf (cooked)        | 87.2       | 155       | 3.4       | 156      | 33.0     | 3.1     | 0.8     | 54   |
| MORACEAE       | <i>Treculia africana</i>                    | African breadfruit      | seed (dry)           | 9.2        | 1555      | 12.6      | -        | -        | 320     | -       | 59   |
| MORINGACEAE    | <i>Moringa oleifera</i>                     | Horseradish tree        | leaf (boiled)        | 87         | 189       | 4.7       | 40       | 31.0     | 2.0     | 0.2     | 47   |
| POACEAE        | <i>Oryza glaberrima</i>                     | Floating rice           | seed                 | 11.3       | 1538      | 7.4       | -        | -        | 3.4     | -       | 11   |
| POACEAE        | <i>Sorghum bicolor</i>                      | Sorghum                 | seed                 | -          | 1459      | 11.1      | -        | -        | -       | -       | 12   |
| SAPINDACEAE    | <i>Nephelium lappaceum</i>                  | Rambutan                | fruit (raw)          | 82.0       | 271       | 1.0       | 0        | 53       | 1.9     | 0.1     | 44   |

| Plant Family | Scientific name                      | Common name | Edible part | Moisture % | Energy kJ | Protein g | Vit A µg | Vit C mg | Iron mg | Zinc mg | Page |
|--------------|--------------------------------------|-------------|-------------|------------|-----------|-----------|----------|----------|---------|---------|------|
| SAPOTACEAE   | <i>Englerophytum magalismontanum</i> | Milk plum   | fruit       | 74.3       | 404       | 0.9       | -        | 13.1     | 0.3     |         | 40   |
| ULMACEAE     | <i>Celtis integrifolia</i>           | Nettle tree | leaf (dry)  | 10.8       | 1058      | 8.0       | -        | -        | 19.7    | -       | 31   |





FOOD PLANT SOLUTIONS  
ROTARY ACTION GROUP  
Solutions to Malnutrition and Food Security

[www.foodplantsolutions.org](http://www.foodplantsolutions.org)