

Potentially Important Food Plants of Malawi



FOOD PLANT SOLUTIONS
ROTARY ACTION GROUP
Solutions to Malnutrition and Food Security

A project of the Rotary Club of
Devonport North and District 9830

www.foodplantsolutions.org

Potentially Important Food Plants of Malawi

Dedication

This book is dedicated to the 3 billion hard working farmers and families around the world who cultivate these, and other, food plants for their own subsistence, and who help conserve them in their rich diversity for other people to enjoy.

Preface

This guide is based on information from the Food Plants International (FPI) database developed by Tasmanian agricultural scientist Bruce French. The source material and guidance for the preparation of the book has been made possible through the support of Food Plants International, the Rotary Clubs of District 9830, particularly the Rotary Club of Devonport North who founded Food Plant Solutions, (previously the Learn2Grow project), and many volunteers who have assisted in various ways.

The selection of plants included in this guide has been developed by Lyndie Hyte working in a voluntary capacity using the selection criteria developed by Food Plant Solutions. These selection criteria focus on the local plants from each of the main food groups with the highest levels of nutrients important to human nutrition and alleviation of malnutrition. It is intended as a **Draft Guide only** to indicate some important food plants that serve as examples for this purpose. Other important nutritious plants may be equally useful, and it is recommended that the FPI database be used to source information on the full range of plants known to occur in Malawi. This guide has been developed with the best intention to create interest and improve understanding of the important local food plants of Malawi, and on the understanding that it will be further edited and augmented by local specialists with appropriate knowledge and understanding of local food plants.

Food Plant Solutions was initiated by the Rotary Club of Devonport North to assist in creating awareness of the edible plant database developed by Food Plants International, and its potential in addressing malnutrition and food security in any country of the world. In June 2007, Food Plant Solutions was established as a project of Rotary District 9830, the Rotary Club of Devonport North and Food Plants International. The primary objective of the project is to increase awareness and understanding of the vast food resource that exists in the form of local plants, well adapted to the prevailing conditions where they naturally occur, and how this resource may be used to address hunger, malnutrition and food security. For more information, visit the website www.foodplantsolutions.org. More detailed or specific information on plants, including references to material by other authors, is available on DVD on request.

Disclaimer: This Field Guide has been produced using information from the “Edible Plants of the World” database compiled by Bruce French of Food Plants International. Although great care has been taken by Food Plants International and Food Plant Solutions, neither organisation, or the people involved in the compilation of the database or this Field Guide:

- makes any expressed or implied representation as to the accuracy of the information contained in the database or the Field Guide, and cannot be held legally responsible or accept liability for any errors or omissions
- can be held responsible for claims arising from the mistaken identity of plants or their inappropriate use
- assume responsibility for sickness, death or other harmful effects resulting from eating or using any plant described in the database or this Field Guide

Always be sure you have the correct plant, and undertake proper preparation methods, by consulting with specialist scientists or local users of the plant. The Food Plants International database, from which the information in this Field Guide is drawn, is a work in progress and is regularly being amended and updated.

Contents

INTRODUCTION	1
STARCHY STAPLES.....	11
LEGUMES.....	23
LEAFY GREENS	31
FRUIT.....	39
VEGETABLES	49
NUTS, SEEDS, HERBS AND OTHER FOODS	58
NUTRITIONAL VALUES OF FOOD PLANTS BY PLANT FAMILY	66

Introduction

Potentially Important Food Plants of Malawi has been produced to provide information on approximately 40 edible plants that are known to grow in Malawi. These plants come from all the major food groups and have been chosen because of their high nutritional value. Many of the plants in this book may be neglected and under-utilised plants. This means they may not be well known. However, because they are high in many beneficial nutrients, and they are already adapted to the environment, and therefore likely to require minimal inputs, they could be important food plants that are likely to be superior to imported foods and plants. Commercially grown plants may also be included in the book, but only if they are significant foods for household consumption. It is hoped people will become confident and informed about how to grow and use these plants as many local food plants provide very good quality food.

Growing food

Growing food to feed a family is, without doubt, one of the most important things anyone can do. The more interest you take in your garden and the more you learn about plants and how to grow them well, the more interesting and fun food gardening becomes.

A country with very special plants

The local food plants of most countries have not been promoted and highlighted in the way they deserve. Visiting a local food market will quickly show what a rich variety of food plants can be grown in this country. Good information about these plants is often still in the minds and experience of local farmers, and has not been written down in books. This can make it hard for the next generation of young people to find out how to grow them.

In many countries, some of the traditional food plants are only harvested from the wild and others are only known in small areas. Others have hundreds of varieties and are the main food for people in different regions. Information on all these plants, their food value and the pest and diseases that damage them is available in the Food Plants International database.

Getting to know plants

People who spend time in gardens and with their food plants get to know them very well. It is a good idea to learn from someone who grows plants well. Each plant grows best in certain conditions and there are often special techniques in getting it to grow well. For example, sweet potato will not form tubers if the soil is too wet, but it may still grow lots of green leaves. Taro will grow in light shade, but sweet potato will not. Ginger can grow in fairly heavy shade. Pruning the tips of betel leaf or pepper vines will cause more side branches to grow and therefore, produce more fruit. Stored yam tubers need special treatment if you want them to put out shoots early. There are lots of unique things about every plant and learning about these helps a good gardener produce more food.


Naming of plants

Many food plants have local names, as well as a common English name. Every type of plant also has its own scientific name. Although the scientific name might not be widely recognised, this is the link by which people in different countries and with different languages can recognise the same plant. We know that many plants are grown in many different countries, but relying on local or common names, we might not recognise the same plant grown in different places. By using scientific names to accurately identify plants, we can get useful information from people in other countries. Wherever possible, plants in this book are named by their common English name and their scientific name.

Local food plants are often very good

People sometimes think that local food plants are not very special and that any food plant that is new or comes from another country must be a lot better. This is often not true. Many of the newer or introduced food plants, such as the round or ballhead cabbages, have very little food value. Many traditional tropical green, leafy vegetables and ferns have 10 times or more food value as ballhead cabbage or lettuce. It is important to find out more information about the food value of different foods if we want to eat well. Citrus fruit, such as lemons and oranges, are often grown for vitamin C that helps keep people healthy. These fruits do not grow well in the tropics-the common guava fruit has three times as much vitamin C and is loved by children. This is just one example that there are often much better choices of local foods with higher levels of important nutrients.

Our bodies need a variety of food plants to enable us to grow, stay healthy and have enough energy to work. Different foods are needed to provide energy, protein, vitamins and minerals. The following diagram highlights the iron content value of some traditional edible, tropical plant leaves, compared with cabbage. Iron is a nutrient that is very important for our bodies and especially our blood. People who are short of iron become anaemic and lack energy.



The relative iron content of some edible leaves

A healthy balanced diet

Good nutrition, or eating a healthy balanced diet, is really very simple. If people eat a wide range of food plants, their bodies will normally get a balanced amount of all the different nutrients they require. If a nutrient is lacking in one food plant, then they are likely to get it from another plant if they are eating a range of food plants. For this reason, everybody should eat a range of different food plants every day. The food group that is especially important for young people is the dark green leaves. Everyone should eat a good serving of dark green leaves every day. They have many vitamins and minerals, as well as protein. There are many spices or flavouring plants that can improve the taste of foods, but taste should be considered separately from food value.

Learning to cook well

Even though some nutrients in food can lose some of their value during cooking, it is normally much safer to cook all food plants, at least for a short time. Bacteria, which cause diarrhoea, can occur in gardens and on food plants. These are killed during cooking. Many plants in the tropics develop cyanide, a chemical that makes them bitter and poisonous. This happens often with cassava (tapioca, manioc) and beans, but can also occur in many other plants. Boiling the food for two

minutes normally destroys cyanide and makes the food safe to eat. Some of the nutrients our bodies need (such as vitamin A for good eyesight) only become available when food is cooked in oil.

Learning to grow “wild” food plants

Many plants grow wild in the bush and are not cultivated by people. We can normally find someone who has taken an interest in them and has learned to grow them. This may be people from a different language group. It may be that in their area they have found better types than the ones that simply grow wild.

Saving better types of plants

If we simply allow plants to grow from seed, the improvements that have been made in finding sweeter or better types may get lost. Some fruit trees are like this and the fruit produced may not be sweet at all. It is often necessary to take cuttings from a tree to be sure the new plant is exactly the same as the old one. If the plants won't easily grow from cuttings simply by sticking a piece of the branch in the ground, there are other ways of helping these plants to form roots and start to grow. One good way is to make a small cut in the bark of a young branch and then wrap soil around the cut and cover it with plastic. With plants like guava, new roots will start to grow from this cut and grow into the soil wrapped around the branch. It can then be cut off and planted. This is called air-layering. A similar method is used with the roots of breadfruit. A shallow root is uncovered and a small cut made from which a new sucker will start to grow. This can be cut off and replanted.

Growing from cuttings and suckers

Many food plants are grown from cuttings and suckers. This is very important, as it allows all the different kinds of yams, taros, bananas, sweet potato and sugarcane to be continually grown and ensures the varieties are preserved. Each plant has its own special propagation method. It is important to use healthy planting material, as diseases can be spread in planting material.

Saving seed

Some food plants are grown from seed. Sometimes this is very easy as the seeds are large, store well, grow easily and grow the same as the original plant. It is more difficult with other plants. Many large fleshy seeds, such as breadfruit, need to be planted while still fresh as they do not store easily. Other seeds do not “breed true” or do not grow into new plants that are the same as the original plants. For example, the fruit may not be as large or sweet or have the same colour or taste. With many of these plants, it may be necessary to find ways of growing them from cuttings or other methods such as grafting. Some plants “inbreed” and get smaller or poorer. This happens when a plant self-pollinates or receives pollen from a close relative. Corn grown in small plots normally does this and the plants grown from seed grown in this situation get smaller and smaller each year. The seed needs to be saved from several different plants with different history and then mixed together before sowing. All the seeds on one cob are related and will inbreed. Some seeds develop a hard seed coat and need to be scratched, soaked in water, or even put into hot water, before they will start to grow. Saving local seeds is often a good idea as they are already adapted to local conditions. For example, seed saved from pumpkins grown locally will produce plants with less pest and disease damage than those grown from imported seed. *If you can't get seeds or planting material from local gardens – it is probably not a suitable local plant!*

Growing a garden of mixed plants

In nature, one variety of one plant never grows alone. There are always lots of different plants of different kinds and sizes, all growing together. Anyone who has ever walked into a tropical jungle will know this very well. The reason people all over the world want to save the rainforest is because it has so many different kinds of plants all growing together. Growing plants in a food garden in a

way similar to how they grow in nature, as a mixed group of plants, is very good agriculture. Mixing plants in a garden usually gives more reliable food production, as any disease from one plant will wash off in the rain onto a different plant, where it cannot survive. Small plants fill the gaps and reduce the need for weeding.

Different types of plants for food security

There is another reason for growing a range of food plants in a local garden or around a village. If something goes wrong, like extreme insect damage to plants, some disease occurring in the garden, or a poor growing season, some plants will be more damaged than others. With a variety of plants, there will still be some food to eat until the other plants recover and grow again. Also, a wide variety of plants will mean that different ones will be maturing at different times, which helps ensure a continuous supply of food. There are shrubs that can be planted as edible hedges around houses, and fruit and nut trees that need to be planted as a gift for your children, several years before they will be able to enjoy them. Some nuts can be stored and eaten when other foods are not available. Most yams will store well for a few months.

Looking after the soil

Gardeners in traditional tropical agriculture usually move their gardens often by shifting to a new piece of land. There are usually three reasons for this:

- In the tropical lowlands, weeds can become a very big problem. There are usually a lot fewer weeds in the first year or two after clearing and burning the land, but weeds increase in the following years.
- Some of the nutrients in the soil are used each year and the soil becomes poorer and plants do not grow as well. There are ways of reducing this loss of nutrients.
- Very small worms called nematodes build up in the soil after a few years and get into the roots, especially of annual vegetable plants, and stop their roots working properly. For example, root knot nematode will cause the roots of plants like tomatoes and beans to become twisted resulting in poor growth of the plant.

Building up the soil

When a new garden has been cleared, it has lots of leaf mulch and other old plant material. This provides plant nutrients for new plants to grow. There is a simple rule for growing plants and improving the soil—"If it has lived once, it can live again." Any old plant material can provide nutrients for new plants to grow, but it must be allowed to rot into mulch or compost for this to happen. If this plant material is burnt, some nutrients, especially phosphorus and potassium ("potash"), get left behind in the ashes for new plants to use, although it also allows these important nutrients to be lost by being washed away by rain. But with burning other important nutrients, such as nitrogen and sulphur, get lost in the smoke and disappear from the garden and soil. These last two plant nutrients are especially important for growing green leaves and when their levels are low, plants grow small or pale green. When nitrogen is lacking, the old leaves of the plant go pale and fall off early, and when sulphur is lacking, the young leaves go pale. Wherever possible, old plant material should be covered with some soil to allow it to rot down and not simply dry out or get burnt.

Poor soils where crops won't grow

When soils are very acid (or sour), plants cannot get the necessary nutrients. Natural chemicals in the soil that are toxic to plants when present at higher levels become soluble, get into plants, and stop them growing. Adding limestone to these soils can improve them. Using compost will not make them less acid, but will keep the plant nutrients in the soil in a more readily available form that plants can use.

Soil nutrients

Plants need 16 different kinds of plant food or nutrients in different amounts to grow properly. A plant that has already been growing will have these nutrients in them and probably even have them in a balanced amount. That is why composting old plant material is so important. Plants usually show some signs or symptoms if any of these nutrients is running out.

One of the most common and important nutrients for plant growth is nitrogen, which actually comes from the air, but gets into plants through the soil. When plants are short of nitrogen, their older leaves often become yellow or pale. When grass family plants, like sugarcane and corn, are short of nitrogen, the centre of the oldest (lowest) leaves starts to develop a dry or dead V-shape. The plant cannot find enough nitrogen in the soil so it gets it from an old leaf to grow a new leaf. This causes the old leaf to die, forming a characteristic V-shape in the centre of the leaf. The plant does not get any bigger as an old leaf dies each time a new leaf is produced. Village farmers often walk through grassland before they clear it for gardens, looking to see if the grass leaves are dry and dead, because they know gardens on this soil won't grow well. It is necessary to use compost or legumes (such as beans) to put nitrogen back into the soil. Growing plants from the bean family (legumes) is the most efficient way to increase the level of nitrogen in the soil.

Corn is a good plant for indicating which nutrients are running short in the soil. If the older leaves go dry along the edges, the soil is running out of potash. If leaves that are normally green develop a bluish colour, the soil is short of phosphorus. Generally, leafy crops need lots of nitrogen, and root crops need lots of potash.

Making compost

Compost is old plant material that has been allowed to rot down into a fine, sweet smelling mulch that is full of nutrients that can be put back on the soil to grow new plants. Making good compost is very simple. A simple heap of plant material can be made in the corner of a garden or near a house. The composting process is carried out by small bacteria that live in the soil and feed on decaying plants. They break down old plant material into compost. These bacteria are living, so they need air, water and food. A good compost heap must have air, so don't cover it with plastic or put it in a container. This makes a foul smelling compost, as different bacteria that don't need air turn it into an acid mixture that preserves it. Good compost must have moisture, so keep the heap damp, but not too wet. The compost bacteria like a balanced diet, which means that both green material and dried material is needed to balance the carbon and nitrogen in the compost pile. If the compost material gets too dry and brown, it will not break down, and if it gets too green, it will go slimy. Using a little bit of compost from an old heap will make sure the right bacteria are there to start the whole process off. As soon as the plant material is broken down to a fine mulch it can be put onto the garden. It is best if it is dug in, but if it is regularly put onto the surface of the garden, worms will mix it into the soil.

Pests


There are a large number of insects that enjoy sharing our food with us! We should not try to kill all these insects as they have an important role to play in keeping everything in nature in balance. What we need to do is to learn to manage these insects so we can all get some food to eat! Some insects are attracted to lights, and if the garden is near village lights some insects can cause a lot of damage. If large areas of one particular crop are planted, insects can breed more quickly and cause a lot of damage. As an example, insects called armyworms can breed up in large numbers on the shade trees of cacao and then move "like an army" into gardens. Some insects are large and breed slowly and can be picked off and removed. The large, green grubs with pointy tips that hide under taro leaves are best controlled by simply picking them off. Some insects, like taro beetles, can be a

serious problem, but the young curl grubs of this insect are tasty if you catch and cook them. Some insects do not like sunlight. The very small moth that damages banana fruit is like this. Simply pulling off the leafy bracts over the banana fruit reduces the damage, as this lets sunlight in and the insect flies away. The best rule for reducing pest damage is to grow healthy plants, as they suffer less damage.


Diseases

The living organisms that cause disease are much smaller than insects. These disease organisms can often only be seen with a microscope. There are three main kinds of disease organisms-fungi, bacteria and viruses. Fungi are like the mushrooms we eat, only very much smaller. They usually make distinct dry spots on leaves and other plant parts. Fungi have spores that often blow in the wind. Bacteria are often smaller and live in damp places. They usually make plants go soft and squashy, and they may cause a smell. Bacteria are mostly spread with rain and in water. Viruses are very, very small and usually make irregular stripes and patterns on leaves and other plant parts. Viruses usually spread in planting material or in the mouths of small sucking insects. One common fungus disease on sweet potato causes the leaves to become wrinkled and twisted. It usually gets worse in old gardens and where soils are running out of nutrients. It doesn't affect all kinds of sweet potato to the same extent. The answer is not to stop the disease, but to improve the soil. The general rule is that healthy plants that are growing well will suffer less damage from disease.


Food value charts for a selection of plants from Malawi




Protein helps the body repair cells and make new ones. Protein is also important for growth and development in children, teens, and pregnant women. Symptoms of protein deficiency include wasting and shrinkage of muscle tissue, and slow growth (in children).




Vitamin A is very important for eyesight and fighting disease, particularly in infants, young children and pregnant women. People who are short of Vitamin A have trouble seeing at night.



Vitamin C helps us avoid sickness, heal wounds, prevent infections and absorb iron from food. Severe vitamin C deficiency increases the risk of scurvy with symptoms such as inflammation of the gums, scaly skin, nosebleed and painful joints.



Iron is important because it helps red blood cells carry oxygen from the lungs to the rest of the body. Low levels of iron cause anaemia, which makes us feel fatigued. Iron is also important to maintain healthy cells, skin, hair and nails. Iron is more available when Vitamin C is also present.



Zinc is particularly important for the health of young children and teenagers, and to help recovery from illness. It is needed for the body's immune system to work properly. It plays a role in cell division, cell growth, wound healing, and the breakdown of carbohydrates. Zinc is also needed for the senses of smell and taste. Zinc deficiency is characterized by stunted growth, loss of appetite, and impaired immune function.

Note regarding plant selection: In compiling these field guides, we acknowledge that some staple foods and commercial crops which are grown widely in the target country may be omitted. Such foods are often in the starchy staple category (e.g. rice, corn). This does not mean that they are not useful, but merely reflects a desire for the Food Plant Solutions project to concentrate on plants that are less well known and/or underutilised.

Starchy staples

English: Taro

Local:

Scientific name: *Colocasia esculenta*

Plant family: ARACEAE

Description: This plant has large flat leaves on the end of upright leaf stalks. It grows up to 1 m high. The leaf stalk or petiole joins the leaf towards the centre of the leaf. The leaves are 20-50 cm long. Near the ground a thickened rounded corm is produced. Around this plant there is normally a ring of small plants called suckers. Many different varieties occur. If left to maturity, a lily type flower is produced in the centre of the plant. It has a spathe 15-30 cm long which is rolled inwards. The flowers are yellow and fused along the stalk. There are many named cultivated varieties. Taro comes in two basic forms. The Dasheen type *Colocasia esculenta* var. *esculenta* and *Colocasia esculenta* var. *antiquorum* or the Eddoe type. The basic difference is the adaptation of the Eddoe type to storage and survival in seasonally dry places, while the dasheen type needs to be maintained in a more or less continuously growing vegetative stage.



Distribution: It is a tropical plant. Taro grows from sea level up to about 2300 m altitude in the tropics. It grows well in humid places. It can stand damp soil and grow under light shade. It suits hardiness zones 9-12.

Use: The corms, petioles and leaves are all edible after cooking. The leaves are also dried and stored. Fresh leaves can be stored for 4-5 days. **Caution:** Some varieties burn the throat due to oxalate crystals.

Cultivation: Taro can be planted from cormels or from the top of the central corm. Other sections of the corm could also be used but this is not commonly done. Flowering of taro and seed production can lead to new cultivars. Flowering can be promoted by the use of gibberellic acid. The general growth pattern is for an increase in top growth, in terms of leaf number, leaf area and petiole length, to continue for about 6 months under tropical lowland conditions then for each of these to decrease and tuber storage to continue to increase. Corm weight increases significantly from 5-11 months. Starch content also increases with time but protein content declines over the corm development period.

Taro can be grown under flooded conditions but root rots develop if the water becomes stagnant. For flooded cultivation, the land is cleared, ploughed, cultivated and puddled. The aim is to get a field that is flat with embankments allowing the impounding of water. Planting is done into 2-5 cm of standing water.

For dryland taro, the soil is prepared by digging, unless a fresh bush fallow is used where the natural friability of the soil allows plants to be put into the undug soil in a small hole that is prepared. Plants are put into a hole 5-7 cm deep or deeper. Mulching to conserve moisture and reduce weed growth is beneficial. Setts from corms normally give higher yield than that from cormels. The greater leaf area and root production may be responsible for this. Setts of about 150 g are optimum.

The time of planting is primarily determined by the availability of moisture. Planting is done shortly after the rainfall has become regular, if seasonally distinct wet and dry occur. Higher rainfall, higher temperatures, and higher hours of sunlight, enhance production and determine seasonality of production.

Evapotranspiration for flooded taro averages about 4 mm per day, ranging from 1.5-7.2 mm, with a total of about 1200 mm for the crop. Intermittent moisture can result in irregular shaped corms. Flooding has been found to be more effective than sprinkler irrigation, or furrow irrigation. Increased suckering, giving greater leaf area, seems to be the reason for this.

Taro is sensitive to weed competition throughout most of its growth, but it is more critical during early growth up to 3-4 months. About 7-9 weedings are required, to keep the crop clean under tropical lowland conditions, where flooding is not used. Due to the decrease in height and leaf area towards the end of the growth cycle when starch accumulation in the corms is maximum, weed competition and weed control are again significant. Mechanical weeding needs to be shallow to avoid damaging the superficial taro roots. A range of herbicides have been recommended in various situations.

Taro produces the highest dry matter yield under full sunlight, but it can still grow under moderate shade. Under shaded conditions it grows more slowly and develops fewer cormels. They require good moisture conditions and have little tolerance for drought. Taro residue has an allelopathic factor which can reduce the germination and growth of other plants, for example, beans.

Taro tends to demand high fertility, and is responsive to additional NPK fertiliser. Higher doses of K increases starch content and higher doses of N increases protein content. Both N and K applications increase oxalic acid content of the tubers.

Spacing affects total yield, and marketable, harvestable yield, of corms. Close spacing increases the corm yield per area, and the shoot yield per area, but decreases the corm yield per plant, and the contribution of sucker corms, to the yield. Where spacings of 30 cm x 30 cm are used, giving about 110,000 plants per hectare, a very large amount of planting material is required, which reduces the net return per unit of planting material. A spacing of 60 cm x 60 cm is more common. Wider spacings of 90 cm x 90 cm reduces overall yield.

Production: Crops mature in 6-18 months. Yields of 5-15 tonnes per hectare are probably average.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
root	66.8	1231	1.96	3	5	0.68	3.2
leaf	85.0	210	5.0	57	90	0.62	0.7
leaf stalk	93.0	101	0.5	180	13	0.9	-
leaf (cooked)	92.2	100	2.7	424	35.5	1.2	0.2

Starchy staples

English: Sweet potato

Local:

Scientific name: *Ipomoea batatas*

Plant family: CONVOLVULACEAE

Description: This is a root crop which produces long creeping vines. The leaves are carried singly along the vine. Leaves can vary considerably from divided like fingers on a hand, to being entire and rounded or heart shaped. Purple trumpet shaped flowers grow at the end of the vine. Fattened tubers are produced under the ground. There are a large number of varieties which vary in leaf shape and colour, tuber shape, colour, texture and in several other ways.



Distribution: A tropical and subtropical plant. They grow from sea level up to about 2700 m altitude in the tropics. Plants can grow with a wide range of rainfall patterns and in different soils. Plants are killed by frost and can't stand water-logging. Plants grow well with temperatures between 21-26°C. It can grow with a pH between 5.2-6.8. Sweet potato are not tolerant to shading. It suits hardiness zones 9-12.

Use: Tubers are boiled or baked. They can be steamed, fried, mashed or dried. They can be fermented into alcoholic drinks. They can also be used in pies, cakes, puddings and candies and jams. They can be used in noodles. The chopped and dried tubers can be boiled with rice or ground into flour and mixed with wheat flour to make cakes or bread. The young leaves are edible.

Cultivation: Vine cuttings are used for planting. In grassland soils it is grown in mounds, ridges or other raised beds. In bush fallow, it is mostly planted in undug loose soils. It needs a sunny position. Tubers won't form if the ground is waterlogged when tubers start to develop. Sweet potato is grown by cuttings of the vine. About 33000 cuttings are required per hectare. These weigh about 500 kg. Vine lengths of about 30 cm are optimum. As long as the vine is adequately inserted in the soil, the length of vine inserted does not significantly affect yield. Fresh sweet potato seeds germinate relatively easily and lead to continuous production of new cultivars under tropical conditions. Excess nitrogen restricts storage root initiation and therefore excess leaves are produced without significant tuber yield. Dry matter percentage increases with increasing age of the crop. Higher dry matter tubers are normally preferred.

Sweet potato are not tolerant to shading. Under shaded conditions, both foliage growth and storage root production are decreased. Some cultivars can be selected for increased production under mild shade but not heavy shade. The survival of cuttings at planting is also reduced under shaded conditions. Under shaded conditions, plant become more climbing and with fewer, larger leaves. With increasing shade, fewer tubers are produced and these grow more slowly. Sweet potato tends to be responsive to potassium fertiliser. Cultivars are often selected for yield under low fertility conditions.

Under lowland conditions in the tropics sweet potato tubers undergo active tuber enlargement from 6-16 weeks. Weed control is essential especially during early stages of growth. The rate of ground coverage by foliage varies greatly with growing conditions and cultivar, but once ground coverage has occurred, weed control is less of a problem. Sweet potato tuber initiation is subject to

aeration in the soil. Either heavy clay soils, waterlogged conditions or other factors reducing aeration can result in poor tuber production. For this reason, sweet potatoes are often grown on mounded beds. In well drained or high organic matter soils, digging or mounding is not as essential. Leaf scab (*Elsinoe batatas*) can significantly reduce yield especially in sites where leaf production is low due to low soil fertility. To reduce sweet potato weevil damage, plants need to be hilled or have the tubers well covered with soil. Cracking soils can allow the weevil access to tubers.

Production: The time to maturity ranges from 5 months to 12 months depending on the variety planted and the altitude at which it is being grown. Yields range from 6-23 t/ha.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber (baked)	72.9	431	1.7	961	24.6	0.5	0.3
tuber (raw)	70.0	387	1.2	709	25	0.7	0.4
tuber (boiled)	72.0	363	1.1	787	15	0.6	0.3
leaf	86.3	168	3.9	105	58	2.9	-

Starchy staples

English: Cassava

Local:

Scientific name: *Manihot esculenta*

Plant family: EUPHORBIACEAE

Description: A plant which can re-grow year after year from the thickened roots. It has several stems. The stems are woody and have some branches. Plants grow up to 3 metres tall. Stalks have distinct scars where leaves have fallen. The leaves tend to be near the ends of branches. The leaves are divided like the fingers on a hand. The leaves have long leaf stalks. The leaves have 3-7 long lobes which can be 20 cm long. These are widest about 1/3 of the distance from the tip and taper towards the base. The colour varies. It produces several long tubers. These can be 50 cm long by 10 cm across. The flowers are on short stalks around a central stalk. They are produced near the ends of branches. The female flowers are near the base of the flower stalk and the male flowers higher up.



Distribution: A tropical plant. Plants grow from sea level up to about 1650 m. In Fiji they grow to 900 m. They can grow in poor soil and can survive drought. It is native to tropical America. It grows between 25°N and 25°S and needs a rainfall above 750 mm. It suits hardiness zones 10-12.

Use: The tubers are eaten after thorough cooking. They are boiled, roasted or made into flour. The starch is used in puddings, soups and dumplings. Young leaves are edible after cooking. They are also sometimes dried and stored. Seeds are also eaten. **Caution:** Bitter kinds of cassava contain poison but this is destroyed on heating. This kind of cassava should be cooked, sun dried, soaked and cooked again.

Cultivation: Cassava is planted from sections of the stalk. Sections about 15-20 cm long of the more mature woody stem are cut and stuck into the ground. They can be completely buried or put at almost any angle and it affects the growth little. Soon roots form and leaves start to sprout from the stalk. Cassava seeds need a soil temperature of 30°C for their germination. Flower and fruit production is more common under lower temperatures such as in highland or less equatorial conditions.

It is not necessary to dig a hole to plant cassava and on many soils where the soil is loose it can be planted without digging the soil first. Cassava does not suit waterlogged soils and preferably they should not be too shallow or stony.

Cassava can be planted at any time of the year but to get started it needs moisture so is often planted near the beginning of the wet season. The crop once established can survive for several months without rain. The ability to tolerate drought varies significantly with cultivar. During drought less and smaller leaves are produced and leaves die off more quickly but storage roots can be increased in the short term.

Because cassava can still grow satisfactorily in poorer soils it is often put last in a rotation after others crops have already been grown on the piece of land. Cassava is more responsive to nitrogen

and potassium than phosphorus under many field situations. Nitrogen can increase cyanide levels. Under very acid conditions with high soluble aluminium levels, cassava has been able to achieve and maintain top growth but with significantly reduced root yields. When drainage is good and soil moisture is adequate, cassava stalks can be planted at any orientation from horizontal to vertical, but in very sandy soils horizontal planting is best and in heavy clay soils vertical planting is best.

Because of the slow growth in early establishment stages, soil loss from erosion with heavy rains can be significant. To avoid this planting should be timed so that the maximum vegetative growth is occurring during the heaviest rains. A leaf area index between 2.5-3.5 is optimal for cassava yield. The critical period for weed control is the time from 2-8 weeks after planting. Cassava tuber bulking is delayed under shaded conditions. Yields are also reduced. In mixed cropping situations using crops which mature early, allowing the cassava time to recover, is one possible strategy. For optimum production shading should be avoided.

Cassava takes about 10-12 months to produce mature tubers in the lowlands tropics although some varieties produce a smaller yield earlier. Yields in the range of 20-45 t/ha have been recorded for 12-14 month crops. The plants can be left growing and the tubers stored in the soil for considerable time. Crops of 24 months duration occur. Once the tubers have been dug they do not keep for more than a few days. Pre-harvest pruning of plants increases the storage time of tubers after harvest.

Spacing and plant density varies with soil climatic conditions and variety. Plant densities from 10000 to 30000 plants per hectare are used. Plants from the higher density crops have been shown to have quick post-harvest deterioration. Mulching has given significant yield increases in some conditions. It also reduces the incidence and damage of some root boring insects.

Production: Plants can be harvested after 10 months in the lowlands. There are some faster growing varieties. Yields in the range of 20-45 t/ha have been recorded for 12-14 month crops.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber	62.8	625	1.4	30	15	0.23	0.48
leaf	82.0	382	7.1	57	275	7.6	-

Starchy staples

English: Country potato

Local:

Scientific name: *Plectranthus rotundifolius*

Plant family: LAMIACEAE

Description: Country Potato is a small annual herb. It grows 15-30 cm tall. Its succulent stems can lie along the ground or curve upwards. The leaves are thick and have a smell like mint. The flowers are small and pale violet. Dark brown tubers are produced in clusters at the base of the stem. The tubers can be up to 20 cm long and 2 cm wide.

Distribution: Country potato is a tropical plant. It grows well in high rainfall areas with low night time temperatures. It is grown as a monsoonal crop.



Use: The tubers of Country Potato are commonly boiled and eaten but they can also be eaten raw. They can be added to curries, baked, or fried into chips, steamed, added to soups or stews or mashed and fried. The leaves are edible and can be used as a flavouring. The ripe fruit can also be eaten.

Cultivation: Country Potato is grown from suckers from germinating tubers. The tubers are put in a nursery bed about 4 cm deep and 10 cm apart. Tubers germinate in 10-15 days once watered. These produce a cluster of sprouts which are then transplanted after about 3 months. The earth is heaped up around the plants to encourage tuber growth. Plant spacings of 20 cm apart in rows 90 cm apart are used.

Plants can also be grown from cuttings. It needs a well drained, sandy loam. It cannot stand water-logging. In heavier soils, plants should be grown on ridges. Plants cannot stand cold, frost or drought. The best soil pH for optimal growth is 6.5-7.0.

Production: Crops reach maturity after about 5-6 months. This may be longer in colder places. Yields of 8-15 tonnes per hectare are average. Tubers should be harvested as soon as they are mature to avoid decay.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber	76	393	1.4	-	1.0	6.0	-

Image sourced from: <http://d3d71ba2asa5oz.cloudfront.net/12001418/images/plectranthusrotundifolius1.jpg>

Starchy staples

English: Greater yam

Local:

Scientific name: *Dioscorea alata*

Plant family: DIOSCOREACEAE

Description: A long angular vine with square stems that twine to the right around support sticks. The stem does not have spines and is often coloured green or purple. The leaves are heart shaped and borne in pairs along the vine. The leaves vary in shape, size and colour with different varieties. Leaves can be 10-30 cm long by 5-20 cm wide. The leaf stalk is 6-12 cm long. The flowers occur in the axils (where the leaf joins the stem) of the upper



leaves. The male flowers are in small heads along branched stalks. These can be 25 cm long and green. The female flowers are in shorter spikes. Many cultivated varieties do not produce fertile seed. The fruit are 3-winged and 2.5 cm long by 3.5 cm wide. The seeds, when they occur, have wings right around them. One large, but often irregular-shaped, tuber occurs under the ground. There are many different varieties. The tubers can vary in shape, size, colour, texture and other ways. Some varieties produce bulbils, or small bulbs, along the vine.

Distribution: It grows in many tropical countries, growing from sea level up to about 1800 m in the tropics. Yams are most important in seasonally dry areas. They need a well-drained soil with reasonable fertility and are, therefore, often planted first in rotations. The maximum temperature is >30°C while the minimum is 20°C. The best temperature range is 25-30°C. Rainfall is often seasonal in yam areas and the maximum needs to be 14-20 weeks rain, with the best being 1,150 mm during the growing season. Yams can tolerate drought, but give best yields with high rainfall. The critical rain period is during the first 5 months. They cannot tolerate water-logging. Yams are influenced by the number of hours of sunlight. Short days (less than 10-11 hours of sunlight) favour tuber development. Yams suit hardiness zones 10-12.

Use: The tubers are boiled, baked or mumued (cooked in the ground).

Cultivation: For general food production, use top pieces of the tuber after they have sprouted, use a branched stick for supporting the vine, space plants about 1 m apart and choose a smooth round variety of yam. This makes harvesting easier, and peeling and food preparation quicker. Varieties that get less leaf spot disease and are less damaged by virus diseases give a more reliable yield. Tubers which are cut and stored in shady places until they form sprouts give improved yields over tubers that are left whole then cut into setts at planting. Because yam tubers have a period of dormancy, tubers do not normally commence regrowth for up to 5-6 months. This means they store, but cannot easily be used for out of season replanting. Dormancy, or inactivity, of the yam tubers can be broken using Calcium Carbide treatment for 5 hours, or by covering tubers with leaves of *Croton aromaticus* or *Averrhoa bilimbi*.

In some kinds, the bulbils that grow along the vine can be used for planting. By using staggered plantings of male and female plants, and then hand-pollinating the flowers, it is possible to get seeds to develop and these can be used to establish new plants. It is common practice in many areas to plant the yam piece upside down. The probable reason for this is to give the shoot and roots time to develop and get established away from the sun and wind so that the plant does not dry out. People in yam areas have their varieties classified as to whether they are planted at the top or the

bottom of the hole, and whether the shoot is pointed up or downwards. A planting depth of 15 cm is best. Normally top pieces give a higher yield than middle pieces of the tubers and these are better than bottom pieces. Top pieces of the yam tuber give earlier and more reliable shoots and the yams mature earlier. These top pieces are also the less attractive part of the tuber for eating, so they are preferred for planting. The larger the sett, the earlier it develops shoots and the larger the yield. Putting plants more closely gives smaller yams, but more total food. Closer spacing is normally used on lighter soils.

Yams should also have sticks to climb up. It is best to have a stick that is twisted or branched because the vine can slip down a very straight stick. Normally, a stick 2 m tall is sufficient. It needs to be a strong stick, firmly fixed in the ground. Yam varieties have varying types of vine growth. This affects where the stick needs to be placed. The fat, irregular yams can have the sticks near the mound, as a thick clump of vines and leaves soon develops. But, if a the stick is put beside the mound of one of the long ceremonial yams, the vine will often reach the top of the stick before it has produced more than a couple of leaves and will then fall back down to produce its leaves on the ground. The stick for these varieties often needs to be put at some distance from the yam hole. The tip can be picked off the vine if branching is wanted earlier.

Light influences the growth of the tubers. If the tubers have light on them often, due to cracks in the soil on hillsides, tubers are smaller. Compact soil or stones means the tubers may be exposed to sunlight. This needs to be avoided as it reduces yield. Yams must have plenty of air in the soil, so they will not normally grow on heavy clay soils or in areas with a lot of soil moisture. The soil can be improved for yam growing by putting leaves and other plant material in the planting hole, by making a mound above the hole, or by planting on a hillside. In some very loose sandy soils, yams can just be planted in flat, unmounded soils without digging a special yam hole, but these situations are not common.

Production: In most places, the yam growth and time to maturity is linked to seasonal rainfall patterns. They are mostly planted just before the first rains, where a 8-10 month rainy season exists. They give better yields in 6-8 month rainy season areas, where they are planted 3 months before the rains. Earlier planting requires larger sett size to withstand drying out. In drier grassland areas, mulching the mounds at planting means fewer plants die and more food is produced. The time to maturity ranges from 5 months on the coast, to 9-10 months at higher altitudes. Yams will store well for over 6 months in a dry, dark, well-ventilated shed. Greater yam is an important root crop of the seasonally dry, hot humid, tropics.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber	76.6	323	2.0	18	10	0.8	0.39

Starchy staples

English: Potato yam

Local:

Scientific name: *Dioscorea bulbifera*

Plant family: DIOSCOREACEAE

Description: A yam with a long smooth stemmed vine, round in cross section and without spines. The vine winds to the left, can climb into trees and grow to long lengths. The large leaves (14-30 cm across and slightly longer than wide) have pointed tips and round bases. About 7 veins arise from the tip of the leaf stalk. It produces often flattened bulbils (potatoes) in the leaf angles along the vine. They can be grey brown or purple. The smaller tuber underground is normally covered with roots. The flowers are large. The male flowers are in spikes up to 20 cm long. The female spikes are usually in pairs. The winged fruit are about 2.5 cm long by 1.5 cm across. The seeds have wings. The bulbils normally have few fibres through the tissue compared to some yam tubers. Many varieties have yellow flesh.



Distribution: An annual tropical plant. It will grow from the coast up to about 1700 m altitude in equatorial zones. It is common near the edge of grassland and forest at mid altitudes. Both wild and cultivated forms occur. It is common near secondary forest at low and medium altitudes.

Use: The cooked tubers aerial bulbils are eaten. Some kinds are bitter and inedible or at least require special processing and cooking. Some varieties are poisonous.

Cultivation: Either the vine bulbils or the underground tubers are planted. The long vines can be trained up trees. The bulbils need a set storage time before sprouting. The leaves die off for 1-4 months each year before re-sprouting from the tuber. Bulbils only grow shoots from one end unless the bulbil is cut into pieces. If larger bulbils are cut, the cut surfaces should be dried and healed in a shady place for 2-3 days before planting. Bulbils are planted 8-12 cm below ground at a spacing of about 100 cm by 100 cm. Nitrogen and potassium fertilisers give greater responses than phosphorus. Friable well drained soils are most suitable. Often little cultivation or mounding is done. A high level of organic matter improves yield. Strong staking is required with branched 2 m stakes, or with trees or living stakes. Vine and tuber growth can be extensive and heavy.

Production: Bulbils or aerial yams are produced as leaves begin to unfold, continue until plant maturity, and often fall. Harvesting can start 3 months after planting but immature tubers have less starch. Underground tubers are normally not harvested until leaf die back. Wounds and damage to the tubers normally heal naturally given dry aerated conditions. Some varieties have seasonally dormant aerial tubers which only grow after an extended storage period. Others germinate quickly.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
tuber	70.8	357	2.7	-	78	3.1	0.4
bulbil	79.4	326	1.4	-	-	2.0	-

Starchy staples

English: Sorghum

Local:

Scientific name: *Sorghum bicolor*

Plant family: POACEAE

Description: Sorghum is a millet grass. A mature sorghum plant resembles maize in its stature. Plants vary in height from 45 cm to 4 m. It is an annual grass with erect solid stems. The stems can be 3 cm across at the base. Prop roots occur at the base of the plant. There are numerous sorghum varieties. Some have one main stem while others produce multiple tillers. More tillers are produced when plants are widely spaced. The nodes on the stem are slightly thickened. Short types have up to 7 leaves while tall late varieties may have up to 24 leaves. The leaf blade can be 30-135 cm long. Leaves are bluish green and waxy. They have a prominent midrib. The large flower panicle can be 20-40 cm long. The flower occurs at the top of the plant. It can stick upright or bend over. The flower can be open or compact. Over 1000 cultivated varieties occur in China.



Distribution: Sorghum is a tropical plant. It suits the savannah zones in the tropics and can tolerate heat and drought. It can recover from drought even as a seedling. It can tolerate water-logging. It can be grown on heavy or light soils. Sorghum requires short day lengths to flower. Many kinds are adapted to specific day length and rainfall patterns. It suits hardiness zones 9-12.

Use: Sorghum seeds are eaten as a cereal. Flour can be made from the grain and then used for porridge or other dishes. It is used for dumplings, fried cakes and drinks. It cannot be used for bread as it contains no gluten. The stems of some kinds are sweet and can be chewed. The grains can be popped and eaten. The sprouted seeds can also be eaten.

Cultivation: Sorghum seeds will germinate soon after harvest. The seeds also store well if kept dry and protected from insects.

Production: Grain is ready for harvest 4-8 weeks after flowering.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	-	1459	11.1	-	-	-	-

Starchy staples

English: Finger millet

Local:

Scientific name: *Eleusine coracana*

Plant family: POACEAE

Description: An annual millet grass. It is robust and forms many tillers or young shoots from the base. It grows 40-120 cm tall. The stems are somewhat flattened and the leaves are narrow. The flower heads are made up of 2-7 finger like spikes, 1.5 cm across and 10-15 cm long. These in turn have about 70 smaller spikes. Each one of these smaller spikes has 4-7 seeds. The seeds are roughly rounded and 1-2 mm across. There are *coracana* and *africana* subsp.



Distribution: It is a very drought resistant tropical plant. For good yields, it needs good soil drainage and adequate moisture. It cannot stand water-logging. It is an important crop in areas where annual rainfall is 900-1250 mm. It especially suits areas with long hot summers. It needs a minimum temperature above 18°C and does best where temperatures are above 27°C. It grows from sea level to 2,400 m altitude in Africa. It is a short day length plant and does best where day length is 12 hours. It can grow in arid places.

Use: The seed are eaten either roasted or ground into flour. This is used for porridge and flat bread. Alcohol is brewed from the grain. The leaves are also edible.

Cultivation: It is grown from seed. Often plants are grown mixed with sorghum or maize. Good soil preparation is needed to reduce weed competition. Seed can be broadcast or drilled. Young plants need to be weeded and thinned. Seed viability drops to about 50 % after 2 years. Spacings of 5 cm apart in rows 30-33 cm apart, or 10-12 cm apart in rows 25 cm apart are recommended. About 25-35 kg of seed per hectare are needed if seed are broadcast. 5-10 kg per hectare are required if seed are drilled. Using fertiliser can dramatically increase yield. 125 kg per hectare of sulphate of ammonia when plants are 15 cm high is used in Uganda.

Production: It is self-pollinating and pollination occurs over 8-10 days. Millet seed stores very well and can be stored without damage for 10 years. Often it is stored on the head. Yields of about 450-900 kg of dried grain per hectare are usual. This can easily be increased to 1650 kg per hectare. Crops take 3-6 months until harvest.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	11.7	1594	6.2	-	-	5.3	-

Legumes

English: Soybean

Local:

Scientific name: *Glycine max*

Plant family: FABACEAE

Description: A small erect bean growing up to 60 cm tall. It grows each year from seed. Straggling kinds can occur. Stems, leaves and pods are softly hairy. The leaves have 3 leaflets. The leaflets have stalks. Flowers are small and white or blue. They occur in groups in the axils of leaves. The pods are broad, flat and hairy. Pods have 2-4 seeds. The seeds can be yellow to black.



Distribution: It is a temperate plant that suits lowland areas. It can be grown from sea level to 2000 m altitude. Many varieties will not flower in the tropics (short days). It needs fertile soil. The best soil acidity is pH 5.5-7.0. It is damaged by frost.

Use: The young pods and ripe seeds are eaten. They are used for flour. The dried seeds are boiled or baked and used in soups, stews and casseroles. The seeds are used for oil. Toasted seeds are eaten like a snack. Strongly roasted seeds are used for coffee. Soy flour is used for noodles, and confectionary. The beans are fermented and used in a range of foods. Sometimes the young leaves are eaten. The seeds are also used for sprouts and for making cooking oil and soya sauce etc. Because soybean contains a trypsin inhibitor they should be cooked and even the sprouts should be lightly cooked.

Cultivation: It is grown from seed. Seeds need to be inoculated with bacteria before planting. Plants need to be about 20 cm apart.

Production: Plants flower about 8 weeks after sowing and pods mature about 16 weeks after sowing. Often plants are pulled up and hung up before threshing out the seed.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	9.0	1701	33.7	55	-	6.1	-
seed (immature)	68.0	584	13.0	16	27	3.8	0.9
sprout	79.5	339	8.5	1.0	8.3	1.3	1.0

Legumes

English: African locust bean

Local:

Scientific name: *Parkia filicoidea*

Plant family: FABACEAE

Description: A deciduous tree that grows up to 35 m tall. It has a spreading flat crown. The trunk has small rounded buttresses. The grey to yellow-brown bark can be scaly or smooth, and becomes dark and cracked with age. The bark has an orange coloured resin. The leaves are feathery. A leaf is made up of 6-9 pairs of leaflets each divided into 16-24 pairs of smaller leaflets. These are about 2 cm long and 5-8 mm wide. The flowers are small and in bright red club shaped heads. These hang down on stalks 30 cm long. The flower heads are up to 8 cm long. The fruit are dark brown to purple pods which hang down in clusters. They are 30-60 cm long and 2 cm wide with their stalk. The pod is narrowed slightly between the seeds. The seeds are red-brown in a dry, mealy, edible, yellow pulp.

Distribution: A tropical and subtropical tree of lowland rainforests. It grows in Africa in forests near streams. It occurs in sub-humid and humid places with an annual rainfall of 950-1750 mm annually. It grows from 250-1370 m above sea level. It can grow in arid places.

Use: The pods and the pulp are eaten. The seeds are boiled and fermented then eaten. This has a strong smell but is removed by frying or roasting. The seeds can also be powdered and used for flavouring soups and rice dishes. The leaves are cooked as used as a vegetable.

Cultivation: Plants can be grown from seed. The pod is crushed and the seed removed from the pulp. The seed they should be boiled briefly, then allowed to cool and soaked for 12 hours, before sowing.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	7.0	1780	32.3	-	6	33.2	-
fruit	13.2	1263	3.4	-	-	3.6	-

Image accessed from:

http://www.westafricanplants.senckenberg.de/images/pictures/fabmimo_parkia_filicoidea_cbch_6118_4049_b8ed36.jpg



Legumes


English: Cowpea

Local:

Scientific name: *Vigna unguiculata*

Plant family: FABACEAE

Description: A creeping bean type plant with straight firm pods. There is a deep tap root and many branches occur from it in the surface of the soil. The root nodules are large and round. The leaves have 3 leaflets. The end leaflet can be 12-16 cm long. The side leaflets are asymmetrical. The stipules at the base of the leaf are large and with spurs at their base. Flowers occur often in pairs on the end of long flowering shoots. Only 2-4 flowers in each stalk produce pods. Flowers are white, yellow or blue. They are large and showy. The pods are about 15 cm long. The seeds are white except for a dark scar.



Distribution: It grows in tropical and subtropical climates. It grows from sea level to 1800 metres altitude in the tropics. Plants can stand high temperatures. Some kinds can tolerate drought. They are sensitive to cold and killed by frost. Plants germinate with a temperature between 11.5-15.5°C. The best growth occurs between 20-35°C. They can grow on a range of soils providing they are well drained. They are a short day plant. They do well in the semiarid tropics. It will not tolerate acid or alkaline soils. It grows in areas with an annual rainfall between 280-410 mm. It can grow in arid places.

Use: The young leaves, young pods and ripe seeds are all eaten. They can be steamed, boiled, stir-fried etc. The leaves can be dried and stored. The dried seeds are used in soups and stews. They are ground into flour or fermented. The seeds are also used for bean sprouts. Roasted seeds are used as a coffee substitute.

Cultivation: It is grown from seeds. Seeds remain viable for several years if carefully stored. A seeding rate of about 20 kg per ha is suitable and seed are sometimes broadcast then thinned.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	11.2	1189	23.5	-	1.5	6.4	-
seed (young, boiled)	75.5	406	3.2	79	2.2	1.1	1.0
leaf	88.4	143	4.2	36	35	4.7	0.3
young pod + seed (boiled)	89.5	142	2.6	45	17.0	0.7	0.2
leaf (boiled)	91.3	92	4.7	29	18	1.1	0.2

Legumes

English: Lablab bean

Local:

Scientific name: *Lablab purpureus*

Plant family: FABACEAE

Description: A climbing bean which can have vines 1-5 m long. It keeps growing from year to year. The stems can be smooth or hairy. Leaves are made up of 3 almost triangular leaflets. The leaflets are 5-15 cm long and 3-14 cm wide. The side leaflets are somewhat asymmetrical. Often the plants are flushed purple. The flowering clusters are 5-20 cm long. Flowers are often white but can vary from red to blue. The pods are flattened, pointed and up to 12 cm long and 2 cm wide. They can be green, purple or white. Inside there are 3-5 white or dark seeds. Seed pods have a wavy margin. The seeds are 0.5-1.5 cm long. (This bean is similar to Lima bean but the keel of the flower is not spirally twisted, the pod ends more bluntly with a long thin style at the end and the hilum on the seed is longer.)



Distribution: It is a tropical and subtropical plant. It mostly grows between 750 m and 2175 m altitude in the tropics. It is drought resistant and can grow in quite low rainfall areas. Some varieties are short day and some are long day kinds. It suits hardiness zones 9-12.

Use: The young pods, ripe seeds and young leaves are edible, cooked. Flowers can be eaten raw, steamed or added to soups and stews. Dried seeds can be cooked as a vegetable. The seeds can also be sprouted then crushed and cooked. The large starchy root is edible. **Caution:** Many types can be poisonous. They should be boiled and the cooking water thrown away.

Cultivation: Seeds are sown at 30 x 60 cm spacing near stakes or trees. About 20 kg of seed per hectare are required. Fertilising with nitrogen and potash until flowering is recommended.

Production: Young pods are ready 4-6 months after planting and seeds 6-8 months. Pods are often harvested over 2 or 3 years. Pollination and seed setting are reduced in cold weather.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	10.0	1428	22.8	-	-	9.0	-
seed (young)	86.9	209	3.0	14	5.1	0.8	0.4
pod (fresh)	86.7	203	3.9	-	1.0	2.4	-

Legumes

English: Mung bean

Local:

Scientific name: *Vigna mungo*

Plant family: FABACEAE

Description: Mung beans are erect, annual herbs that grow up to 80 cm tall. Sometimes the stems are twining. The stems have a dense covering of yellow or rusty hairs. The leaves are divided into 3 oval leaflets. Each leaflet is 3-10 cm long by 1-5.5 cm wide. The leaf stalks are long but the leaflets stalks are short. The flowers are yellow, 1-2 cm long and occur in dense clusters. The fruit is a hairy pod 4-7 cm long and 5-6 mm wide. Each pod contains 4-10 small, black seeds. The seeds are 4-4.5 mm long by 4 mm wide and have square ends.



Distribution: Mung beans are grown mainly in coastal areas of the tropics but will probably grow up to 1800 m altitude. It suits dry areas. It is drought resistant. It is grown in areas with rainfall of 900 mm per year. It cannot stand frost or long periods of cloud. It is not suited to the wet tropics. It suits hardiness zones 10-12.

Use: The ripe seeds can be eaten. They can be fried and roasted and eaten as a snack. The young pods and young leaves are also edible. The beans are also used for bean sprouts. The seeds can be used in lentil soup, parched and ground into flour for porridge, or baked into bread. The roasted and ground seeds are also used in a spice mixture.

Cultivation: Mung beans are grown from seed. Seed collection is easy. Seed can be broadcast or sown 25 cm apart.

Production: Flowering of mung bean commences after 6 weeks. Plants are self pollinated. Pods are ready to harvest 2-4 months after planting. Pods shatter easily. It is easiest to pull the whole plant, dry them for a week then thresh out the seeds. Seed yields of 450-560 kg per hectare after 80-120 days are common.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (raw)	12	981	22.0	6	-	8.0	-
seed (sprouted)	93.4	88	2	1	11.4	0.7	0.5

Legumes

English: Jack bean

Local:

Scientific name: *Canavalia ensiformis*

Plant family: FABACEAE

Description: A perennial climber, although short kinds do occur. Often it is a more bushy plant than the sword bean. Plants grow up to 1.5 m long. Stems can be hairy. Leaves have 3 leaflets. The leaflets are oval and 5.7-20 cm long by 3.2-11.5 cm wide. The leaf tends to be wedge shaped at the base. The leaf stalks are 2.5-11 cm long. Flowers are red/purple. They occur on flower clusters 5-12 cm long and with flower cluster stalks which are 10-34 cm long. The individual flower stalks are 2-5 mm long. Pods are long and sword shaped. Pods can be 15-35 cm long. Seeds are white with a light brown hilum half as long as the seed. Seeds are 2 cm long, by 1 cm across.

Distribution: It grows in tropical and subtropical places. It requires a fairly high temperature (15°-30°C). It will possibly grow up to 900 m altitude. It is fairly drought resistant and also has some resistance to water-logging and salt in the soil. It can tolerate shade. It can tolerate pH from 4.5-8.0 but does best at about 6.1. The optimum mean annual temperature is 14.4°-27.8°C. Seed germinate between 24-27.5°C. It is a short day plant growing well with a day length of 10-12 hours of sunlight. It can grow in arid places.



Use: The leaves and top shoots are eaten. The very young pods are boiled and eaten. The flowers can be eaten. The young seeds are eaten boiled, roasted, or peeled and cooked. The seeds are also fermented. The ripe seeds are roasted and used as a coffee substitute. **Caution:** The ripe seeds can contain poison and need to be well cooked and the water changed before eating. They are also often left under running water or fermented.

Cultivation: It is grown from seeds. Seeds need to be 2 cm deep. A spacing of about 60 cm is suitable. Plants preferably need a support to climb over. It benefits from a fertile soil but adding nitrogen depresses yield.

Production: Green pods are produced in 3-4 months, but ripe seeds need 6-9 months. Yield of seeds can range from 700-5400 kg/ha.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	10.0	1423	20.4	160	-	4.9	-
pod (fresh)	88.0	155	2.4	-	-	-	-

Legumes

English: Pigeon pea

Local:

Scientific name: *Cajanus cajan*

Plant family: FABACEAE

Description: An upright perennial shrubby legume that can live for 3-4 years. They can grow up to 4 m tall and spread to 1.5 m wide. It has a bushy appearance and a strong deep taproot. The root nodules are round and sometimes lobed. The leaf consists of 3 narrow, green leaflets which are silvery-green underneath. The end leaflet is larger with a longer leaf stalk. The pea shaped flowers are red and yellow and occur on branched flower stalks which stick upwards in the axils of leaves. Pods are long, straight and narrow, often with 4-8 seeds. Seeds vary in shape, size and colour. The pods are slightly hairy. Pods are often 4-8 cm long and have a beak at the end. Pods are constricted between the seeds. Many varieties of pigeon pea occur. Some are dwarf and day length neutral.



Distribution: A tropical plant that requires a tropical or subtropical climate. Plants grow from sea level up to about 1,800 m in the tropics. They can tolerate drought and are suited to a drier climate. They can grow in places with less than 600 mm rainfall per year. They do less well in the wet tropics. They suffer in waterlogged soils and are damaged by frost. It can also tolerate heat. It will grow on poor soils cannot grow on salty soils. It can grow in arid places and suits hardiness zones 10-12.

Use: Young leaves, shoots and pods are eaten. The pods can be used in curries. The leaves and shoots as potherbs. Young seeds are cooked and eaten like peas. Ripe seeds are also cooked and eaten in soups and curries. Bean sprouts can be produced and eaten. Preparation of the seeds for dahl is somewhat complicated.

Cultivation: They are grown from seeds. It is best to sow seeds where the plants are to grow. Seeds normally germinate easily and well. Before sowing seed it helps to soak them in cold water for one day. Seeds store well if kept cool and dry. A spacing of 1.5 m x 1.5 m is suitable. Plants can be cut back and allowed to re-grow. Plants can also be grown from cuttings.

Production: Plants are fast growing. Pods are ready after 5 months. Mature seeds take about 8 months. Plants will often live for 3-4 years. Plants are cross pollinated by insects, or self pollinated.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	10.0	1449	19.5	55	-	15.0	-
pod (young)	64.4	477	8.7	-	-	2.0	-
seed (young, boiled)	71.8	464	6.0	2	28.1	1.6	0.8

Legumes


Common name: African yam bean

Local:

Scientific name: *Sphenostylis stenocarpa*

Plant family: FABACEAE

Description: A vigorous climbing vine. It grows 1.5-2 m high. The leaves have 3 leaflets. They are 14 cm long and 5 cm wide. The flowers are pink, purple or greenish-white. They are 2.5 cm long. They occur on stout stalks in the axils of leaves. The seed pods are smooth and 25-30 cm long by 1-1.5 cm wide. They are flat but have both edges raised. The seeds vary in shape, size and colour. They can be 1 cm long by 0.7 cm wide. They can be cream or brown. Small narrow tubers grow under the ground. They can be 5-7.5 cm long and weigh 50-150 g. The flesh is white and watery.



Distribution: It is a tropical plant that grows from sea level up to 1800 m altitude. It grows in grassland and woodland and sometimes in marshy sites. It can grow in arid places.

Use: The pods, leaves, seeds and tubers are cooked and eaten. They are used in soups or with maize or rice. The hard seeds need to be soaked in water for 12 hours before cooking and being ground. The tubers are cooked and eaten.

Cultivation: It can be grown from seed or tubers.

Production: Tubers are ready for harvest about 8 months after planting.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	9.0	1470	19.2	-	-	-	-
seed (boiled)	67.9	485	7	-	-	1.3	1.1
tuber	64.0	542	3.8	-	-	-	-

Image accessed from:

https://upload.wikimedia.org/wikipedia/commons/thumb/4/48/Sphenostylis_angustifolia%2C_veldblomstappie%2C_Faerie_Glen_NR%2C_a.jpg/1200px-Sphenostylis_angustifolia%2C_veldblomstappie%2C_Faerie_Glen_NR%2C_a.jpg

Leafy greens

English: Goat's horns

Local:

Scientific name: *Sida cordifolia*

Plant family: MALVACEAE

Description: An erect, woody shrub that grows about 0.4-1 m high. It keeps growing from year to year. It is covered with short and long hairs that make the plant feel soft. The leaf stalk is 1-2.5 cm long. The leaves are one after the other and heart shaped at the base. They are toothed at the edge and 1.5-4.5 cm long. The flowers are yellow and occur in the axils of the leaves. The fruit are about 6-8 mm across and have 20 fine bristles on the top.



Distribution: A tropical plant that grows in open waste places in the tropics and sub-tropics. It is common and widely distributed in the Philippines. It grows in hot arid places with a marked dry season. It grows in places with an annual rainfall below 520 mm. It grows in dry sandy soils and can grow in salty soils. It grows below 1100 m altitude. It can tolerate shade and can grow in arid places.

Use: The leaves are edible when cooked.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	6.6	1296	24.2	-	-	79.8	-

Image accessed from

[http://upload.wikimedia.org/wikipedia/commons/f/f4/Sida_cordifolia \(Bala\) in Hyderabad, AP W IMG 9420.jpg](http://upload.wikimedia.org/wikipedia/commons/f/f4/Sida_cordifolia_(Bala)_in_Hyderabad,_AP_W_IMG_9420.jpg)

Leafy greens

English: Quail grass

Local:

Scientific name: *Phyllanthus maderaspatensis*

Plant family: PHYLLANTHACEAE

Description: A herb or small shrub that can grow for more than year. It grows 10-60 cm tall. Leaves are broadly sword shaped and 1-3 cm long by up to 1 cm wide. The fruit are round with 3 lobes. They are 3 mm across.

Distribution: It is a tropical and subtropical plant that grows on sandy soils and limestone. It grows from sea level to 1500 m above sea level.

Use: The leaves are cooked in curries with pulses.



Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	60.9	552	3.0	-	-	59.4	1.0

Image sourced from: https://en.wikipedia.org/wiki/Phyllanthus_maderaspatensis

Leafy greens

Common name: Sunhemp

Local:

Scientific name: *Crotalaria juncea*

Plant family: FABACEAE

Description: An evergreen shrub. It grows to 2.5 m high and spreads to 2 m across. The stem is erect, with silky hairs on the branches. The plant has a strong taproot. The roots have root nodules which have branches and lobes and are 25 mm across. The leaves are narrow, simple and spirally arranged. The leaves can be 12 cm long and with short leaf stalks. The flowers are bright yellow and pea shaped. They occur in clusters of up to 20 blooms. The fruit is a short, inflated, light yellow pod. It is about 3 cm long and 1 cm wide. It is covered with soft hairs, has a groove on the upper surface and a beak at the end. The mature seeds are loose in the pod. They are dark grey, broad and flattened and hooked.



Distribution: A tropical plant. It prefers light to medium well drained soils. It needs an open sunny position. It is drought resistant but frost tender. In Nepal it grows to about 1300 m altitude.

Use: The leaves have been recorded as being eaten. The flowers are pickled.

Cultivation: Plants can be grown from seed. Seed need treatment to break the hard seed coat before planting. Plants can also be grown from cuttings.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	10.2	-	30.1	-	1.39	28.9	-

Image sourced from: <https://www.feedipedia.org/node/313>

Leafy greens

English: Smooth loofah

Local:

Scientific name: *Luffa cylindrica*

Plant family: CUCURBITACEAE

Description: A pumpkin family plant. It is an annual climber up to 10 m long. The stem is five angled and slightly hairy. The tendrils have 2 or 3 branches. Leaves are 10-20 cm across with 5-7 lobes. Male and female flowers are separate and yellow. The male flowers occur as 4-20 flowers together while female flowers are solitary in the leaf axils. Flowers open in the early morning. The fruit is fairly smooth and cylindrical and can be 30-60 cm long. The seeds are black, flat and smooth and 10-15 mm long.



Distribution: A tropical plant that grows well in the tropical lowlands but will also grow in more temperate places. It does best with temperatures of 25-30°C. It is better suited to the drier season as too much rainfall during flowering and fruiting is harmful. Soils should be well drained and moderately rich. It grows in areas with an annual rainfall of 1000-1800 mm. In Zimbabwe it grows up to 1500 m above sea level. It can grow in arid places. It suits hardiness zones 9-12.

Use: The young fruit are eaten as a vegetable. They are skinned and have the centre removed. They can also be sliced and dried for later use. They can be pickled or used in soups, stews and curries. The seeds yield an edible oil after extraction. The seeds are roasted with salt and eaten as a snack. The young leaves and flowers are edible. They are blanched by covering to make them white.

Caution: Older fruit are bitter and fibrous and contain poisonous substances.

Cultivation: Plants are grown from seed which are collected from ripe fruit. Seed are sown 4-5 cm deep and plants are put 1 metre apart. They can be sown in seed boxes and transplanted when 15 cm high. It is best to have a trellis for the plant to climb on or be left to climb over trees. They are often pollinated by insects but can be hand pollinated in the early morning.

Production: Fruit are harvested for sponges when fully mature. Young fruit are ready 2-3 months after planting while fruit mature 4-5 months after planting.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	94.3	79	1.1	-	-	0.7	-
leaf	90	113	5.1	-	95	11.5	-

Leafy greens

English: Yellow ipomoea

Local:

Scientific name: *Ipomoea obscura*

Plant family: CONVULVULACEAE

Description: A slender trailing herb that lies along the ground. It can be a climber or twining. It has a taproot and can keep growing from year to year. The leaf stalks are 1.5 cm long. The leaf blades vary but are long and tapering to the tip with a broadly heart shape base. They are 4 cm long. The flowers occur singly or as a few together in the axils of leaves. The flowers are funnel shaped and 4 cm long and 3 cm across. They are pale yellow or white.



Distribution: It is a tropical plant. It grows up to 1800 m above sea level. It grows in woodland, grassland, savannah and coastal sands. It can grow in arid places.

Use: The leaves are cooked and eaten as a relish. The leaves are added to soup.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	56.6	569	8.8	-	-	28.8	1.1

Image sourced from: https://en.wikipedia.org/wiki/Ipomoea_obscura

Leafy greens

English: Hibiscus

Local:

Scientific name: *Hibiscus rosa-sinensis*

Plant family: MALVACEAE

Description: A shrubby, evergreen, woody, shrub used for hedges. It grows 2-5 m tall. The bark is grey and flaky and has fine stripes. The leaves are bright green and oval with long tips. The edges are entire on the lower leaves. The upper leaves are coarsely toothed. The flowers occur singly in the axils of leaves. Flowers can be single or double. They are bell shaped and 10-15 cm across. There are a range of colours. The fruit are rounded capsules with many seeds inside. The capsules are beaked. Plants usually do not produce fruit in the hot humid tropics.



Distribution: A tropical and subtropical plant, common as an ornamental throughout the tropics. It originally came from China. It thrives on any type of soil. Different types are adapted to sunny or shady places. It grows in open, moist places. It grows where average temperatures are 15-30°C. It is very sensitive to frost and can grow from sea level to 1000 m altitude. It requires a minimum rainfall of 700 mm per year and suits hardiness zones 9-11.

Use: The leaves are eaten cooked. In some places they are pounded before cooking. The flowers are eaten raw or pickled. They are also added to drinks. They are used to colour foods including preserved fruit, sliced pineapple, agar-agar jellies, and cooked vegetables. The fresh flower ovary is eaten.

Cultivation: It is mostly grown from cuttings.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf (dry)	6.4	1339	25.9	-	-	19.6	8.9
leaf (fresh)	76.0	321	2.3	-	-	-	-

Leafy greens

English: Catkin blooming

Local:

Scientific name: *Opilia amentacea*

Plant family: OPILIACEAE

Description: A shrub or woody climber. It grows off other trees and plants. It grows to 4-10 m tall and has stems 20 cm across. The bark is rough and light grey. It has furrows along it and is corky. The aerial branches often hang downwards. The leaves are fairly smooth and leathery. They are 5-14 cm long by 2-5 cm wide. The midrib is prominent underneath the leaf. The leaf has a pointed tip. The leaf stalk is 0.3-0.7 cm long. The new leaves are bright shiny green. The base of the leaves is slightly curved backwards. The flowers are very small and yellow green. They are star shaped. They have a sweet scent. Many flowers occur together on short stalks around a central stem. These occur in the axils of leaves and are 2-3.5 cm long. The white-fleshed, edible fruit can occur singly or in clusters and are oval and fleshy. They are 1.5-3 cm long by 1.2-1.8 cm wide. They are pale yellow or orange when ripe. They have one seed inside. The seed is 21 mm long by 15 mm wide.



Distribution: A tropical plant that grows in tropical Asia. They occur near the beach in monsoon areas. They are often on sandy soil. They need fresh water so are often near streams. It can grow in arid places.

Use: The fruit are eaten fresh. **Caution.** If eaten in large quantities, the fruit can irritate the lips and tongue. Leaves are cooked as a vegetable.

Cultivation: It can be grown from fresh seed. The seed need to be placed on the ground surface, not buried.

Production: It fruits in the wet season. In Tanzania, leaves are collected from April to November.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	9.2	-	14.8	-	3.9	15.7	3.2

Image accessed from:

http://www.westafricanplants.senckenberg.de/images/pictures/opil_opilia_amentacea_rvbli_4_1163_e5e841.jpg

Leafy greens

Common name: Prickly amaranth

Local:

Scientific name: *Amaranthus spinosus*

Plant family: AMARANTHACEAE

Description: An annual plant that grows 60-100 cm tall. It can be erect or lie over. The stems can be either nearly round or angular. The plant branches from the base upwards. Leaves are fairly smooth or hairless but can be tinged purple. The leaf stalk can be 0.5-10 cm long. The leaf blade is oval or sword shaped and 2-7 cm long by 0.6-3 cm wide. There can be a short tip at the top end and it gradually tapers to the base. Flower clusters occur at the sides and these can be single or arranged in compound spikes. The flower clusters at the top can be 3-10 cm long. The top flowers often droop over. Parts of the flowers in the clusters of the leaves form sharp spines 0.5-2 cm long. The upper flowers are male and the lower flowers are female. The seeds are 1-1.2 mm across and flattened.



Distribution: A tropical plant that grows world wide from the tropics to the warm temperate zone. It can grow in sun or light shade. In Tanzania it grows from sea level to 1800 m altitude, in areas with 800-1300 mm annual rainfall. It grows well in moist, damp soil and can also grow in arid places.

Use: The leaves are edible when cooked. The seeds are ground into flour and cooked. **Caution:** This plant can accumulate poisonous nitrates if grown with high nitrogen inorganic fertilisers.

Cultivation: Plants are often self sown but can be grown from seed.

Production: Leaves are often picked early in the season before spines develop. Leaves can be dried and stored.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	91.7	84	3.6	109	46	14.4	0.3

Fruit

English: Boabab

Local:

Scientific name: *Adansonia digitata*

Plant family: BOMBACACEAE

Description: A large tree. It grows up to 25 m tall. It loses its leaves during the year. The branches are thick, angular and spread out wide. The trunk is short and stout and can be 10-14 m around. Often the trunk has deep grooves or is fluted. The bark is smooth and grey but can be rough and wrinkled. The leaves spread out like fingers on a hand. There are 5-9 leaflets. Often the leaves are crowded near the ends of branches. The flowers are large and 12-15 cm across. The petals are white and the stamens are purple. The fruit hangs singly on a long stalk. The fruit has a woody shell. This can be 20-30 cm long and 10 cm across. Inside the fruit are hard brown seeds. They are about 15 mm long. The seeds are in a yellow white floury pulp. The pulp is edible. The thick roots end in fattened tubers.



Distribution: It is a tropical plant that grows in the lowlands. It grows in the hot dry regions of tropical Africa, such as the Sahel. It survives well in dry climates. It grows where rainfall is 100-1000 mm a year. It can tolerate fire. It grows where the annual temperatures are 20-30°C. In most places it grows below 900 m altitude but occasionally grows to 1,500 m altitude. It requires good drainage. It can grow in arid places and suits hardiness zones 11-12.

Use: The young leaves are eaten as a cooked vegetable. The dried leaves are also used to thicken soups. The fruit pulp is eaten raw. It is also used for a drink. The flowers are eaten raw or cooked. The seeds can be eaten fresh or dried and ground into flour then added to soups. They yield a cooking oil. The shoots of germinating seeds are eaten. The young tender roots are eaten. The fattened root tubers are cooked and eaten. The bark is eaten and the dried leaves are used as flavouring.

Cultivation: Trees are grown from seed. The seed remain viable for several years but before planting the seeds must be treated to break the hard seed coat, by soaking the seeds in hot water for several minutes or by cutting the seed coat. Seeds that float in water should not be used. Seeds can be planted in nurseries in plastic bags then transplanted after 6 months. Plants can also be grown from cuttings.

Production: Trees grow quickly reaching 2 m in 2 years. Trees produce fruit after 2-15 years. The plant is pollinated by bats, insects and winds. Trees can last 600 or more years. Fruit can be stored for about a year.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
nut (dry)	7.8	1832	33.7	-		13.9	-
fruit	16.0	1212	2.2	-	360	7.4	6.7
leaf	77.0	290	3.8	-	50	-	-

Fruit

English: Apricot

Local:

Scientific name: *Prunus armeniaca*

Plant family: ROSACEAE

Description: A medium-sized deciduous tree. It grows to 10 m tall. It is a broad spreading shape. The bark is dark red-brown and smooth and shiny. The leaves have stalks. The leaves are broadly oval. They are 10 cm long by 6 cm wide. They are rounded at the base and taper to the tip. They have shallow round teeth along the edge. The leaves are glossy and dark green. The flowers are 2.5 cm across and pale pink or white. They are almost without stalks and have 5 petals. They occur singly on old shoots. The fruit is fleshy and rounded. It is yellow and can be flushed with red. There is a hard covering over the seed. This stone is smooth. The seeds are oval. The flesh is edible. The seeds are edible (but contain toxins).



Distribution: It is native to C. Asia and N. China. It grows in cool areas with hot dry summers. They do best on a free draining soil. In Nepal, plants grow from 2500-3500 m altitude. It does well in areas with 100 cm rainfall, cool winters with 300-900 chilling hours below 7°C and a frost free spring. A soil pH of 6-6.8 is suitable. It needs good sunlight. It suits hardiness zones 5-10.

Use: The ripe fruit are eaten. The kernels can be eaten. If sweet they are eaten fresh and if bitter they are roasted. The seed oil is used for cooking. The fruit are also used for juice and are dried and eaten dry. They are also used for jam, and in pastries, pies, cakes, and picked. **Caution:** The bitter kinds of apricot kernels contain amygdalin which releases cyanide and is poisonous unless removed by cooking.

Cultivation: Plants can be grown from seed but are often grafted onto rootstocks. The flowers are self-fertile. Plants require some winter chilling.

Production: Trees fruit in 3-4 years. Fruit are picked by hand.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit (raw)	86	117	0.6	96	7	0.4	0.1
fruit (boiled)	82.6	92	0.4	59	5	0.3	0.1

Image sourced from: <https://www.gardenia.net/plant/prunus-armeniaca-harcot>

Fruit


English: Mango

Local:

Scientific name: *Mangifera indica*

Plant family: ANACARDIACEAE

Description: An erect, branched evergreen tree. It can grow to 10-40 m high and is long lived. (Trees grown by vegetative means are smaller and more compact.) Trees spread to 15 m across. It has strong deep roots. The trunk is thick. The bark is greyish-brown. The leaves are simple and shaped like a spear. Some kinds of mangoes have leaves with a wavy edge. They can be 10-30 cm long and 2-10 cm wide. They are arranged in spirals. The leaf stalk is 1-10 cm long and flattened. Leaves are often brightly coloured and brownish-red when young. These tender leaves which are produced in flushes become stiff and dark-green when mature. The flower stalks are at the ends of branches. They are 10-50 cm long and branching. Up to 6,000 flowers can occur on a stalk. Most of these are male and up to 35% have both male and female flower parts. Fruit are green, yellow or red and 2.5-30 cm long. The fruit hang down on long stalks. The outside layer of the seed is hard and fibrous and there is one seed inside. Several embryos can develop from one seed by asexual reproduction. The fruit shape and colour vary as well as the amount of fibre and the flavour. India has many varieties and they cannot tolerate humidity.



Distribution: A tropical and subtropical plant. It grows in the lowlands. It grows from sea level up to 1300 m altitude in the tropics. It does best in areas below 700 m and with a dry season. Rain and high humidity at flowering reduces fruit set. It thrives best where temperatures are about 25°C but will grow with temperatures from 10-42°C. Temperatures of 0°C will damage young trees and flowers. Low temperatures (10-20°C) at flowering time will reduce fruiting. As temperatures get lower due to latitude or altitude, fruit maturity is later and trees become more likely to only have good crops every second year. Mangoes can grow on a range of soils. In wetter areas soils with less clay are better. They can withstand occasional flooding. A soil pH of 5.5-6.5 is best. Soils with pH above 7.5 cause plants to develop iron deficiency. It grows in the Sahel. It can grow in arid places. It suits hardiness zones 11-12.

Use: Ripe fruit are eaten raw. Unripe fruit is pickled. Seeds can be eaten cooked. They are boiled or roasted. They are made into meal by powdering. Young leaves can be eaten raw or cooked. Amchur is made from the dried unripe fruit. This is used in curries, and pickles and chutneys. The seed kernels are used for famine food in India. They are boiled, roasted or soaked to remove the bitterness. **Caution:** The sap from the tree or fruit can cause skin problems with some people.

Cultivation: Trees are grown by planting fresh seed and they can be transplanted. Mangoes vary in their ability to breed true from seed. When more than one seedling emerges from the seed some of these are asexual and breed true. Clean seed germinate best if they are treated at 50°C for 20 minutes, then planted on their edge with the round bulge upwards and near the soil surface. The husk around the seed should be removed. Seeds germinate in 3-6 weeks. The strongest growing seedlings from this seed are used and the others thrown away. The seedlings from the folds of the seed are vegetative while the seedling from the centre of the seedling near the stalk end may be sexual and show variation from type. Other seeds only produce one seedling and these normally

vary and can be different from the parent tree. Plants can be propagated by budding, or by grafting using in-arching. This is not easy and care is required. In wetter places, flowers need to be protected with fungicides to enable fruit to form. If organic manure is used this should not be directly in the planting hole nor immediately against the new plant. Young transplanted seedlings need regular watering. A spacing of 6-12 m between plants is used. Wind protection is advisable to prevent fruit rubbing and getting damaged. Trees should only ever be lightly pruned as fruit develop on new growth and heavy pruning can reduce flowering. Flowering can be brought about by foliar sprays of potassium nitrate.

Production: Seeds germinate after about 20 days. Seedling trees produce after 4-6 years and increase in production up to 20 years. Trees often bear better each second year. Rain at flowering reduces fruit setting. Fruiting is at the end of the year. Fruit take 4-5 months to mature. Fruit vary in weight from 200-1000 g. Trees can produce one million flowers but only 500 fruit. Trees last for many years.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	83.0	253	0.5	54	30	0.5	0.04
leaf	82.1	226	3.9	-	60	2.8	-

Fruit

English: Cape gooseberry

Local:

Scientific name: *Physalis peruviana*

Plant family: SOLANACEAE

Description: A perennial herb that grows 45-90 cm tall. They are often grown as annuals. It is hairy and slightly branched. The spreading branches are purplish and ribbed. The leaf blade is 6-15 cm long by 4-10 cm wide. The leaves are heart shaped at the base and taper to the tip. They are slightly wavy and toothed along the edge. The flowers occur singly and hang down in the axils of leaves. The flowers are white with violet anthers and slightly spotted petals. The fruit is a berry 1-1.5 cm across. They are orange-yellow or pale brown. This is inside an inflated husk. The seeds are yellow and 2 mm across. There are several named cultivated varieties.



Distribution: A temperate plant that grows in the tropical highlands. It suits warm climates and does best in warm sunny conditions. It needs well drained soil. Plants are not killed by a slight frost but it grows best free from severe frosts and strong winds. In Indonesia plants are found from 700-2300 m altitude, but fruit best above 1500 m. It can grow in arid places and suits hardiness zones 8-10.

Use: The ripe fruit are eaten fresh or cooked. They are used for jam. They can be dried, preserved, stewed, pureed, or used in pies, cakes, jellies and sauces. Roasted seeds are pickled. The leaves have been used instead of hops in beer. The leaves are also used as a potherb.

Cultivation: Plants are grown from seed that is broadcast over the soil. Seeds should be sown 1.5 cm deep in loose soil. Seed germinate irregularly. Plants should be spaced 45 cm apart. In the tropics, plants keep growing from year to year, but in the subtropics they regrow from seed each year. Plants can be grown from softwood cuttings from the upper parts of the shoots. Seedlings can be transplanted.

Production: Plants produce fruit in 1 year.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit (mature)	84.2	201	2.0	36	30	1.5	-

Fruit

English: Tamarillo

Local:

Scientific name: *Cyphomandra betacea*

Plant family: SOLANACEAE

Description: A small, soft-wooded fruit tree in the tomato and tobacco family. It grows as a shrub up to 4 m tall and is shallow rooted. The brittle stem is 5-10 cm across. The leaves are large (25 cm x 12 cm) and soft and heart-shaped at the base. The sweet smelling flowers are 1-2 cm across and occur in loose clusters near the ends of branches. Flowers are normally self-pollinating but need wind. The red or orange egg-shaped fruit are 6-12 cm long and hang off the ends of the branches. The skin of the fruit is somewhat tough, but the flesh around the seeds is soft and juicy. The edible flesh varies from yellow to dark purple. The edible seeds are black, thin and nearly flat. There are several named cultivated varieties.



Distribution: A subtropical plant suited to tropical highlands. It grows at 750-2200 m altitude in the equatorial tropics, and up to 3000 m in the continental tropics. It is more cold hardy than tomato, and does best where average temperatures are 15-21°C. It cannot stand much frost, although mature trees will survive short periods of light frosts. It can't stand water-logging or drought. Trees can grow under shade but do better in sun, unless it is hot and dry. Deep, fertile, permeable, disease-free soil is best. Plants don't fruit at low altitudes in the tropics due to constant high temperatures. They need shelter from wind. It suits hardiness zones 9-11.

Use: The fruit is eaten raw or cooked. The layer just under the skin can be bitter. The layer around the seeds is sweet. The seeds can be eaten, or strained out. The outside skin is easily removed by immersing fruit in boiling water for 1-2 minutes. The fruit can be boiled, stewed, grilled, baked or pickled, or used in jams, jellies, chutneys, conserves, pies, preserves and sauces.

Cultivation: They can be grown from seed or cuttings. Seeds grow better if they are washed and dried, then placed in a freezer for 24 hours before planting out. Seeds produce a high branched erect tree. Cuttings produce a lower bushier plant. Cuttings of 60-90 cm long stalks are suitable and should be taken from 1-2 year old wood which is 1-2 cm thick. Root cuttings can also be used. It has shallow roots, so needs careful weeding. A spacing of 3 m apart is sufficient. Because the roots are easily damaged by nematodes, plants grafted or budded onto nematode resistant rootstocks will live longer. Root rot fungus can make it difficult to get plants established in old gardens. High humidity can also cause stem rots. Young plants can be pruned to produce lower branches and fruiting. Fruit is produced on new branches, so branches which have fruited can be pruned out. Flowers are self-compatible so pollination by wind can occur within the one plant.

Production: Fruiting commences in the second year when grown from seed. Trees bear hundreds of fruit throughout the year. Yields of 20 kg per plant are possible. Trees live 5-6 years, although if infested with root knot nematodes, may live only 3-4 years. Fruit are ready to harvest when the red or yellow colour develops. Fruit can be stored above 3°C in a refrigerator for 10 weeks.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	86.2	113	2.0	500	28	0.7	-

Fruit

English: Small-leaved white raisin

Local:

Scientific name: *Grewia tenax*

Plant family: MALVACEAE

Description: A shrub that grows up to 2 m tall. It often lies along the ground. The leaves are small and nearly round. They are 5 cm long. They have 5 main veins. The tip of the leaf is pointed and the edge has teeth. The lower leaf surface has hairs. The flowers are white and occur singly. They are 2 cm across. They are on long slender branches. The fruit are orange-red, smooth and edible. They have 1-4 lobes. They are the size of a small maize grain.



Distribution: A tropical plant that grows in arid zones. It occurs in very dry woodland and semi-desert scrub. It grows on rocky and gravelly soils. It grows in the Sahel. It is often near temporary pools. It grows in areas with over 200 mm rainfall. It can tolerate salt. In East Africa it grows between sea level and 1500 m altitude.

Use: The fruit are eaten fresh and raw. They are also dried for eating later. They are added to grains in porridge. A drink is made by soaking the fruit overnight then pressing, sieving and sweetening the juice. The seeds are edible.

Cultivation: Plants can be grown from seeds.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit (dry)	9.2	1157	5.5				
fruit	59.1	-	4.5	-	161	125	-

Image accessed from:

[http://www.southernafricanplants.net/photocollection/batch005/medium/G/TILLIACEAE Grewia tenax Arandis 20090215 072 \(1\).jpg](http://www.southernafricanplants.net/photocollection/batch005/medium/G/TILLIACEAE Grewia tenax Arandis 20090215 072 (1).jpg)

Fruit

English: Mandarin

Local:

Scientific name: *Citrus reticulata*

Plant family: RUTACEAE

Description: A small, evergreen tree that grows 4-8 m tall and 2 m across. The stem is erect, branching and thorny. The leaves are dark green, and long and narrow in shape. They are 3-4 cm long. There is only a narrow wing on the leaf stalk. It has a few or no spines. The flowers are white and star-shaped. They are 2.5-4 cm across and have a scent. Fruit are almost round and the skin peels off easily. The fruit are 4-8 cm long. The flesh is red, juicy and sweet.



Distribution: It is grown in many tropical countries. It is the hardiest of the citrus. It grows from sea level up to 2300 m altitude in the tropics. It does best between 800 m and 1200 m altitude. A well-drained soil is needed. It also prefers a drier climate. It is drought and frost resistant. It needs a temperature above 3-5°C. It suits hardiness zones 9-11.

Use: The fruit are eaten fresh.

Cultivation: Trees are often grown from seed. Some breed true from seed. Seedling trees take a long time to start producing fruit. Budded trees are best. A spacing of about 8 m between trees is suitable. Several seedlings can grow from one seed. Using seedlings of seeds with three or more shoots helps produce trees true to type. Cuttings or layering can also be used.

Production: Fruit tend to be produced seasonally. The season is often from April to August in the southern hemisphere.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	87.6	184	1.5	42	136	0.8	-

Fruit

English: Small wild medlar

Local:

Scientific name: *Vangueria infausta*

Plant family: RUBIACEAE

Description: A shrub or small deciduous tree. It grows 3-4.5 m tall. It can be 7 m tall. It has a smooth grey trunk. The bark becomes rough with age. The branches are short and stout. They are soft and covered with red hair when young. The leaves are light green and leathery. The leaves are 5-30 cm long and 3.8-18 cm wide. They are often egg-shaped. The tips can be round or blunt and they are densely hairy. The veins are like nets. The flowers are greenish-white or yellow. They are small. They grow in clusters. The fruit are round and 3.8 cm across. They have a shallow crown on top and are covered with a light brown leathery skin. They have 3 seeds inside. Each seed is in a separate compartment.



Distribution: A tropical plant. It grows in the lowlands and the highlands. It grows on stony and sandy ground. It grows in shady places. It can tolerate frost. It can tolerate drought. It grows up to 3333 m above sea level. It can grow in arid places.

Use: The fruit are eaten raw. The fruit can be used to make a sauce or a pudding. When sugar is added, the fruit looks and tastes like apple sauce. The fruit can be dried and stored. It is later soaked and cooked. The seeds are cracked and the kernels extracted and eaten. They are roasted. They are also used as flavouring. The leaves are cooked as a green vegetable.

Cultivation: Plants can be grown from seeds. The fruit are allowed to dry and then soaked in water before planting. Using fertiliser, manure or early irrigation reduces the survival and early growth of young plants. They are adapted to low fertility and seasonal rainfall.

Production: It grows quickly. Fruit are edible January to April.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	64.4	498	1.4	-	11.5	24.4	7.1

Image accessed from: <http://www.plantzafrica.com/planttuv/voteplant.php>

Vegetables

English: Bullrush millet

Local:

Scientific name: *Pennisetum glaucum*

Plant family: POACEAE

Description: An annual grass that grows to 3 m tall. The leaf blades are 20-100 cm long by 2-5 cm wide. The flower is dense and 40-50 cm long by 1.2-1.5 cm wide. They also vary in shape and size. Plants that tiller produce smaller heads. The species varies a lot. There are 13 cultivated, 15 weed and 6 wild races of this grass. It has a cylindrical ear like a bullrush. The grains are small and round and have a shiny grey colour like pearls. There are thousands of cultivated varieties.

Distribution: A tropical plant that suits regions with a short growing season. It grows in areas with less than 600 mm of rainfall. It is replaced with sorghum between 600-1200 mm rainfall and then by finger millet or maize above 1200 mm rainfall. It is important in the drier areas of India and Pakistan. It can grow in arid places.

Use: The seeds are eaten like rice. They are also ground into flour and made into bread and cakes. They are used to make alcoholic drinks. They are mixed with other grains and seeds to make fermented foods. Some kinds have sweet stalks that are chewed. The young ears can be roasted and eaten like sweet corn.

Cultivation: Plants are grown from seed. It is usually sown directly into the field. The plant density is adjusted to suit rainfall and soil fertility. The spacing is 45 cm apart up to 200 cm apart. It is also intercropped with other crops such as cowpea, sorghum and peanut. Crops are normally weeded 2 or 3 times.

Production: It takes from 75-180 days to maturity. The heads can be picked by hand or the plant removed. Some types need to be picked 2 or 3 times as heads mature.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	11.6	1442	10.5	-	-	6.5	1.7



Vegetables


English: Lentils

Local:

Scientific name: *Lens culinaris*

Plant family: FABACEAE

Description: A slender, annual plant. They grow to about 25-40 cm high. It is erect with many branches. Plants are softly hairy. The leaves are compound with leaflets along the stalk. There is usually a tendril at the end. There are 4-7 pairs of leaflets and these do not have a stalk. They are sword shaped and 1.3 cm long. The flowers are in the axils of leaves. There are 1-4 flowers in a group. The flower stalk is slender. The flowers are small and up to 8 mm long. The flowers are bluish. The pods are oblong and 1.3 cm long. There are 1-2 seeds per pod. The pods are flat and the seeds are about 3-6 mm across. There are also large seeded kinds with seeds 6-9 mm across. The seeds are lens shaped, round and curved out on both sides. The seeds become reddish brown when ripe. There are several named cultivated varieties.



Distribution: A plant of warm temperate and tropical zones. It prefers a sandy soil in a warm position. It produces most seed when grown on poorer soils. They grow in subtropical, warm temperate and high altitude tropical places. In India they grow from sea level to 3500 m altitude. In Nepal they grow to about 1000 m altitude. In Ethiopia it grows between 1600-2350 m above sea level. They can grow on a range of soils. It suits hardiness zones 7-11.

Use: The seeds are cooked, sprouted or eaten raw. Young seedpods can be cooked and eaten. The ground seed can be used with cereals. The seeds are often eaten in soups and stews. They are served as Dahl in India. Lentil flour can be mixed with cereal flour to bake bread. The sprouted seeds are eaten in salads, vegetable dishes and soups.

Cultivation: Seed are sown where they are to grow. Plants are normally self pollinated but cross pollination can occur. In India it is often grown mixed with rice. When grown as a pure stand it can be broadcast or planted in rows.

Production: Yields of 2 ton per hectare are possible. For sprouts, the seeds are soaked for 12 hours in warm water then allowed to sprout for 5 days. Crops mature in about 3.5 months.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
sprouted & cooked	68.7	423	8.8	4	12.6	3.1	3.1
split & boiled	72.1	420	7.6	20	-	2.4	1.0

Vegetables

English: Okra

Local:

Scientific name: *Abelmoschus esculentus*

Plant family: MALVACEAE

Description: A tropical annual herb that grows erect, often with hairy stems. It mostly grows about 1 m tall but can be 3.5 m tall. It becomes woody at the base. The leaves have long stalks up to 30 cm long. Leaves vary in shape but are roughly heart shaped with lobes and teeth along the edge. Upper leaves are more deeply divided than lower ones. The flowers are yellow with red hearts. The fruits are green, long and ribbed. The seeds are 4-5 mm across. They are round and dark green.



Distribution: A tropical plant that suits the hot humid tropical lowlands but is unsuited to the highlands. It is very sensitive to frost. It can grow in salty soils. It grows best where temperatures are 20-36°C. It can grow well in dry climates with irrigation. It suits hot humid environments. It does best on well drained well manured soils but will grow on many soils. A soil pH of 5.5-7.0 is best.

Use: Pods are eaten cooked. They are slimy, but less so if fried. Dried powdered seeds can be used in soups as a thickener. They can also be pickled. Young leaves can be eaten cooked. They can be dried and stored. Flowers can also be eaten. Okra is frozen and canned. The seeds are roasted and used as a coffee substitute.

Cultivation: They are grown from seeds, which are easy to collect. They need high temperatures for germination (over 20°C) and a sunny position. Often seeds are soaked for 24 hours before sowing to give quick germination. Seeds are sown 1.5-2.5 cm deep with 2-3 seeds per hole. Later these are thinned out to one plant. Seeds can be sown in nurseries and plants transplanted. Pinching out the tops of plants when 30 cm high encourages branching. A spacing of about 90 x 45 cm is suitable. About 8-10 kg of seed are required for one hectare. Most kinds respond to fertiliser. Seeds do not breed true and can cross with other kinds of okra growing nearby. This is not normally a problem but simply means plants and fruit are not all the same.

Production: Plants maintain production if the fruits are harvested regularly. Plants are ready to harvest 8-10 weeks after sowing. Seed yields of 500-800 kg per hectare are recorded. Pod yields of 4-6 tonnes per hectare occur. It takes 2-4 months from sowing to harvest of young pods. Pods develop 5-10 days after flowering. Pod harvests can continue for 1-2 months. Leaving pods on the plants stops new pods developing.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	9.2	1721	23.7	-	-	-	-
leaf	81.0	235	4.4	116	59	0.7	-
pod (fresh)	88.0	151	2.1	185	47	1.2	-
fruit (cooked)	90.0	134	1.9	58	16.3	0.5	0.6

Vegetables

English: Pumpkin

Local:

Scientific name: *Cucurbita maxima*

Plant family: CUCURBITACEAE

Description: A pumpkin family plant. It is a creeping vine with tendrils. It is an annual plant. The stems are soft and round in cross section. The leaves are large and hang loose. They are dark green and kidney shaped. The edges of the leaves are entire. There are large nodes at the base of the leaf. The tendrils are fairly stout and are divided half way along their length into many branches. Male flowers are carried on long upright stalks. The 5 petals are united into a long yellow tube. The female flowers are larger



than the male and are fewer in number and carried on shorter stalks. The fruit varies in size, colour and patterns on the skin. They can be round, oval or flattened, with yellow, orange or green skin. The surface can be smooth or rough and warty. The flesh is yellow and edible. The seeds are in the centre. The seeds are white or brown. They are flattened but plump and have a slanting scar at the top. The seeds are edible. (*C. moschata* does not have hairy stems but has fruit with a thickened stalk near where it joins the fruit.) There are a large number of cultivated varieties.

Distribution: A subtropical plant that grows from sea level to 2400 m altitude. They need a fertile soil. *C. moschata* is better suited to coastal areas. They are frost sensitive but better suited to cooler areas than *C. moschata*. It can grow in arid places. It suits hardiness zones 8-11.

Use: The young leaf tips are eaten cooked. They can also be dried and stored. The fruit can be eaten cooked. They are baked, boiled, fried, steamed or mashed. They are used in pies and cakes. The seeds are edible, raw or roasted. They are also ground into a meal. The male flowers are eaten after removing the stamen and calyx.

Cultivation: They are grown from seed. Usually 2 or 3 seeds are planted together in a mound. The distance apart depends on the cultivar. Some kinds are better for leaf tips. It is good to save seed of adapted varieties.

Production: Fruit are ready for harvest after about 3-4 months. Seed can be saved from fruit for re-sowing, but as pumpkins cross-pollinate, different types become mixed.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	6.9	2264	24.5	38	1.9	14.9	7.5
fruit	69.6	439	1.4	-	-	-	-
leaf	88.0	160	4.9	260	28	2.5	0.9
flower	88.7	107	1.4	173	14	0.8	0.1

Vegetables

English: Jute

Local:

Scientific name: *Corchorus olitorius*

Plant family: MALVACEAE

Description: An annual plant. It is upright, branching, and slightly woody. Plants vary in height, shape, leafiness and hairiness. Plants grown for leaves are usually only 30 cm tall. They also have many branches. Leaves are shiny and have leaf stalks. The leaves have teeth along the edge. The tips of the lowest leaves in each side, have long bristle like structures. Small clusters of yellow flowers grow in the axils of the leaves. The fruit are ridged capsules. They can be 7 cm long. These have partitions across them between the seeds. A ripe capsules contains 180-230 seeds. The seeds are dull grey and with four faces and one long point. Each seed has one pale line along it.



Distribution: A tropical plant. It is mostly coastal, below 250 m altitude. Temperatures of 22-35°C are suitable. It can stand both drought (2-3 weeks) and water-logging, except when young. A well-drained soil is best. They require humus-rich soils. A soil pH of 5.5-7.0 is best, but they can grow in soils with pH up to 8.5. They also need adequate moisture for good leaf production. A rainfall of 1000 mm is suitable. A high relative humidity (80-90%) is best. It produces seeds when day lengths are short. It grows in most African and Asian countries.

Use: The young leaves and stem tops are eaten cooked. They are slimy unless fried. They are also used to make a thick soup. Leaves can be sun dried, pounded to flour, then stored for a long time.

Cultivation: Plants grow from seed, and they can be transplanted. Seeds are often broadcast into fine seed beds at the beginning of the wet season. Mixing the small seeds with sand makes it easier to sow them evenly. Often seeds are slow to start growing. This can be overcome by soaking them in hot water. A spacing of 20-30 cm between plants is suitable. For vigorous varieties this could be increased to 45-50 cm. Seeds are saved from pods for re-sowing.

Production: First leaves can be harvested after 5-6 weeks. Tips about 20-30 cm long are picked. Production of edible green tips, is not large. 7-8 kg of leaf tips can be harvested from 3-8 pickings over 3-4 months. Seeds can be collected after 13-15 weeks. If seeds of a particular variety are desired, it is necessary to grow these plants 16 m away from other plants, to avoid cross pollination. Seeds can be stored for 8-12 months in well-sealed jars.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf (raw)	80.4	244	4.5	574	80	7.2	-
leaf (cooked)	87.2	155	3.4	156	33.0	3.1	0.8

Vegetables

English: Marrow

Local:

Scientific name: *Cucurbita pepo*

Plant family: CUCURBITACEAE

Description: A bristly hairy annual vine in the pumpkin family. It has branched tendrils. The stems are angular and prickly. The leaves are roughly triangular. The leaves have 5 lobes which are pointed at the end and are toothed around the edge. Male and female plants are separate on the same plant. Male flowers are carried on long grooved flower stalks. Female flowers are borne on shorter more angular stalks. The fruit stalks have furrows along them but are not fattened near the stalk. The fruit vary in shape, size and colour. Often they are oval and yellow and 20 cm long by 15 cm wide. The seeds are smaller than pumpkin and easy to separate from the tissue. The scar at their tip is rounded or horizontal, not oblique. There are a large number of cultivated varieties.



Distribution: A subtropical plant. They are more suited to drier areas. They are frost sensitive, and grow best with day temperatures between 24-29°C and night temperatures of 16-24°C. It suits tropical highland regions. It suits hardiness zones 8-11.

Use: The young fruit are cooked and eaten. They can be steamed, boiled or fried. They are used in pies, soups, stews and cakes. The young leaves and the ripe seeds can also be eaten cooked. The seeds are dried, salted and toasted and eaten as a snack food. The seeds can also be pressed to produce oil. The sprouted seeds are used in salads. Flowers and flower buds can be eaten boiled. They can be dried for later use.

Cultivation: They are grown from seeds. The seeds germinate after one week. They can be grown from cuttings. They are best planted on mounds. A spacing of 2-3 m between plants is needed. Hand pollination assists fruit setting. Plants can also be grown from cuttings as plants root at the nodes.

Production: The first usable immature fruit are ready 7-8 weeks after planting.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	3.7	2266	29.4	-	-	7.3	-
leaf	89.0	113	4.0	180	80	0.8	-
fruit (mature)	92.0	105	1.6	17	16	2.4	-
fruit	91.3	102	1.1	-	12	0.8	0.2
yellow fruit	92.0	97	1.0	180	8	1.4	-
immature fruit (raw)	92.0	92	1.5	-	9	0.4	0.1

Vegetables

English: Horseradish tree

Local:

Scientific name: *Moringa oleifera*

Plant family: MORINGACEAE

Description: A small, soft-wooded tree that grows 9-12 m tall. The tree loses its leaves during the year. The bark is grey, thick, corky and peels off in patches. The leaves are pale green and the leaf is divided 3 times. The whole leaf is 30-60 cm long and the leaflets are usually oval and 1-2 cm long. The leaflets are jointed with a gland near the joint. The flowers are pale yellow. They occur in long sprays 30 cm long. Each flower has 5 petals and of these one is erect and 4 are bent backwards. The fruit is a long capsule 30-100 cm long by 2 cm wide. The seed capsules are up to 45 cm long. They are roughly triangular in shape. The seeds have 3 wings. Often the fruiting kinds are grown as annual plants.



Distribution: A tropical and subtropical plant. They suit the dry lowland areas and grow up to 1350 m altitude in the tropics. They are not hardy to frost. They cannot tolerate water-logging. A pH of 6-7.5 is suitable. It can grow in arid places. It suits hardiness zones 9-12.

Use: The young tops and leaves are eaten cooked. They are eaten as potherbs or used in soups and curries. They can be dried and stored for later use. The very young long pods are eaten cooked, especially in curries and soup. They are also pickled. The young seeds are eaten roasted or fried. Sometimes the roots are used as a horseradish substitute. A gum from the bark is used as seasoning. The bark is used for tea. The roots, leaves, flowers and fruits are eaten cooked in water and mixed with salt and chili peppers. The oil expressed from the seeds is used in salads.

Cultivation: It is best to grow plants from 1 metre long cuttings but they can be grown from seed. They can be used as a hedge and pruned regularly to produce more leaves. Properly dried seed can be stored for a long time in sealed containers in a cool place. Normally perennial types are grown from cuttings and annual types are grown from seed.

Production: Trees are fast growing. They can be pruned or topped. With one variety the tree flowers and fruits continuously while with the other variety there are flowers and fruit once per year. The fruit ripens 3 months after flowering. Annual types produce fruit 6 months after planting. Leaves are best dried in the shade to retain more of their Vitamin A.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
leaf	76.4	302	5.0	197	165	3.6	-
flower	84.2	205	3.3	-	-	5.2	-
leaf (boiled)	87	189	4.7	40	31.0	2.0	0.2
pod (raw)	88.2	155	2.1	7	141	0.4	0.5
seed	6.5	-	46.6	-	-	-	-

Vegetables

English: Bitter cucumber

Local:

Scientific name: *Momordica charantia*

Plant family: CUCURBITACEAE

Description: A pumpkin family plant. It is a slender annual climber with flowers of both sexes on the one plant. It has simple tendrils and vines can be 4 m long. It has bright green lobed leaves 5-12 cm long on thin leaf stalks 3-10 cm long. The flowers have a sweet smell and 5 small, yellow petals. Fruit are green when young and orange when ripe. The fruit have a lumpy appearance, with ridges along its length and when fully ripe burst open. It has bright red covering on the seeds inside. The seeds are pale brown and 10-16 mm long and 7-10 mm wide. Considerable variation in the fruit occurs between varieties.



Distribution: A tropical plant that grows from sea level up to about 500 m and will probably grow to 1000 m altitude in tropical regions. They require a well-drained soil preferably rich in organic matter. Seeds do not germinate below 15°C. Plants grow best with temperatures of 18-35°C. A soil pH of 6.5 is best. It suits hardiness zones 9-12.

Use: The young bitter fruit are cooked and eaten. They are boiled, stuffed, fried or pickled. They are used in soups, stews and stir-fried dishes. The seed mass of the ripe fruit is used as a food flavouring. The leaves are also cooked and eaten as a flavouring. The tender shoots and leaves are sometimes eaten. **Caution:** The leaves are considered to cause diarrhoea and vomiting.

Cultivation: Plants are grown from seed. For large scale plantings, 6-7 kg of seed are required for planting one hectare. Seeds are planted at 50 cm spacing in the place where the plants are to grow and need a stick to climb up. Often plants are grown on raised beds 2 m apart with 0.5 m between plants. The seed has a hard seed coat and germinates slowly. Soaking seeds for 24 hours before sowing gives a quicker more even germination. Regular watering is required.

Production: Fruit are ready to harvest 45-55 days after planting. Fruit should be harvested when young and tender. Once fruit have begun to change colour to yellow they are past maturity for eating. Early removal of young fruit also ensures continuous fruit setting. This can allow 6-8 successive pickings of fruit. Fruit on the plant are sometimes wrapped in paper to prevent fruit fly damage. Seed well stored can remain viable for 4-5 years. The young bitter fruit are cooked and eaten. The fruit is blanched or soaked in salt water to reduce the bitter taste.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	8.6	2020	18.6	-	-	-	-
leaf (raw)	84.7	252	5.0	44	170	7.1	0.3
leaf tip (boiled)	88.7	146	3.6	173	57	1.0	0.3
fruit	93.6	105	1.2	-	-	0.2	-
pod (boiled)	94.0	79	0.8	11	33	0.4	0.8
pod (raw)	94.0	71	1.0	380	84	0.4	0.8

Nuts, seeds, herbs and other foods

English: Milk plum

Local:

Scientific name: *Englerophytum magalismontanum*

Plant family: SAPOTACEAE

Description: A shrub or small tree that usually grows up to 15 m tall, although can occasionally grow up to 33 m. Often the roots are twisted around a rock. It has a leafy rounded crown. The trunk is thin and has flutes along it. The stem is mostly short and thick and the branches can touch the ground. Often the trunk has wart-like growths. The bark is grey and rough. The firm, leathery leaves are simple and alternate and are often crowded towards the ends of branches. The leaves are oval and dark green and shiny on top but brown and hairy underneath. The



young shoots and twigs have dense rusty hairs. The upper leaf surface often has a bloom. The under surface of the leaves has silver or red hairs. The leaves are usually 5-10 cm long but can be 30 cm long. The leaf stalks are 0.6-2.5 cm long. Flowers and fruit are produced on old wood. The flower buds occur in clusters, often on knobs on the branches and trunk. The fruit are 2.5 cm long. They are bright red when ripe. There is a small point at the end. The fruit have a leathery skin and 2 stones inside. The flesh is purple and they have a sticky latex. The seeds are oval and 1.7 x 1.4 cm.

Distribution: A tropical plant that occurs near quartzite and granite rocks. It can grow in the lowlands and up to 1800 m in South Africa. They mostly grow in coastal areas in deciduous woodland. They are somewhat drought resistant and can tolerate frost. They grow on a range of soils. They do best on dry well drained soils. It can grow in arid places.

Use: The fruit can be eaten raw when immature or when ripe. They are also used for beer, wine and jams.

Cultivation: They can be grown from seed or cuttings. Fresh seed germinate quickly. Cuttings from old wood can be used. Fruit are ready to harvest 4-5 months after flowering.

Production: Trees grow fairly quickly. In Tanzania, fruit are collected from April to June.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
fruit	74.3	404	0.9	-	13.1	0.03	

Nuts, seeds, herbs and other foods

English: Cashew

Local:

Scientific name: *Anacardium occidentale*

Plant family: ANACARDIACEAE

Description: An evergreen tree, with spreading branches, growing 7-14 m tall. The canopy can spread to 12 m. The roots grow deeply and spread widely. The shiny leaves are pale green and large. They are 10-15 cm long by 6-8 cm wide. They have fine veins. The flowers are produced on the ends of the branches. They are red in colour. The kidney-shaped nut is about 3 cm long and is borne below the "apple" which is really a fleshy stalk.



Distribution: It is a tropical plant that suits the lowland tropics but will grow up to about 1,200 m altitude. It only bears well in dry areas because of blight of the flowers. It grows best in temperatures of 22-26°C. A rainfall of 1750 mm per year is considered suitable but good yields have been obtained with rainfall of 750 mm. It can grow on poor soils but needs good drainage.

Use: The fleshy "apple" is edible but acid until very ripe. It is used for jams, drinks, candy, chutney and pickles. The nut is eaten after roasting. The young shoots and leaves are edible. They are picked during the rainy season and eaten fresh with hot and spicy dishes. **Caution:** The oil of the nut can blister the skin until roasted. The apple is used to make spirits.

Cultivation: It is usually grown from seeds. Seeds germinate poorly and slowly. Only nuts which sink in water (or a solution of 150 g of sugar in a litre of water) should be planted. Seeds are sun dried for 2-3 days to improve germination. Seeds can be sown in a nursery then transplanted, or more commonly, are sown directly. Trees are spaced 7-10 m apart. The crop is cross pollinated mostly by insects. For good production, complete fertiliser or appropriate organic material should be applied. Pruning to shape the tree is often undertaken in the first 2-3 years. Cashews are often planted scattered in gardens or amongst other trees. Clearing under the tree prevents fire and makes finding nuts easier. Allowing nuts to fall before harvesting ensures only ripe nuts are collected. Resin in the cashew nut shell can damage hands and discolour the nuts. Roasting the nuts before removing the kernel avoids this.

Production: Trees commence bearing after 3 years. Fruit production is seasonal, normally October-January. Mature nuts are produced in 2-3 months. Yields of 80-200 kg of nuts per hectare are normal. Trees reach maximum production after 10 years and last for about 100 years.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
nut	4.0	2478	17.5	-	-	2.8	4.8
leaf	69.9	418	5.2	-	-	-	-
fruit	84.7	213	0.8	0.12	265	1.0	0.2

Nuts, seeds, herbs and other foods

English: Almond

Local:

Scientific name: *Prunus dulcis*

Plant family: ROSACEAE

Description: A deciduous tree. It grows 7.5-12 m high. It can spread 7.5-9 m wide. It grows straight then spreads later. The bark is dark grey and cracks. The bark has yellow rings of pores around the trunk. The leaves are alternate and long pointed. They are 12 cm long by 4 cm wide. They hang down. The leaves have fine soft teeth around the edge. The leaves are folded along the midrib. It loses its leaves during the year. The flowers appear before the leaves. The flowers occur either singly or in pairs. They are pink and 5 cm across. The fruit are green and contain edible nuts. The fruit are 6 cm long. The flesh is dry and leathery. There is one stone and the white seed inside is edible.



Distribution: It is native to North Africa, Central and South-West Asia. They are frost hardy and suit a Mediterranean climate. They need a warm summer to fully ripen. It suits hardiness zones 7-10.

Use: The oil from the kernel can be bitter and contains HCN which needs to be removed by heating. The oil is used as food flavouring. The kernels of sweet kinds are eaten. They can be eaten fresh or dried, crushed, flaked, ground and used as ingredients. The nuts can be blended with water to form almond milk, made into almond butter or pressed for their edible oil. The oil is used in salads.

Cultivation: Almond trees cannot pollinate themselves. For good fruit set pollinating cultivars are planted along with others.

Production: Trees can bear for 50 years. A tree can produce 10-15 kg per year.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
nut	5.3	2418	21.3	1	-	4.3	3.4

Nuts, seeds, herbs and other foods

English: Sesame

Local:

Scientific name: *Sesamum indicum*

Plant family: PEDALIACEAE

Description: A small, erect annual plant. It is very branched and grows 1-2 m tall. The stem is stout, 4 sided and furrowed along its length. It is densely covered with fine, downy, glandular hairs that vary in shape. The lower leaves have long stalks and are spear shaped, often with lobes or a toothed edge. The leaf stalks are 3-11 cm long. The leaf blade is 4-20 cm long by 2-10 cm wide. Upper leaves are narrow and oblong. They are 0.5-2.5 cm wide. The flowers occur in the axils of upper leaves, either on their own, or



in groups of 2 or 3. They can be white, pink, purplish and with yellow spots and stripes. The fruit can be smooth or rough and there are 2 chambers in the capsule. The fruit are brown or purple. They are oblong and deeply grooved. The seeds are small and oval. They are 3 mm by 1.5 mm and vary in colour from white, yellow, grey, red, brown or black. The fully ripe pods burst open.

Distribution: A tropical plant that suits the hot, dry, semi-arid tropics and sub-tropics. It can tolerate short periods of drought once established. It needs a temperature of 20-24°C in early growth, then 27°C for ripening. It grows from sea level to about 1200 m in areas with an annual rainfall of 400-1000 mm. Soils need to be well drained. It is very intolerant of water-logging. It cannot stand high humidity and needs frost free conditions. It needs a dry period for seed drying. It does not like acid soils. It grows in open sunny places. It can grow in arid places.

Use: The seeds are eaten. They are used in soups or fried or boiled. They are used in tahini and hummus. Seeds are eaten in the form of sweetmeats. Roasted seeds are used in pickles. They are also put on bread. Oil from the seeds is used in cooking and on salads. The refuse from the seed after the oil has been extracted is boiled in water and made into soup.

Cultivation: Plants are grown from seed. Seed will not germinate below 21°C. Seeds are broadcast on well prepared land and then harrowed in using a light harrow, or sown 2-15 cm apart in rows 20-45 cm apart. Plants can be thinned or weeded during early growth to produce a better crop. Seeding rates of 9-11 kg/ha are used. Some varieties shatter easily.

Production: Yields of 340-500 kg/ha are average. Plants reach maturity in 80-180 days. Crops are harvested as the leaves begin to drop. Plants are cut and stooked or dried in racks. The hull is removed by soaking in water overnight, then partly dried and rubbed against a rough surface.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	4.7	2397	17.7	1	-	14.6	7.8
leaf (raw)	85.5	188	3.4	-	-	-	-
oil	-	3683	0.2	-	-	-	-

Nuts, seeds, herbs and other foods

English: Sunflower

Local:

Scientific name: *Helianthus annuus*

Plant family: ASTERACEAE

Description: An upright annual plant that ranges in height from 1-4 m. It has a strong tap root. Plants are mostly unbranched, but may have some branches. The stems are hairy. The leaves are large and oval to heart shaped with teeth around the edges. They are roughly hairy and mid to dark green. Leaves can be 10-40 cm long by 5-20 cm wide. The leaf stalk is long. The flowers are yellow and daisy like, and 9-20 cm across. Sometimes they are tinged red or purple.



Distribution: A temperate plant that suits the highlands of the tropics and can stand a light frost. It needs a well drained, rich soil. It is drought and frost resistant. Sunflower grow from the equator to 55°N latitude. It does not suit the wet tropics. It cannot tolerate very acid soils. It can grow in arid places. It suits hardiness zones 4-11. It is widely distributed in many environments.

Use: An edible oil is extracted from the seeds and used for cooking. Sometimes seeds are eaten raw or roasted. The seeds can be ground into a meal for use in bread and cakes. They are also dried, roasted and ground and used as a coffee substitute. The seeds are boiled with water and honey to make a drink. The germinated seeds are fermented into a yogurt or cheese.

Cultivation: Plants are grown from seed. Only well-filled seed should be planted. It is easy to save your own seed as dry seed stores well. A plant spacing of 1 m by 0.5 m is suitable. Seeds are sown at a depth of 2-4 cm. Mature heads are collected by hand, dried and then threshed.

Production: Time to maturity is usually 4-5 months. Seeds are ready to eat when the flower starts to wither.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	5.4	2385	22.8	5	1.4	6.8	5.1

Nuts, seeds, herbs and other foods

English: Peanut

Local:

Scientific name: *Arachis hypogea*

Plant family: FABACEAE

Description: Peanuts grow on spreading bushy plants up to about 40 cm high. The leaves are made up of 2 pairs of oppositely arranged leaflets. Flowers are produced in the axils of the leaves. Two main kinds of peanuts occur. The runner kind (Virginia peanut) has a vegetative or leafy branch between each fruiting branch and therefore produces a spreading bush. The bunch type (Spanish-Valencia peanuts) produces fruiting branches in a sequence one after the other along the branches. They grow as a more upright plant and grow more quickly. Pods are produced on long stalks which extend under the ground and they contain between 2-6 seeds. The stalk or peg from the flower grows down into the soil and then produces the pod and seed under the ground. The flowers need to be no more than 18 cm from the soil surface for the seed pod to develop underground.



Distribution: Peanuts grow in tropical and subtropical areas. They grow well from sea level up to about 1650 metres in the equatorial tropics. They require temperatures of 24-33°C. Plants are killed by frost. They need a well-drained soil and cannot stand water-logging and often require raised garden beds. Peanuts need 300-500 mm of rain during the growing season. Dry weather is needed near harvest.

Use: The seeds can be eaten raw, cooked or sprouted. They are boiled, steamed, roasted, salted or made into peanut butter or flour. The young leaves and unripe pods are edible after cooking. An edible oil is extracted from the seeds. The remaining meal can also be eaten.

Cultivation: Peanuts require soil with good levels of calcium and boron or they produce empty pods. Peanuts have nitrogen fixing root nodule bacteria and therefore can give good yields in soils where nitrogen is low. The nuts are normally removed from the shell before planting and are sown 2-3 cm deep, with 10 cm between plants and 60-80 cm between rows. The soil needs to be weeded and loose by the time the flowers are produced to allow the peg for the seed pods to penetrate the soil.

Production: Flowering can commence in 30 days and it takes 3.5-5 months until maturity. Peanuts are harvested by pulling out the plant when the top of the plants die down. After harvesting, they should be left to dry in the sun for 3-4 days. Virginia peanuts have a longer growing season and the seeds need to be stored for 30 days before they will start to re-grow.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed (dry)	4.5	2364	24.3	-	-	2.0	3.0
seed (fresh)	45	1394	15	-	10	1.5	-
leaf	78.5	228	4.4	-	-	4.2	-

Nuts, seeds, herbs and other foods

English: Turmeric

Local:

Scientific name: *Curcuma longa*

Plant family: ZINGIBERACEAE

Description: A herb in the ginger family which continues growing from year to year. For harvesting, it is grown as an annual crop. It grows up to 1 m high and spreads to 50 cm across. The stem is erect. It has a bright orange, sweet smelling, fattened stem (or rhizome) under the ground. The main fattened part is about 5-8 cm long by 1-5 cm across. It has thinner branches off it. These are thickened with secondary tubers. The leaves are in a cluster and have leaf stalks that clasp the stem. The leaves are sword-shaped and have long leaf stalks. Leaves are long, flat and bright green. The flowers are dull yellow. They occur in clusters of 3-5. The cultivated turmeric is normally a sterile triploid (three sets of chromosomes) plant that does not bear fruit.



Distribution: It is a tropical plant and is widespread in coastal areas up to 1000 m altitude in the Philippines. It needs a hot, moist climate and a well-drained soil. The soil needs to be loose for good rhizomes. Plants can grow in the shade but the yield is lower. Plants are drought and frost tender. It suits hardiness zones 10-11.

Use: The orange-coloured rhizome is used as seasoning, especially in curry. It is also used for food colouring and can be used fresh or boiled and dried. The young shoots are also eaten. It is a minor spice.

Cultivation: Plants are normally grown from pieces of the rhizome. Pieces with one or two buds are used. Larger side tubers are best. They are planted 6-8 cm deep. The distance between plants needs to be 30-40 cm. They can also be grown from seed where these occur. The roots are harvested as the leaves start to die back.

Production: Leaves appear above ground in about 4 weeks. The crop is harvested in about 9-10 months when the lower leaves turn yellow. When the roots are harvested, they are put in boiling water for an hour. This assists drying and avoids sprouting. They are then dried in the sun. The outer layer is removed by rubbing.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
rhizome (dry)	11.4	1481	7.8	0	25.9	41.4	4.4
root (fresh)	88.2	192	1.2	-	-	2.3	-

Nuts, seeds, herbs and other foods

English: Teff

Local:

Scientific name: *Eragrostis tef*

Plant family: POACEAE

Description: It is an annual tufted millet grass that grows 60-120 cm tall. It has a slender stem and long narrow smooth leaves. The flowers occur in loose open panicles 15-35 cm long. The branches are very thin and droop over. The seeds are very small (1-1.5 mm long). There are about 2500-3000 seeds per gram. Brown and white-seeded types are recognised.



Distribution: A drought resistant tropical grass. It grows in places with a distinct dry season. It grows best at about 2000 m altitude in Ethiopia in temperatures of 25-28°C. It is grown from 1700-2800 m. Brown teff is grown at the higher locations. The rainfall in this region is about 950-1000 mm. It can be grown with rainfall of 400 mm. Soils should be permeable. It can tolerate frost and can grow in arid places.

Use: Seeds are ground into flour and cooked in a variety of ways. It can be used in stews or to make unleavened bread. This is called *injera* in Ethiopia.

Cultivation: Teff is best grown in fallowed land or after legume crops. Land preparation needs to be very thorough. A fine firm weed-free seed bed is needed. Seed are mostly broadcast. Driving sheep or cattle over the land is used to trample in the seed. Seed is sown at 25-30 kg per hectare. Nitrogen fertiliser is recommended. It is usually harvested with sickles.

Production: It is fast growing. Plants take 90-120 days for early varieties and 120-160 days for late maturing varieties. Yields range between 300 kg and 3000 kg per hectare. Seeds can be stored for many years as a reserve food supply.

Food Value: Per 100 g edible portion

Edible part	Moisture %	Energy kJ	Protein g	proVit A µg	proVit C mg	Iron mg	Zinc mg
seed	9.3	1541	8.9	-	-	9.9	20

Nutritional values of food plants by plant Family

Plant Family	Scientific name	Common name	Edible part	Moisture %	Energy kJ	Protein g	Vit A µg	Vit C mg	Iron mg	Zinc mg	Page
AMARANTHACEAE	<i>Amaranthus spinosus</i>	Prickly amaranth	leaf	91.7	84	3.6	109	46	14.4	0.3	38
ANACARDIACEAE	<i>Mangifera indica</i>	Mango	fruit	83.0	253	0.5	54	30	0.5	0.04	41
ANACARDIACEAE	<i>Anacardium occidentale</i>	Cashew	nut	4.0	2478	17.5	-	-	2.8	4.8	59
ARACEAE	<i>Colocasia esculenta</i>	Taro	root	66.8	1231	1.96	3	5	0.68	3.2	11
ASTERACEAE	<i>Helianthus annuus</i>	Sunflower	seed	5.4	2385	22.8	5	1.4	6.8	5.1	62
BOMBACACEAE	<i>Adansonia digitate</i>	Boabab	fruit	16.0	1212	2.2	-	360	7.4	6.7	39
CONVOLVULACEAE	<i>Ipomoea batatas</i>	Sweet potato	tuber (baked)	72.9	431	1.7	961	24.6	0.5	0.3	13
CONVOLVULACEAE	<i>Ipomoea obscura</i>	Yellow ipomoea	leaf	56.6	569	8.8	-	-	28.8	1.1	35
CUCURBITACEAE	<i>Momordica charantia</i>	Bitter cucumber	pod (boiled)	94.0	79	0.8	11	33	0.4	0.8	56
CUCURBITACEAE	<i>Luffa cylindrica</i>	Smooth loofah	fruit	94.3	79	1.1	-	-	0.7	-	34
CUCURBITACEAE	<i>Cucurbita maxima</i>	Pumpkin	leaf	88.0	160	4.9	260	28	2.5	0.9	52
CUCURBITACEAE	<i>Cucurbita pepo</i>	Marrow	fruit (mature)	92.0	105	1.6	17	16	2.4	-	54
DIOSCOREACEAE	<i>Dioscorea alata</i>	Greater yam	tuber	76.6	323	2.0	18	10	0.8	0.39	18
DIOSCOREACEAE	<i>Dioscorea bulbifera</i>	Potato yam	tuber	70.8	357	2.7	-	78	3.1	0.4	20
EUPHORBIACEAE	<i>Manihot esculenta</i>	Cassava	tuber	62.8	625	1.4	30	15	0.23	0.48	15
FABACEAE	<i>Glycine max</i>	Soybean	seed	9.0	1701	33.7	55	-	6.1	-	23
FABACEAE	<i>Parkia filicoidea</i>	African locust bean	seed (dry)	7.0	1780	32.3	-	6	33.2	-	24
FABACEAE	<i>Vigna unguiculata</i>	Cowpea	seed (dry)	11.2	1189	23.5	-	1.5	6.4	-	25
FABACEAE	<i>Lablab purpureus</i>	Lablab bean	seed (young)	86.9	209	3.0	14	5.1	0.8	0.4	26
FABACEAE	<i>Vigna mungo</i>	Mung bean	seed (raw)	12	981	22.0	6	-	8.0	-	27
FABACEAE	<i>Canavalia ensiformis</i>	Jack bean	seed	10.0	1423	20.4	160	-	4.9	-	28
FABACEAE	<i>Cajanus cajan</i>	Pigeon pea	pod (young)	64.4	477	8.7	-	-	2.0	-	29
FABACEAE	<i>Sphenostylis stenocarpa</i>	African yam bean	seed	9.0	1470	19.2	-	-	-	-	30
FABACEAE	<i>Crotalaria juncea</i>	Sunhemp	leaf	10.2	-	30.1	-	1.39	28.9	-	33
FABACEAE	<i>Lens culinaris</i>	Lentils	split & boiled	72.1	420	7.6	20	-	2.4	1.0	50
FABACEAE	<i>Arachis hypogea</i>	Peanut	seed (dry)	4.5	2364	24.3	-	-	2.0	3.0	63
LAMIACEAE	<i>Plectranthus rotundifolius</i>	Country potato	tuber	76	393	1.4	-	1.0	6.0	-	17
MALVACEAE	<i>Sida cordifolia</i>	Goat's horns	leaf	6.6	1296	24.2	-	-	79.8	-	31
MALVACEAE	<i>Hibiscus rosa-sinensis</i>	Hibiscus	leaf (dry)	6.4	1339	25.9	-	-	19.6	8.9	36
MALVACEAE	<i>Grewia tenax</i>	Small-leaved white raisin	fruit	59.1	-	4.5	-	161	125	-	46
MALVACEAE	<i>Abelmoschus esculentus</i>	Okra	fruit (cooked)	90.0	134	1.9	58	16.3	0.5	0.6	51
MALVACEAE	<i>Corchorus olitorius</i>	Jute	leaf (cooked)	87.2	155	3.4	156	33.0	3.1	0.8	53
MORINGACEAE	<i>Moringa oleifera</i>	Horseradish tree	leaf (boiled)	87	189	4.7	40	31.0	2.0	0.2	55
OPILIACEAE	<i>Opilia amentacea</i>	Catkin blooming	leaf	9.2	-	14.8	-	3.9	15.7	3.2	37
PEDALIACEAE	<i>Sesamum indicum</i>	Sesame	seed (dry)	4.7	2397	17.7	1	-	14.6	7.8	61
PHYLLANTHACEAE	<i>Phyllanthus maderaspatensis</i>	Quail grass	leaf	60.9	552	3.0	-	-	59.4	1.0	32
POACEAE	<i>Sorghum bicolor</i>	Sorghum	seed	-	1459	11.1	-	-	-	-	21
POACEAE	<i>Eleusine coracana</i>	Finger millet	seed	11.7	1594	6.2	-	-	5.3	-	22
POACEAE	<i>Pennisetum glaucum</i>	Bullrush millet	seed	11.6	1442	10.5	-	-	6.5	1.7	49
POACEAE	<i>Eragrostis tef</i>	Teff	seed	9.3	1541	8.9	-	-	9.9	20	65
ROSACEAE	<i>Prunus armeniaca</i>	Apricot	fruit (raw)	86	117	0.6	96	7	0.4	0.1	40
ROSACEAE	<i>Prunus dulcis</i>	Almond	nut	5.3	2418	21.3	1	-	4.3	3.4	60
RUBIACEAE	<i>Vangueria infausta</i>	Small wild medlar	fruit	64.4	498	1.4	-	11.5	24.4	7.1	48

Plant Family	Scientific name	Common name	Edible part	Moisture %	Energy kJ	Protein g	Vit A µg	Vit C mg	Iron mg	Zinc mg	Page
RUTACEAE	<i>Citrus reticulata</i>	Mandarin	fruit	87.6	184	1.5	42	136	0.8	-	47
SAPOTACEAE	<i>Englerophytum magalismontanum</i>	Milk plum	fruit	74.3	404	0.9	-	13.1	0.03	-	58
SOLANACEAE	<i>Physalis peruviana</i>	Cape gooseberry	fruit (mature)	84.2	201	2.0	36	30	1.5	-	43
SOLANACEAE	<i>Cyphomandra betacea</i>	Tamarillo	fruit	86.2	113	2.0	500	28	0.7	-	44
ZINGIBERACEAE	<i>Curcuma longa</i>	Turmeric	rhizome (dry)	11.4	1481	7.8	0	25.9	41.4	4.4	64



FOOD PLANT SOLUTIONS
ROTARY ACTION GROUP
Solutions to Malnutrition and Food Security

www.foodplantsolutions.org